

The effect of different treatment durations of clopidogrel in patients with non-ST-segment elevation acute coronary syndromes: a systematic review and value of information analysis

W Rogowski,¹ J Burch,^{2*} S Palmer,³
C Craigs,² S Golder² and N Woolacott²

¹Helmholtz Zentrum München, German Research Center for Environmental Health, Institute of Health Economics and Health Care Management, Neuherberg

²Centre for Reviews and Dissemination, University of York

³Health Economics Centre for Health Economics, University of York

*Corresponding author




Executive summary

Health Technology Assessment 2009; Vol. 13: No. 31

DOI: 10.3310/hta13310

Health Technology Assessment
NIHR HTA programme
www.hta.ac.uk





Executive summary

Background

Acute coronary syndrome (ACS) is a fissuring or rupturing of atheromatous plaques leading to occlusive thrombi in the arteries. Non-ST-elevation-ACS (NSTEMI-ACS) can be classified as unstable angina with undetectable markers but with electrocardiogram changes, or non-ST-elevation myocardial infarction (NSTEMI) where there is evidence of myocardial necrosis. Sixteen-year survival rates for men aged 50–59 years are 34% with a history of myocardial infarction (MI) and 53% with a history of angina, compared with 72% of those with no history of coronary disease. For patients with confirmed NSTEMI-ACS, UK guidelines recommend early treatment with antiplatelets, which are effective in preventing ischaemic vascular events in patients at increased risk. Guidance by the National Institute for Health and Clinical Excellence (NICE) in 2004 was based in part on a Technology Assessment Report undertaken by the Centre for Reviews and Dissemination (CRD) and the Centre for Health Economics (CHE), and published as a Health Technology Assessment (HTA) report (Main *et al.*, 2004). The report presented the results of a systematic review assessing the clinical effectiveness and cost-effectiveness of clopidogrel in combination with aspirin for people with NSTEMI-ACS. Only one relevant trial was identified for inclusion in the systematic review [the Clopidogrel in Unstable angina to prevent Recurrent Events (CURE) trial]. For patients with NSTEMI-ACS at moderate to high risk of ischaemic events treated with clopidogrel, the NICE guidance recommended that it be given in combination with aspirin.

Objectives

The objective of this research project was to update the previous model, and formally assess the potential value and feasibility of further research to address the optimal duration of clopidogrel treatment using value of information (VOI) analysis and a Bayesian decision theoretic approach. In line with this we aimed to update the previous systematic review of the use of clopidogrel in combination with aspirin for patients with NSTEMI-ACS, investigating the optimal duration of treatment and effects of withdrawal from treatment.

Methods

We conducted a systematic review of the clinical effectiveness and cost-effectiveness literature. Ten electronic databases and internet resources were searched from 2003 to February 2007, including MEDLINE, MEDLINE In-Process, EMBASE, BIOSIS, CENTRAL and CINAHL. Randomised controlled trials (RCTs) of clopidogrel plus aspirin compared with aspirin alone were used to evaluate clinical effectiveness and safety. Inclusion criteria were broadened to include any comparator trial for duration of treatment studies, and any study design conducted in patients with NSTEMI-ACS, percutaneous coronary intervention (PCI), stroke, peripheral artery disease (PAD) or ST-elevation myocardial infarction (STEMI) for evidence of rebound (a reactivation of the condition or concentration of adverse events) on withdrawal of treatment. The primary outcomes for the evaluation of efficacy, safety and the duration of treatment were non-fatal MI, ischaemic heart disease (IHD) without MI, death and bleeding complications.

The systematic reviews were used to assist in updating the existing model in order to provide a more robust approach to evaluating the cost-effectiveness of alternative durations of clopidogrel. The previous work was also extended to include a formal assessment of the potential value of further research using VOI approaches. These approaches were applied to estimate the expected costs of decision uncertainty predicted by the model and the maximum value that can be placed on additional research aimed at reducing this uncertainty. The costs of decision uncertainty were quantified using the expected value of perfect information (EVPI). These were used to help identify the potential design and value of further research which could be undertaken in this area. Consideration was also given to the potential impact that the introduction of a generic version of clopidogrel may have on the VOI results.

Results

Two RCTs were included for the review of clinical effectiveness and safety. The only RCTs identified that evaluated different durations of

clopidogrel treatments were conducted in patients with stroke, PAD, STEMI or PCI. Two small RCTs and one uncontrolled retrospective cohort study were identified for the review of rebound after thienopyridine withdrawal in patients with medically-treated NSTEMI-ACS. When the criteria were broadened, five RCTs, two observational cohorts, nine case series and 33 case reports were identified in patients post-PCI, and two case series and two case reports were identified in patients with stroke, PAD or STEMI.

The CURE trial reported that the proportion of patients experiencing cardiovascular death, MI or stroke was lower in the clopidogrel group at 30 days [relative risk (RR) 0.79; 95% confidence interval (CI) 0.67–0.92] and from 30 days to 12 months (RR 0.82; 95% CI 0.70–0.95). Overall, clopidogrel seems to be effective in reducing adverse cardiovascular events in patients with NSTEMI-ACS at intermediate (RR 0.86; 95% CI 0.75–0.98) and high (RR 0.77; 95% CI 0.64–0.93) risk of ischaemic events, and there is evidence that clopidogrel increases the risk of bleeding when compared with aspirin in patients with intermediate risk of ischaemic events (RR 1.44; 95% CI 1.12–1.86). A post hoc analysis indicated that the treatment effect in the first 3 months may be greater than in later periods; however, this analysis comprised non-randomised comparisons. There were no direct comparisons of the effectiveness of different durations of clopidogrel treatment in patients with NSTEMI-ACS. The evidence available relating to the potential rebound effect on withdrawal of clopidogrel therapy in patients with NSTEMI-ACS was limited and provided no conclusive evidence of its presence or absence.

In terms of the cost-effectiveness of alternative durations of clopidogrel, the updated model reinforced the conclusions from the earlier analysis. That is, a policy of 12 months of clopidogrel for patients with NSTEMI-ACS appears to be cost-effective both in 'average' patients (i.e. based on the average across all patient risks considered) and in the subgroup of higher-risk patients (presence of any of the following: age > 70, presence of ST depression or diabetes), compared with shorter-term durations. The incremental cost-effectiveness ratio (ICER) of 12 months' duration ranged from £13,380 to £20,661 per additional quality-adjusted life-year (QALY) across the different scenarios considered. However, for lower-risk patients (absence of any of the risk factors) treatment with clopidogrel beyond 3 months does not appear to be cost-effective. The ICER of 12 months' treatment with clopidogrel varied between

£49,436 and £58,691 per QALY. These conclusions appeared robust to alternative assumptions related to whether the relative effect of clopidogrel was assumed to remain constant over time or where the treatment effect in the first 3 months was assumed to be greater than in later periods.

Estimates of EVPI were markedly higher for the combined analysis of all patients (representing an average of the risks) and for analysis of high-risk patients alone, compared with those for lower-risk patients (ranging between £48.69 million and £108.4 million at a threshold of £30,000 per QALY). It was also acknowledged that more recent changes in routine clinical practice in the UK has shifted to the extent that the CURE trial itself (or the model presented here) may no longer be considered to be representative of current practice for groups at high risk, and as such the EVPI results for this group of patients may be overstated.

At a threshold of £20,000–£30,000 per QALY, total EVPI ranged between £3.27 million and £20.38 million in the lower-risk group. Given that a trial is unlikely to be able to report until after the entry of generic clopidogrel, equivalent EVPI estimates for this scenario ranged between £10.8 million and £11.9 million. The expected value of partial perfect information (EVPPPI) calculations demonstrated that approximately 40–45% of this value was related to the treatment effectiveness parameters for clopidogrel (i.e. those for which an RCT would be required).

Limitations and uncertainties

Our review was limited by the lack of available data. Although one additional trial was identified that provided information on the clinical effectiveness of clopidogrel in patients with NSTEMI-ACS, this trial was likely to be underpowered and reported limited results. Thus the CURE trial remains the primary source of data.

No studies directly compared different durations of clopidogrel treatment, and insufficient evidence was identified to adequately assess the clinical significance of any rebound effect after withdrawal of clopidogrel in these patients. Therefore, there is still a large degree of uncertainty surrounding both the optimal duration of clopidogrel treatment and the impact of withdrawal of clopidogrel treatment, which can only be addressed by further research.

The cost-effectiveness and VOI analyses are subject to a number of potential limitations. These relate not only to the limitations noted above pertaining

to the clinical effectiveness data, representing important assumptions and parameters of the model, but also to the uncertainty surrounding a range of other factors. Firstly, the issue of risk stratification is clearly an important consideration. However, it should be noted that the pragmatic approach to risk stratification applied in the decision model (due to limited patient numbers and information available from the epidemiological data used) dichotomised the population into two separate risk categories (higher- and lower-risk patients). This meant that consideration could not be given to a wider categorisation (i.e. including a third group to represent patients at intermediate risk). Similarly, these definitions are not directly comparable with other risk stratification approaches that have been applied elsewhere. Indeed, it should be recognised that the sample of patients included in the epidemiological data set were all hospitalised for NSTEMI-ACS and hence are likely to be more representative of patients at intermediate to high risk using conventional classifications. Thus, the interpretation of the results in low- and high-risk groups should be seen in this context. Secondly, changes in routine clinical practice (particularly for the high-risk group) may mean that the results presented here are more reliable for the lower-risk group. Finally, the results of the VOI demonstrate considerable variation in the potential value of further research. More importantly, the EVPI results present an upper bound to further research and hence do not provide both a necessary and a sufficient condition, even if the cost of trial fell below this amount. This is because a trial will resolve only a proportion of the uncertainty and, as such, the amount of uncertainty that is likely to be resolved would have to be assessed against the cost of the trial to ensure that any further research was considered an efficient use of resources.

Conclusions

- Clopidogrel combined with aspirin reduces adverse cardiovascular events in comparison with aspirin alone in patients with NSTEMI-ACS, but may increase the risk of bleeding.
- The optimal duration of clopidogrel treatment in patients with NSTEMI-ACS is uncertain and requires further research.
- There is some evidence that a rebound effect occurs following the withdrawal of thienopyridine treatment, but its clinical significance is uncertain.
- The results of the updated decision model suggest that durations of clopidogrel treatment

beyond 3 months do not appear to be cost-effective in patients at lower risk. However, for an average-risk patient (and in higher-risk patients), 12 months of treatment with clopidogrel appear to be cost-effective.

- These conclusions appeared robust to alternative assumptions related to whether the treatment effect remained constant over a 12-month period or was assumed to decline after 3 months.
- There is considerable variation in the costs of uncertainty surrounding the different scenarios and populations considered. The validity of these may also be less reliable in the higher-risk groups owing to changes in clinical practice. The results in the lower-risk group suggested that the upper bound of the value of a future trial was between £10.8 million and £11.9 million (and of this total, approximately 40–45% related to parameters for which a randomised design would be essential).

Recommendations for research

An adequately powered, well-conducted RCT that directly compares different durations of clopidogrel treatment in patients with NSTEMI-ACS would ideally be required to provide more robust evidence in relation to the impact of clopidogrel withdrawal. The use of an RCT would minimise possible biases associated with establishing causality with any potential rebound effect and providing robust estimates of the relative effect of alternative durations of treatment. However, the design and cost of this trial need to be evaluated carefully in relation to the VOI estimates reported here and against other uses of NHS resources. In lower-risk groups, for which shorter durations of clopidogrel appear more cost-effective, it would seem unlikely that an adequately powered RCT would be considered to provide value for money owing to the significant cost that would be required to undertake such a study and the cost of the uncertainty that such a trial might resolve.

Publication

Rogowski W, Burch J, Palmer S, Craigs C, Golder S, Woolacott N. The effect of different treatment durations of clopidogrel in patients with non-ST-segment elevation acute coronary syndromes: a systematic review and value of information analysis. *Health Technol Assess* 2009;**13**(31).



How to obtain copies of this and other HTA programme reports

An electronic version of this publication, in Adobe Acrobat format, is available for downloading free of charge for personal use from the HTA website (www.hta.ac.uk). A fully searchable CD-ROM is also available (see below).

Printed copies of HTA monographs cost £20 each (post and packing free in the UK) to both public and private sector purchasers from our Despatch Agents.

Non-UK purchasers will have to pay a small fee for post and packing. For European countries the cost is £2 per monograph and for the rest of the world £3 per monograph.

You can order HTA monographs from our Despatch Agents:

- fax (with **credit card** or **official purchase order**)
- post (with **credit card** or **official purchase order** or **cheque**)
- phone during office hours (**credit card** only).

Additionally the HTA website allows you **either** to pay securely by credit card **or** to print out your order and then post or fax it.

Contact details are as follows:

HTA Despatch
c/o Direct Mail Works Ltd
4 Oakwood Business Centre
Downley, HAVANT PO9 2NP, UK

Email: orders@hta.ac.uk
Tel: 02392 492 000
Fax: 02392 478 555
Fax from outside the UK: +44 2392 478 555

NHS libraries can subscribe free of charge. Public libraries can subscribe at a very reduced cost of £100 for each volume (normally comprising 30–40 titles). The commercial subscription rate is £300 per volume. Please see our website for details. Subscriptions can be purchased only for the current or forthcoming volume.

Payment methods

Paying by cheque

If you pay by cheque, the cheque must be in **pounds sterling**, made payable to *Direct Mail Works Ltd* and drawn on a bank with a UK address.

Paying by credit card

The following cards are accepted by phone, fax, post or via the website ordering pages: Delta, Eurocard, Mastercard, Solo, Switch and Visa. We advise against sending credit card details in a plain email.

Paying by official purchase order

You can post or fax these, but they must be from public bodies (i.e. NHS or universities) within the UK. We cannot at present accept purchase orders from commercial companies or from outside the UK.

How do I get a copy of HTA on CD?

Please use the form on the HTA website (www.hta.ac.uk/htacd.htm). Or contact Direct Mail Works (see contact details above) by email, post, fax or phone. *HTA on CD* is currently free of charge worldwide.

The website also provides information about the HTA programme and lists the membership of the various committees.

NIHR Health Technology Assessment programme

The Health Technology Assessment (HTA) programme, part of the National Institute for Health Research (NIHR), was set up in 1993. It produces high-quality research information on the effectiveness, costs and broader impact of health technologies for those who use, manage and provide care in the NHS. 'Health technologies' are broadly defined as all interventions used to promote health, prevent and treat disease, and improve rehabilitation and long-term care.

The research findings from the HTA programme directly influence decision-making bodies such as the National Institute for Health and Clinical Excellence (NICE) and the National Screening Committee (NSC). HTA findings also help to improve the quality of clinical practice in the NHS indirectly in that they form a key component of the 'National Knowledge Service'.

The HTA programme is needs led in that it fills gaps in the evidence needed by the NHS. There are three routes to the start of projects.

First is the commissioned route. Suggestions for research are actively sought from people working in the NHS, from the public and consumer groups and from professional bodies such as royal colleges and NHS trusts. These suggestions are carefully prioritised by panels of independent experts (including NHS service users). The HTA programme then commissions the research by competitive tender.

Second, the HTA programme provides grants for clinical trials for researchers who identify research questions. These are assessed for importance to patients and the NHS, and scientific rigour.

Third, through its Technology Assessment Report (TAR) call-off contract, the HTA programme commissions bespoke reports, principally for NICE, but also for other policy-makers. TARs bring together evidence on the value of specific technologies.

Some HTA research projects, including TARs, may take only months, others need several years. They can cost from as little as £40,000 to over £1 million, and may involve synthesising existing evidence, undertaking a trial, or other research collecting new data to answer a research problem.

The final reports from HTA projects are peer reviewed by a number of independent expert referees before publication in the widely read journal series *Health Technology Assessment*.

Criteria for inclusion in the HTA journal series

Reports are published in the HTA journal series if (1) they have resulted from work for the HTA programme, and (2) they are of a sufficiently high scientific quality as assessed by the referees and editors.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search, appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

The research reported in this issue of the journal was commissioned and funded by the HTA programme on behalf of NICE as project number 07/19/01. The protocol was agreed in April 2007. The assessment report began editorial review in May 2008 and was accepted for publication in December 2008. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The HTA editors and publisher have tried to ensure the accuracy of the authors' report and would like to thank the referees for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this report.

The views expressed in this publication are those of the authors and not necessarily those of the HTA programme or the Department of Health.

Editor-in-Chief: Professor Tom Walley CBE
Series Editors: Dr Aileen Clarke, Dr Chris Hyde, Dr John Powell,
Dr Rob Riemsma and Professor Ken Stein

ISSN 1366-5278

© 2009 Queen's Printer and Controller of HMSO

This monograph may be freely reproduced for the purposes of private research and study and may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising.

Applications for commercial reproduction should be addressed to: NETSCC, Health Technology Assessment, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Published by Prepress Projects Ltd, Perth, Scotland (www.prepress-projects.co.uk), on behalf of NETSCC, HTA.

Printed on acid-free paper in the UK by Henry Ling Ltd, The Dorchester Press, Dorset.