

Pilot trial and process evaluation of a multi-level smoking prevention intervention in further education settings

Fletcher A^{1*}, Willmott M², Langford R², White J³, Poole R¹, Brown R¹, Young H¹, Moore G¹, Murphy S¹, Townson J³, Hollingworth W², Campbell R², Bonell C⁴

*Corresponding author

¹ DECIPHer, School of Social Sciences, Cardiff University, UK

² DECIPHer, School of Social and Community Medicine, University of Bristol, Bristol, UK

³ Cardiff Centre for Trials Research, School of Medicine, Cardiff University, UK

⁴ Department of Social and Environmental Health Research, London School of Hygiene and Tropical Medicine, UK

Important

A 'first look' scientific summary is created from the original author-supplied summary once the normal NIHR Journals Library peer and editorial review processes are complete. The summary has undergone full peer and editorial review as documented at NIHR Journals Library website and may undergo rewrite during the publication process. The order of authors was correct at editorial sign-off stage.

A final version (which has undergone a rigorous copy-edit and proofreading) will publish as part of a fuller account of the research in a forthcoming issue of the Public Health Research journal.

Any queries about this 'first look' version of the scientific summary should be addressed to the NIHR Journals Library Editorial Office – journals.library@nihr.ac.uk

The research reported in this 'first look' scientific summary was funded by the PHR programme as project number 13/42/02. For more information visit <https://www.journalslibrary.nihr.ac.uk/programmes/phr/134202/#/>

The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The PHR editors have tried to ensure the accuracy of the authors' work and would like to thank the reviewers for their constructive comments however; they do not accept liability for damages or losses arising from material published in this scientific summary.

This 'first look' scientific summary presents independent research funded by the National Institute for Health Research (NIHR). The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, NETSCC, the PHR programme or the Department of Health. If there are verbatim quotations included in this publication the views and opinions expressed by the interviewees are those of the interviewees and do not necessarily reflect those of the authors, those of the NHS, the NIHR, NETSCC, the PHR programme or the Department of Health.

© Queen's Printer and Controller of HMSO 2017. This work was produced by Fletcher *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health. This 'first look' scientific summary may be freely reproduced for the purposes of private research and study and extracts may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Scientific summary

Background

Smoking is a major cause of preventable illness, premature death and health inequalities in the UK. Preventing young people from taking up smoking is vital to maintain and accelerate recent declines in smoking rates. Although much research has been undertaken to develop and evaluate school-based prevention interventions targeting 11-15 year-olds, smoking continues to grow rapidly amongst older youth. With over 1.5 million British 16-18 year olds now enrolled in further education (FE) courses, new smoking prevention interventions are required that target FE settings (e.g. general FE colleges, 'sixth form' colleges attached to secondary schools, etc.). As well as being a period in the life-course when smoking often begins, the transition to FE itself can increase the risk of smoking as young people are exposed to new sources of peer influence and have more independence from their parents. However, research evidence about preventing smoking among FE students is sparse with few evaluations of smoking prevention interventions in FE colleges to date.

To address this gap, "The Filter FE" intervention and logic model was co-designed by ASH Wales and the research team to apply the educational, training and social media resources from ASH Wales' *The Filter* youth project to FE settings in 2014-15. The Filter FE is a novel, multi-level intervention targeting 16, 17 and 18 year-old students in FE settings, delivered by trained staff working on ASH Wales' *The Filter* youth project. Informed by systematic reviews of smoking prevention interventions delivered in schools and other settings, the intervention was designed to integrate the following prevention methods and approaches in FE settings: preventing the sale of tobacco to under 18 year-olds in local shops; implementing tobacco-free campus policies; training FE staff to deliver smoke free messages and support institutional change; publicising *The Filter* youth project's online social marketing campaigns, advice and support services; and on-site youth work activities to provide credible educational messages, address norms, and promote resistance skills, as well as signposting to cessation services. To facilitate scale-ability and sustainability across UK FE settings (including large institutions), the intervention involves standardised processes and activities balanced with opportunities for a local tailoring of activities.

© Queen's Printer and Controller of HMSO 2017. This work was produced by Fletcher *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health. This 'first look' scientific summary may be freely reproduced for the purposes of private research and study and extracts may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Study aim, objectives and research questions

The aim of the pilot trial was to evaluate the feasibility and acceptability of implementing and trialling the Filter FE intervention. The study had three objectives.

The first objective was to assess whether pre-specified feasibility and acceptability criteria were met prior to progressing to a larger, phase III trial to examine effectiveness. To meet this objective, data were collected and analysed to address these research questions (RQ):

- RQ1. Did the intervention activities occur as planned in (at least) two out of three intervention settings?;
- RQ2. Were the intervention activities delivered with high fidelity across all settings?;
- RQ3. Was the intervention acceptable to the majority of FE managers, staff, students and the intervention delivery team?;
- RQ4. Was randomisation acceptable to FE managers?;
- RQ5. Did (at least) two out of three colleges from each of the intervention and control arms continue to participate in the study at 1-year follow-up?;
- RQ6. Do student survey response rates suggest that we could recruit and retain at least 70% of new students in both arms in a subsequent effectiveness trial?

The second objective was to explore the experiences of FE students, staff and the intervention team regarding the pilot intervention and trial design, including how the logic model, intervention content and data collection methods could be refined. In order to meet this objective data were collected and analysed to address the following RQ:

- RQ7. What are students', staff and intervention team members' experiences of the intervention and views about its potential impacts on health?;
- RQ8. What are the barriers and facilitators to implementation and how do these vary according to college context and/or other factors?;
- RQ9. Were there any unexpected consequences?;
- RQ10. How acceptable were the data collection methods to students and staff and do participants think longer term follow-up via email or phone interview would be feasible?;
- RQ11. What resources and partnerships are necessary for a phase III trial?

The third objective was to pilot primary, secondary and intermediate outcome measures and economic evaluation methods prior to a potential effectiveness trial. It was not an objective of the pilot study to assess intervention effects and the study was not designed or powered to do so. Data were collected and analysed to address the following RQ:

- RQ12. Does the primary outcome measure (smoking weekly or more) have an acceptable completion rate, adequate validity and minimise floor/ceiling effects?;
- RQ13. Do cotinine concentrations of saliva samples indicate any evidence of response bias between arms in self-reported smoking status?;
- RQ14. Was it feasible and acceptable to measure all the secondary and intermediate outcomes of interest at baseline and follow-up?;
- RQ15. Is it feasible to assess cost effectiveness using a cost utility analysis within a phase III trial?

Methods

A cluster randomised controlled pilot trial and process evaluation was undertaken in six FE settings in Wales (purposely sampled to examine delivery and trial methods in a range of institutional contexts) with allocation to: the Filter FE intervention (three FE settings) or continuation of normal practice (three FE settings). The following criteria were used to purposely sample FE settings and stratify the allocation: large FE college campuses (new intake more than 500) (n=2); small FE college campuses (new intake fewer than 500) (n=2) 'sixth form' colleges attached to schools (n=2).

In order to assess the feasibility and acceptability of delivering and trialling the intervention according to pre-specified criteria (objective 1), we collected a range of quantitative and qualitative data via: semi-structured observations of staff training sessions (n=1 per intervention setting), group-based youth work sessions (n=1 per intervention setting) and college websites and social media channels (n=2 per intervention setting); interviews with FE college managers (n=5) and the intervention team (n=6); and documentary evidence (e.g. college policies, intervention team records, etc.). The retention of FE settings and response rates were assessed using student survey data.

To explore participants' experiences of implementing and trialling the Filter FE intervention (objective 2), qualitative process data were collected via: interviews with FE college managers (n=5) and the intervention team (n=6); focus groups with students (n=11) and staff (n=5), and semi-structured observations of intervention settings. These qualitative data were transcribed verbatim and analysed using techniques associated with thematic content analysis and grounded theory. The coding framework included both deductive codes, derived from key research questions and relevant progression criteria, and inductive codes, identifying other relevant themes emerging from the data

In addition to examining intervention and trial feasibility and acceptability, primary, secondary, intermediate (process) outcomes and economic evaluation methods were piloted (objective 3). Surveys of new students enrolling at the participating FE settings in September 2014 (baseline) and September 2015 (one year follow-up) were used to examine the pilot primary (self-reported smoking weekly or more) and secondary outcome measures (self-reported lifetime smoking, use of cannabis in the past 30 days; frequent cannabis use, high risk alcohol use, and health-related quality of life). The following additional pilot secondary outcomes for baseline smokers were also examined: cessation; number of cigarettes/week; and, nicotine dependence. Informed by the intervention logic model, multiple sources of data were also collected at baseline and follow-up to pilot intermediate (process) outcomes at multiple levels: the restriction of the availability of tobacco in local shops was assessed via 'mystery shopper' audits; changes to the institutional environment and policies were assessed via structured observations and analysis of college policy documents; students' knowledge, norm and social/situational self-efficacy and resistance skills were assessed via the student survey. Potential economic analyses methods were assessed, including the use of EQ-5D-5L health related quality of life measure. It was not feasible to collect saliva samples from students to assess the validity of self-reported smoking status at follow-up.

Results

The intervention was not delivered in full at any of the three intervention settings, with no implementation of some community and college-level components, and low fidelity of the social media component across sites. The staff training reached a total of 28 staff and youth

© Queen's Printer and Controller of HMSO 2017. This work was produced by Fletcher *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health. This 'first look' scientific summary may be freely reproduced for the purposes of private research and study and extracts may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

work activities were attended by 190 students across the three sites (<10% of all staff and students). Lower than intended recruitment to these activities was largely the result of lack of demand from staff at intervention settings and, although those who did attend were observed to be engaged, low levels of acceptability were reported across FE sites. The intervention team reported additional challenges to recruitment due to the short lead-in time prior to implementation and high intervention-team staff turnover during the pilot study. The process evaluation also found that planned institutional policy review activities did not occur at any of the sites, with limited evidence of changes to smoking policies post-intervention. This was again associated with limited preparation time for intervention delivery as well as issues relating to management of intervention, which also impacted limited community-level activities targeting local shops.

Six colleges were randomised into the two trial arms and all were retained at the one-year follow-up. Recruitment and retention of students was challenging, despite the use of the multiple methods and incentives. In September 2014, 1,320 students out of an estimated total sample of 2,363 participated in the baseline survey. Of these 14.0% (n=185) were ineligible as they were aged under 16 or over 18 and 5 students provided no data, leaving a baseline sample of 1,130 (47.8%). Although this equates to a response rate of less than 50%, the number of potentially eligible students at baseline (n=2,363) was provided by each institution and *over-estimates* the actual number of new students aged 16-18 in that setting and therefore *underestimates the true response rate*, especially in large FE settings, due to: students enrolling in principle prior to September but not registering at the start of term, deferring or dropping out in early September; inclusion of students who study across multiple campuses but whose primary campus is not the study site; and, the inclusion of some students aged over 18 due to incomplete information at enrolment. In September 2015, 412 eligible students completed the follow-up survey (36.5% of baseline respondents; 17.4% of all potentially eligible students at baseline).

The second objective was to explore the experiences of students, staff and the intervention team. Qualitative data indicated that implementation was limited by various factors, including staff and students uncertainty about the need for, and appropriateness of, smoking prevention activities in FE settings, the management of intervention, the high turnover of

intervention team staff and the short lead-in time prior to implementation. Although support was expressed for the involvement of external health agencies in the FE setting, the majority of staff and students perceived that FE is 'too late' for smoking prevention activities, with current smokers better served by cessation activities and resistance from non-smokers to educational messages with high degrees of familiarity. Significantly, the act of intervention was itself a source of resistance, with both staff and students suggesting that such approaches contrast with institutional cultures in the FE sector aimed at promoting personal responsibility and developing autonomy in a population transitioning from, more constrained, schools. The emphasis on freedom of choice was expressed via students' right to smoke.

The third objective was to pilot primary, secondary and intermediate outcome measures and economic methods. There were low levels of missing data for all pilot primary and secondary outcomes from the student baseline surveys completed in September 2014 (n=1,130 eligible participants) and one year follow-up surveys completed in September 2015 (n=412 eligible participants). The prevalence of weekly smoking at baseline was 20.6% and 17.2% at follow-up. Of the 336 students who were not a weekly smoker at baseline, only 21 (6.3%) reported being a weekly smoker at follow-up. The trial arms were not well balanced for the indicative primary and secondary outcome measures at baseline or follow up due to the small number of clusters and heterogeneity between clusters (e.g. sixth form and community colleges). It was feasible to assess changes in intermediate (process) outcome (e.g. smoking norms/attitudes, self-efficacy, situational resistance skills, etc.) and economic measures (EQ-5D-5L, health service use) over time. At follow-up, the quantitative process outcomes identified that most students attempting to purchase tobacco were still able to do so. Only 5.1% students were aware of *The Filter* project at follow-up, although the proportion was higher in intervention group (7.1%) than the control group (2.9%).

Conclusion and recommendations

This one-year pilot study is the first reported evaluation of a universal smoking prevention intervention in an FE context to date and the first cluster RCT in FE settings in the UK. It was not feasible to implement the Filter FE intervention as planned and the methods used had low levels of acceptability among students and staff. FE settings do not appear to be a

conducive environment for smoking prevention intervention activities, although weaknesses in the management of this intervention also further hindered implementation in this pilot. A larger cluster RCT to examine the effectiveness and cost-effectiveness of this intervention is not recommended. The very low prevalence of smoking uptake suggests that further consideration is needed on whether prevention or cessation activities would be most effective in FE and other educational settings. Findings should be considered in relation to evidence on age of onset for young smokers. It was feasible to recruit, randomise and retain FE settings within a cluster RCT design. FE managers valued the opportunity to be involved in health research and accepted randomisation. However, further methodological work is recommended to improve student recruitment and retention rates if RCTs are to be conducted in this setting.