

PROTOCOL TITLE:

**Understanding barriers and outcomes of unspecified
(altruistic) kidney donation (BOUnD); a multicentre
prospective cohort study**

BOUnD

Sponsor

Jennifer Boston
Guy's & St Thomas' Foundation NHS Trust
R&D Department
16th Floor, Tower Wing
Great Maze pond
London SE1 9RT
Ext Tel: 02071887188 ext. 89811
Fax: 02071881295
Email: Jennifer.boston@gstt.nhs.uk

Funder

Name of Sponsor Organisation: NIHR
Name of Sponsor Representative:
Mr Lewis Bradley
National Institute for Health Research
Evaluation, Trials and Studies
Coordinating Centre
University of Southampton
Alpha House, Enterprise Road
Southampton SO16 7NS
023 8059 7802

Chief Investigator

Mr Nizam Mamode
Department of Renal Transplant Surgery,
6th Floor, Borough Wing, Guy's Hospital,
Great Maze Pond SE1 9RT, London
020 7188 1543
Nizam.Mamode@gstt.nhs.uk

Study Manager

Rebecca Gare
Department of Renal Transplant Surgery,
6th Floor, Borough Wing, Guy's Hospital,
Great Maze Pond SE1 9RT, London
020 71887188, ext. 52409
rebecca.gare@gstt.nhs.uk

Co-applicants

Mr Petrut Gogalniceanu

Principal Investigator

Department of Renal Transplant Surgery, 6th Floor, Borough Wing, Guy's Hospital, Great Maze Pond
SE1 9RT, London

020 7188 1543

petrut.gogalniceanu@gstt.nhs.uk

Ms Hannah Maple

Clinical Lecturer, Guy's & St Thomas's NHS Foundation Trust

6th Floor, Borough Wing

Guy's Hospital

Great Maze Pond

London SE1 9RT

United Kingdom

hannah.maple@gstt.nhs.uk

Professor Heather Draper

Professor of Biomedical Ethics - University of Birmingham

University of Birmingham h.j.a.draper@bham.ac.uk

School of Health and Population Sciences

0121 414 6941

h.draper@bham.ac.uk

Dr Sam Norton

Study Statistician- Lecturer in Research Methods & Statistics, Psychology Department, Institute of Psychiatry,
King's College London

Health Psychology, Institute of Psychiatry, Psychology & Neuroscience Kings College London, 5th floor

Bermondsey Wing, Guy's Hospital Campus, London SE1 9RT

sam.norton@kcl.ac.uk

Dr Jo Chilcot

Lecturer in Health Psychology

King's College London

5th Floor, Bermondsey Wing, Guy's Hospital, UK

020 7188 2597

joseph.chilcot@kcl.ac.uk

Ms Annie Mitchell

University of Plymouth

Clinical Director and Associate Professor - Plymouth University

Room 504, Rolle Building, Drake Circus, Plymouth, Devon, PL4 8AA

annie.mitchell@plymouth.ac.uk

01752 586657

Professor Paul McCrone

Professor in Health Economics

Department, Institute of Psychiatry, Psychology & Neuroscience (IoPPN), King's College London

paul.mccrone@kcl.ac.uk

0207 8480874

Ms Lisa Jane Burnapp

Lead Nurse- Living Donation, Organ Donation and Transplantation; Consultant Nurse Living Kidney Donation
- Guy's and St. Thomas' NHS Foundation Trust
6th Floor, Borough Wing, Guy's Hospital
Great Maze Pond, London SE1 9RT, UK
Lisa.Burnapp@nhsbt.nhs.uk
020 7188 7188

Mr Paul J Gibbs

Portsmouth Hospitals NHS Trust paul.gibbs@porthosp.nhs.uk
Consultant Renal Transplant and Vascular Surgeon - Portsmouth Hospitals NHS Trust
Queen Alexandra Hospital Southwick Hill Road Cosham, PO6 3LY
023 9228 6400

Dr Alexis Clarke

University of Plymouth
Research Fellow-Clinical Psychologist - Plymouth Community Healthcare
Rolle Building, Drake Circus, Plymouth, Devon, PL4 8AA
alexisclarke@plymouth.ac.uk

Study Synopsis

Title	Understanding barriers and outcomes of unspecified (altruistic) kidney donation (BOUnD); a multicentre prospective cohort study.
Protocol Short Title/Acronym	BOUnD (Barriers and Outcomes in Unspecified kidney Donation)
Protocol Version number and Date	RQ2 Protocol v2.1, 25/05/2016
Is the study a Pilot?	No
Study Hypothesis	(i) Regional differences in unspecified (altruistic) kidney donor rates will be explained by prospective donor experience e.g. depending on donor interaction with staff members, local expertise and resources. (ii) There is no detrimental impact of unspecified donation on mental and physical health.
Study Duration	December 2015 – April 2018
Methodology	Prospective, mixed-method cohort study recruiting unspecified potential donors (and a directed donor control group). Participants will be recruited to a prospective donor phase shortly after first enquiring about donation (hypothesis i). Those that proceed to donation will continue to a second phase focusing on outcomes over 1 year (hypothesis ii). Nested qualitative studies will explore experiences of the process in donors and non-donors using structured interviews. Focus groups will be used to guide questionnaire design and interview topic guide.
Sponsor name	Guy's and St.Thomas' NHS Foundation Trust R&D Office
Chief Investigator	Prof Nizam Mamode
REC number	15/SC/0637
Medical condition or disease under investigation	Unspecified (altruistic) living kidney donation
Purpose of study	To identify methods of improving the process of unspecified (altruistic) donation in the UK and inform the development of national guidelines
Primary objective	Physical and mental-health related quality of life, anxiety, depression, life satisfaction and self-esteem
Secondary objective (s)	1. Barriers to unspecified kidney donation 2. Economic outcomes of unspecified kidney donation
Number of Subjects/Patients	(i) 16 - 24 participants (focus groups); (ii) 1024 participants (questionnaires), as follows: Test group: 224 unspecified donors that proceed to donate and 400 unspecified donors that withdraw. Control group: 200 directed / specified donor controls that proceed to donate and 200 directed / specified potential donors who did not proceed; (iii) 45 participants (interviews),
Study Design	Prospective cohort study
Endpoints	(i) time from enquiry to donation (for those that proceed to donation) (ii) mental and physical health at 3 and 12 months post donation / withdrawal, compared to directed donor controls
Inclusion Criteria	Individuals contacting a UK transplant centre wishing to become specified or unspecified kidney donors or those that have already begun the work-up process
Exclusion Criteria	Foreign nationals that are unable to donate altruistically in their countries of residence or prisoners
Statistical Methodology and Analysis	Quantitative analysis: (i) Time-to-event analysis using Cox regression; (ii) propensity score weighted mean differences at 12 months using

Glossary of Terms and Abbreviations

AE	Adverse Event
AR	Adverse Reaction
ASR	Annual Safety Report
CA	Competent Authority
CI	Chief Investigator
CRF	Case Report Form
CRO	Contract Research Organisation
DMC	Data Monitoring Committee
EC	European Commission
GAfREC	Governance Arrangements for NHS Research Ethics Committees
ICF	Informed Consent Form
ISRCTN	International Standard Randomised Controlled Trial Number
MA	Marketing Authorisation
MS	Member State
Main REC	Main Research Ethics Committee
NHS R&D	National Health Service Research & Development
PI	Principle Investigator
QA	Quality Assurance
QC	Quality Control
Participant	An individual who takes part in a clinical trial
RCT	Randomised Controlled Trial
REC	Research Ethics Committee
SAE	Serious Adverse Event
SDV	Source Document Verification
SOP	Standard Operating Procedure
SSA	Site Specific Assessment
TMG	Trial Management Group
TSC	Trial Steering Committee

1. Introduction

Over one third of all kidney transplants taking place in the UK today are from living donors. A growing subset of living donors are individuals who choose to donate a kidney to someone that they have not previously met; so called ‘unspecified’ or ‘non-directed altruistic’ donors. Over 200 unspecified donations have taken place in the UK to date since this was introduced in 2006 and this type of living donation is becoming more routine, currently accounting for approximately 7% of living donations (1).

Despite this increase, the concept of unspecified kidney donation (UKD) remains uncomfortable for some clinicians, principally due to concerns about the motivations, characteristics and outcomes of these donors. In a recent study of clinicians’ views, 78% of French physicians were opposed to the practice of unspecified donation (2). In our previous qualitative work, we have found some evidence that this makes donation more difficult or stressful for some potential donors (3-5). Furthermore, we recently performed a large study of a national cohort of all 148 UKDs in the UK over the first five years of the programme, and compared them with a regional sample of 148 specified kidney donors (SKDs - those who donate to someone with whom they have an emotional relationship) (6). This study did not find an excess of poor psychosocial or physical outcomes in UKDs; however the response rate was 74%, with variable retrospective follow-up, and therefore it is impossible to be certain that donors with significant pathology were not missed- indeed, these are the very donors (for example, with depression) that might be expected to fail to respond. The study did highlight broad regional variations in the numbers of UKDs performed and has highlighted differences in the assessment process, which may explain the differences seen across the country. Indeed, 45% of all unspecified donations were performed in 3 centres. There is some evidence from other studies that attitudes from transplant professionals may be a barrier to donation (7-9). Both living donor nurses and psychiatric assessors involved in UKD have expressed concerns about the lack of practice guidance in this area, lack of clear guidance could be a further barrier to donation (4,5).

Through our qualitative work we have also found that barriers to donation may exist within families where there is tension over the decision to donate altruistically and there may be a role for transplant services to support families in this situation (3). We have recently been awarded a grant from the British Renal Society and British Kidney Patients Association to explore this. This work is due to commence prior to this study and will inform this research.

The UKD participants in our PPI sessions and previous qualitative study identified a number of issues in the process that they felt acted as deterrents and may have affected the decision by others to donate (3). They found difficulties in knowing how to make initial contact with the transplant centre. The negative attitudes of transplant professionals were also off-putting and this continued whilst donors were in hospital, with some experiencing ignorance and hostility from ward staff which made them feel guilty for “choosing to become a patient”. The length of the workup process was also commonly an issue, which donors found frustrating. Indeed, when considering living donor chains, most donors would have liked to have participated had it been easier and the timing more predictable. Many were working or had other commitments and the unpredictability of when the donation would take place meant that many were not in a position to oblige. The psychiatric assessment (which is no longer legally mandatory but is considered current best practice) was also a difficult experience for donors who felt that they had to prove their sanity (3).

Unspecified kidney donation is apparently more costly than specified donation, as it is resource intensive, with a large number of enquiries and assessments, and a low proportion who proceed to donate. In Portsmouth (the largest centre for unspecified donation), for example, of 149 referrals, 27 have donated and a further 27 are in work-up, giving a drop-put rate of at least 64%. Nevertheless, a kidney from a UKD may be a particularly valuable resource, since it can be used to provide a high quality, long lasting transplant to those who are otherwise difficult to transplant. The National Kidney Sharing Scheme, for example, involves kidney exchanges between pairs of donor and recipients who cannot otherwise proceed due to immunological incompatibility. A kidney from a UKD can be used to convert these exchanges into a ‘chain’ primed by the UKD; the UKD donates to recipient A, and her donor donates to recipient B, and so on

(Appendix 2). In the US this has resulted in 30 transplants occurring from a single UKD (10). In the UK, 47% of UKDs have been used to prime short chains of two transplants, and the UK Living Donor Strategy aims for 75% to be used for chains, with 3 transplants in each chain, by 2020 (11). Thus, assuming UKDs rise to 200 per year, they would result in 450 transplants annually, which is almost half the current annual living donor transplantation rate. Despite this, no economic analysis of unspecified donation has been performed. This is particularly important since, if it is shown to have a significant economic benefit, extra resources could be allocated by NHS Blood and Transplant, as happened with SKDs over the last decade.

We therefore wish to perform a comprehensive assessment of the unspecified donor programme in the UK, in order to determine the extent and reasons for variation in practice, ascertain barriers to donation, and determine the economic costs and benefits of an unspecified donation. We will also assess outcomes after unspecified donation, in order to provide detailed evidence for transplant teams' decisions about potential donors.

2 Study Objectives and Design

2.1. Study Objectives

Aims: This study aims to perform a comprehensive assessment of unspecified altruistic donor programme in the UK to explore variation between centres and identify barriers and facilitators to donation for those that have expressed a willingness to do so.

Objectives:

- (i) Identify and explain regional variations in unspecified kidney donation (UKD), based on donor interaction with staff members, local expertise and resources, and other economic variables
- (ii) Establish prospectively the psychosocial, physical and economic outcomes of individuals undertaking unspecified kidney donation, compared to specified donors.

Outcomes

Primary outcomes: Physical and mental health-related quality of life.

Psychosocial health outcomes:

- quality of life (SF-12)
- anxiety (General Anxiety Disorder-7 (GAD-7))
- depressive symptoms (Patient Health Questionnaire-9 (PHQ-9))
- life satisfaction (Satisfaction With Life Scale)
- self-esteem (Rosenberg Self-Esteem Scale)
- Decision Regret Scale
- Flourishing Scale
- in house questionnaire

Physical health outcomes

NHSBT pre and post donation physiological and clinical outcomes

Secondary outcomes:

- Barriers to donation (qualitative data from interviews and focus groups)
- Healthcare resource utilisation data (Client Service Receipt Inventory (CSRI))

2.2 Recruitment Strategy

The primary study group will comprise all those who approach a transplant team in any UK transplant centre, offering to donate a kidney to a stranger over a three year period (2015 – 2018). Follow-up will take place up to 2020.

The control study group will comprise all those who approach a transplant team across the UK offering to donate a kidney to a family member or friend (“specified donors”).

The study will use the same national professional transplantation network to collaborate with transplant co-ordinators and living donor nurses willing to participate in the recruitment process. Participant recruitment will take place subsequent to local R&D approval and transplant centres being identified and approved as participant identification centres (PIC).

UK Transplant co-ordinators will be briefed regarding the aims, objectives and recruitment criteria of the study. Communication and liaison with local transplant co-ordinators will be through Ms Lisa Burnapp (Lead Nurse - Living Donation, Organ Donation and Transplantation, NHSBT), who is a collaborator in the study.

Focus Groups Recruitment.

The Focus Groups represent the smallest aspect of the study and serve to help fine-tune the questionnaire design and interview topic guides. As such, only two focus groups will take place in centres such as Guy’s Hospital (London) or Plymouth Derriford Hospital, where the study team has long-standing collaboration links with the donation teams. The local transplant co-ordinator or living donation nurse specialist (living donation team) will approach individuals that have recently donated or have withdrawn from donation and explore whether they would be interested in considering the study. Those that would be interested will be given the contact details of our research team or asked if they would agree to be contacted by us. The research team would then be able to provide further information and lead the consent and recruitment process.


Interview and Questionnaire Recruitment

Transplant co-ordinators at each of the 23 UK transplant centres will ask potential donors approaching their units if they would be interested in participating in the study. This does not equate to being consented into the study, but simply facilitates further information gathering regarding our work. This will occur either at the initial telephone contact between the potential donor and the transplant centre or at the first clinic consultation with the transplant co-ordinator, depending on local practice protocols. Potential donors already being worked-up will also be given the opportunity to contact the study team for recruitment into the study.

Once a potential donor agrees to find out more about the study, the local transplant co-ordinator will facilitate contact with the research team by either giving the team’s contact details to the potential donor, or (with the donor’s permission) pass on their preferred contact details to the research team. The study’s manager will be notified of individuals interested in the study. The research team will contact potential participants by phone, email or post to provide further information, discuss the study and provide participant information sheets. Those that indicate a willingness to participate will be enrolled in the study subsequent to completion of the relevant consent forms. The emphasis of the study is to cause minimal inconvenience to local transplant units and human resources. As such, once a transplant co-ordinator has facilitated the contact between the potential donor and the study team, no further involvement will be needed and all subsequent administrative and research work will be co-ordinated by the study’s manager or investigators.

The control group will consist of individuals who are donating to friend or relative (specified donors). Control (specified) donors will be recruited in a similar consecutive manner by transplant co-ordinators. Control donors

that do not proceed to specified donation will still be asked to participate in the study in order to provide robust data for comparison.



2.3 Study Design

Focus Groups

Two focus groups with potential donors will be undertaken. One focus group will involve those that have proceeded to donation. The other will involve potential donors that have withdrawn or been withdrawn from the donation process. 8-12 participants will take part in each focus group. The focus groups will not involve control participants. The physical location of the focus groups will be a suitable hospital venue, such as a conference room or a postgraduate centre. Recruitment will be undertaken as described above. The focus group discussion will be audio-recorded and transcribed for future analysis.

Data regarding socio-demographic (including the area postcode), physical, psychological characteristics, and resource use variables will be collected at baseline (shortly after contacting the transplant centre).

Questionnaires


The questionnaire part of the study will have four research populations on which questionnaire data will be collected at four intervention points (Q1-Q4): baseline, preoperatively and at 3 and 12 months post-donation in the form of a study questionnaire bundle. Additional data (such as gender, age or ethnic group) will be collected at the time of recruitment into the study.

Q1. Baseline data will be collected within the first week of recruitment to assess the participant prior to the work-up process.

Q2. Pre-operative data will be collected in the 2 weeks preceding donation surgery (+/- 3 days). This will not be collected on the day of surgery in order to avoid confounding errors. This will mark the end of the work-up period. For those that withdraw from the study a longer period of time may be needed to capture these participants. In this case, the Q2 intervention point will span from the time of withdrawal to 3 weeks post withdrawal.

Q3. 3 months post donation or withdrawal

Q4. 12 months post donation or withdrawal




The four study populations will include:


1. Those that **proceed to donation** ('unspecified donors') – Test 1 population
2. Those that **do not to donate** ('unspecified non-donors') due to donor's choice or withdrawal by the clinical assessors – Test 2 population
3. Those that undergo **living donation to a known individual** ('specified donors'), which will act as **control group 1**
4. Those **intending to donate to a known individual that do not donate** ('specified non-donors'), which will act as control group 2

To ensure feasibility of the study questionnaire burden will be tested and considered in conjunction with the PPI group.

Questionnaire validation will be carried out by asking 20-30 volunteers that are previous kidney donors or future specified donors to review the in-house questionnaires. This will involve a facilitated think-aloud exercise to identify any face validity issues related to the newly developed questions. This exercise will result only in minor changes to the question structure, phrasing or answering methods. The questionnaire content validity will have already been validated by 15 members of the research team who will review the developed questionnaires on at least three occasions.



Participants who do not proceed to donation will be identified either by regularly checking with their local transplant coordinators (every two weeks) or by self-referral to the study team. The study researchers will then ascertain whether the patient self-withdrew or was withdrawn by the clinical team. The following data collection algorithm will be used:



Interviews


Qualitative interviews will also be completed with a sample of 15 donors who completed their donation, 15 who withdrew and 15 who were withdrawn by the transplant team from the process. Participants will be asked about their experience of the donation process and services, barriers and enablers to donation and outcomes from either donating or withdrawing from the process. The interview questions have been informed by our previous grounded qualitative work, focus groups and current research. Participants will be purposively sampled to ensure a range of demographics and experiences are captured.

Interviews will take place at 3 months following donation or withdrawal from the process.

Other Study Data

Linkage to the NHS Blood and Transplant records will provide physiological outcome data as well as information donation procedure for all donors. Physiological data will be collected pre- and post-donation, as well as at 12-months following donation, as per national donor follow-up protocol. This is described in Appendix 4. NHSBT data will be collected retrospectively once a participant completes the 12-month questionnaire, or earlier should they choose to withdraw from the study. Consent to obtain NHSBT data will be obtained through the initial study participation consent form. Subsequent to this a formal request for data access from the NHSBT will be made.

2.4 Study Outline



2.5 Trial Statistics and Data Analysis

The study endpoints will be:

- time from enquiry to donation (for those that proceed to donation)
- mental and physical health at 3 and 12 months post donation, compared to directed donor controls

All primary analyses will be undertaken by the study statistician and investigator / research fellow in accordance with a predetermined analysis plan.

Descriptive analysis will be used to describe the proportion of people who withdraw or proceed to donation, and the reasons for failing to proceed. The analysis will include all individuals enquiring about donation, with the dependent variable an indicator for each proceeding to donation. Centre-level structural and attitudinal factors identified in our parallel study (IRAS 170483) will be included in the models to determine whether these variables explain variation in donation rates.

Descriptive analysis will be used to compare baseline variables for individuals that express an interest in donation that: i) the transplant team withdraw from donation; ii) those who decide not to proceed; iii) those that proceed to donation; and iv) the specified kidney donor control group, who either proceed or do not proceed to donate. Linear or logistic mixed-effects models will be used to estimate difference in outcome variables at the 3 and 12 months follow-up assessments between the groups at the outcome assessments. Group membership and follow-up assessment (time) will be included in the models as dummy variables. Interaction terms for group and time will allow for assessments of differences at individual time points. Models will adjust for potential demographic confounders measured at baseline (e.g. age, sex, education, ethnicity) and the baseline level of the outcome variable. Missing outcome data is under the assumption that data is missing at random. Sensitivity analysis will be performed to assess this assumption.

The analysis of qualitative data will be performed using the Framework (thematic) approach as described above.

3. Sample Size and Selection

Focus Groups

The two focus groups will recruit 8-12 potential or actual unspecified kidney donors each. These will be volunteers identified by UK transplant co-ordinators.

Questionnaires

Consecutive people contacting each of the transplant centres in the UK between April 2015 and Feb 2018 will be recruited to participate in the study. Based on current trends we conservatively estimate that there will be at least 279 kidney transplants from unspecified altruistic donors during that period. Indeed, there were 107 UKD in the UK in 2013. Assuming that the proportion of individuals contacting transplant centres who go on to donate remains stable (36%, based on data from Portsmouth in 2012), we expect that 780 people considering unspecified altruistic donation will contact transplant centres during that period. Based on our previous retrospective study, we expect at least a 80% recruitment rate- that is, 624 in total, of which 224 will go on to donate). This recruitment rate is higher than is typical for longitudinal studies but justifiable given the population being studied. A sample size of 624 will provide sufficient precision to estimate the 95% confidence interval for proceeding to donation to within $\pm 4\%$ overall, and to within $\pm 18\%$ for each centre. In summary we aim to recruit 224 who have undergone unspecified donation and 400 who failed to donate.

The control group will recruit 200 people who are donating to friend or relative (specified donors) and 200 individuals that intend to donate to a friend or relative but do not (specified non-donors). Based on our retrospective study we expect a recruitment rate of 80%. Therefore we will need to approach 500 specified donors. Given a stable rate of approximately 1000 specified donations per year across the UK, we anticipate that we will be able to recruit the control group using the same three-year recruitment window as the main cohort. If there is no difference between the unspecified altruistic and specified donors on the physical and psychological variables at 12 months, it will be possible to determine that the lower limit of a one-sided 95% confidence interval will be above the non-inferiority limit of a standardised mean difference of 0.3, which is deemed to be the smallest acceptable clinically meaningful difference – this allows for 20% missing data due to drop-out, at a significance level of 5% with 90% power (14). These individuals will be recruited through transplant co-ordinators nationally.

Interviews

Qualitative interviews will also be completed with a sample of 15 donors who completed their donation, 15 who withdrew and 15 who are withdrawn by the transplant team from the process. These individuals will be identified from the initial cohort of patients that approached transplant centres with the intention to donate altruistically.

Recruitment Targets

The following recruitment targets have been set:

Focus Groups: 2 focus groups of 8-12 previous donors and 8-12 non-donors.

Total: 16-24 participants

Questionnaires:

624 potential donors (test population)

224 who have donated

400 who did not donate

200 specified donors (control population 1) and 200 specified non-donors (control population 2)

Total: 1024 participants

Interviews:

15 potential donors that donated

15 potential donors that did not donate (self-withdrawn)

15 potential donors that did not donate (withdrawn by clinical team)

Total: 45 participants

Inclusion criteria

Any individual contacting a transplant centre to enquire about unspecified donation, who proceeds beyond the initial telephone conversation, and is able to give informed consent will be considered as a potential study participant. Non-English speakers will be included and adequate translation facilities will be provided. Individuals who have already begun the donation work-up process at the time of study commencement will also be eligible for recruitment provided they are more than 2 weeks away from donation. Control participants will be recruited from those individuals contacting a centre in order to donate to a known individual.

Exclusion criteria

Any individual who declines to participate at any stage will be excluded from the study. Individuals lacking capacity will also be excluded as will any individual not eligible to donate in UK. This includes foreign nationals who are unable to donate altruistically in their country of residence or prisoners.

4. Study procedures

4.1 Consent

The study research fellow or study manager will be notified of any eligible individuals by UK transplant co-ordinators. Potential participants will be invited to participate in the study by phone or post and will be provided with an information sheet prior to the consent process. Separate consent forms have been designed for each of the three study arms (focus groups, questionnaires and interviews). Where necessary these will be translated or explained by an interpreter. Individuals who agree to participate will be asked to complete a baseline assessment, in either paper or online format. Pre-operative assessments will be completed one week prior to donation.

The following study documents have been created (Appendix 5):

- PIS Unspecified Donors Focus Group
- PIS Unspecified Donors Questionnaire and Interview Group
- PIS Specified (Control) Donors Questionnaire and Interview Group
- Consent Form Unspecified Donors Focus Group
- Consent Form Unspecified Donors Interview Group

- Consent Form Unspecified Donors Questionnaire Group
- Consent Form Specified (Control) Donors Questionnaire Group

4.2 Follow up Procedures

Follow-up assessments will be sent by post (and made available to complete online). To minimise loss to follow up anyone who has failed to return their 12 month follow up assessments within 14 days will be contacted by phone with the aim of collecting information on at least the primary outcome variable.

4.3 Maximizing Response Rates

Unspecified donors are highly motivated individuals, who, in our experience, are enthusiastic about participation in studies that may help other donors. The response rate of 74% in our previous study, whilst too low for definitive conclusions in a retrospective study, is nevertheless higher than expected for a questionnaire survey (6).

However, it is vital that response rates are high enough to accurately capture outcomes, and we aim to achieve this as follows:

- I. Participants presenting for donation will be contacted directly by the research fellow or study manager (usually by telephone or email). Non-responders will be contacted on 2 occasions, including using an alternative method (such as a written letter and/or telephone calls outside standard working hours).
- II. Participants will be given the opportunity to return documents in a freepost envelope or by completing an online form.
- III. The trial manager will contact all 23 transplanting centres on a regular basis to ensure that those who present for unspecified donation have been considered for inclusion in the study.
- IV. One team member (LB) already has close and regular contact with donor co-ordinators (who are the first point of contact for any donor presenting at a transplant centre) in all transplanting centres. She will send reminders to all co-ordinators regularly to ensure continued referral of potential participants.

We will monitor the success of this approach using the internal pilot study described in the protocol.

4.4 End of Study Definition

Completion of the final questionnaire (at 12 months) of participants recruited in the 3-year period will mark the end of the study.

5. Laboratories

No laboratory facilities will be used for this study

6. Assessment of Safety

6.1 Emotional or Psychological Distress

If any clinical concern is identified by the research team from the questionnaires or interviews (for example suicidal thoughts, or severe depression) the local clinical team (transplant donor co-ordinator) will be informed with a view to referral to the local psychological or counselling service; we used this approach previously in our retrospective study. The provision of this facility is part of our commitment to good practice and we do not anticipate this will be needed. In the unlikely event that concern is raised about a participant who has withdrawn from the donation process early in the assessment period and has no further contact with their local transplant

unit, we will contact their GP directly. The GP contact details will be collected as part of the recruitment baseline data.

6.2 Impact of study on decision to donate

Focus Groups. The Focus Groups will be with altruistic kidney donors who have already completed or have withdrawn from the donation process. As such, the study will not be able to impact on their decision to donate from the perspective of the focus group alone.

Interviews. The qualitative interviews will be performed prospectively and take place three months after donation or withdrawal from the donation process. This is to avoid any undue influence on the participant's decision to donate. The interviews will be conducted by experienced qualitative researchers who have interviewed both altruistic kidney donors and donors withdrawn from the process. The REC applications associated with these previous projects are: Understanding Barriers and enablers to altruistic kidney donation v1.14/SW/1105 and 10/H0203/11-Understanding the experiences of altruistic kidney donors. (Clarke, A., Mitchell, A., & Abraham, C. 2013. Understanding donation experiences of unspecified (altruistic) kidney donors. British Journal of Health Psychology.)

Questionnaires. The questionnaires which will be used are validated and widely used research tools which are regularly employed in the fields of social science and psychology. We have previously used similar tools in our research with no significant impact on the participants' mental or physical health.

6.3 Sensitive questions

Additional questions that will be formulated as a result of the focus group data (in addition to the already validated questionnaires) will be discussed amongst the study research group that consists of psychologists, transplant surgeons and nurses who are highly aware and sensitive to the process of altruistic donation as a result of their extensive clinical experience. Furthermore, two members of the 'Give a Kidney Charity', who represent the altruistic donor community, will review and be involved in the development of any further questions. Any new questions that would potentially impact on the decision to donate will be excluded from the questionnaire bundle

6.4 Ethics Reporting

Reports of related and unexpected SAEs will be submitted to the Main REC within 15 days of the chief investigator becoming aware of the event, using the NRES template. The form will be completed in typescript and signed by the chief investigator. The Coordinator of the main REC will acknowledge receipt of safety reports within 30 days. A copy of the SAE notification and acknowledgement receipt should be sent to the R&D Directorate.

No SAE are expected for this study.

7. Trial Steering Committee

7.1 Study Steering Committee

The study does not have a Data Monitoring Committee, but there is a Study Steering Committee (SSC), which will have the following responsibilities:

- i) To provide advice, through its Chair, to the Chief Investigator, the Project Sponsor, the Project Funder, the Host Institution and the Contractor on all appropriate aspects of the project
- ii) To concentrate on progress of the project, adherence to the protocol, patient safety (where appropriate) and the consideration of new information of relevance to the research question
- iii) The rights, safety and well-being of the participants are the most important considerations and should prevail over the interests of science and society
- iv) To ensure appropriate ethical and other approvals are obtained in line with the project plan

- v) To agree proposals for substantial protocol amendments and provide advice to the sponsor and funder regarding approvals of such amendments
- vi) To provide advice to the investigators on all aspects of the project

The SSC will be constituted as follows: An independent chairperson, an independent statistician, one member (not directly involved with the study) from within the Trust, one member external to the Trust, and two service users. An observer from the sponsor and from the CLRN will be invited to attend.

The SSC will meet at 4 to 6 monthly intervals, or more frequently if the Chairperson deems this to be necessary.

The study steering committee has the following members:

Chair
Prof Kenneth Farrington
Consultant Renal Physician
ken.farrington@nhs.net

Independent statistician
Dr Matthew Robb
NHBST

One member (not directly involved with the study) from within the Trust
Dr David Game
Consultant Renal Physician

One member external to the Trust:
Dr Sian Griffin
Consultant Renal Physician

Previous Service Users
Mr Peter Cordwell

Mr Nicholas Crace

7.2 Trial Management Committee

The Trial Management Committee manages the project on a regular basis. It consists of members of the project team and meets at 3 to 6 month intervals. Minutes and agendas are issued in the regular manner. The committee has two permanent PPI members representing the 'Give a Kidney' Charity.

8. Ethics & Regulatory Approvals

15/SC/0637

South Central – Berkshire B Research Ethics Committee Health Research Authority

Bristol HRA Centre, Level 3, Block B,
Whitefriars, Lewins Mead, Bristol BS1 2NT
HRA (Bristol Centre): 0117 342 1382 | www.hra.nhs.uk, nrescommittee.southcentral-berkshireb@nhs.net

9. Data Handling

Confidentiality

A database will be constructed by the Guys and St. Thomas Biomedical Research Centre. Online or paper questionnaires and interview transcripts will be transferred to the database, held on a secure server at either Guys Hospital or Plymouth University, in an anonymous fashion, with password protected access. Access to the database will be limited to the study researchers, Chief investigator and study manager. Participant data will be managed in accordance with the Data Protection Act, NHS Caldicott Guardian, The Research Governance Framework for Health and Social Care and Research Ethics Committee Approval.

Each patient will have a unique study identity number which will avoid the use of patients' hospital numbers, NHS numbers, dates of birth or names. The Chief Investigator will have a separate key linking the study identification number with identity of the study participants. The study key information will be kept in a separate password secure and locked environment.

Back-up will be performed automatically by the Trust systems, and data archiving will be undertaken by the Kings Health Partners Joint Clinical Trials Office, according to their standard operating procedures.

Record Retention and Archiving

All records will be kept in secure conditions. When the research trial is complete the records are kept for a further 5 years.

Compliance

The CI will ensure that the trial is conducted in compliance with the principles of the Declaration of Helsinki (1996), and in accordance with all applicable regulatory requirements including but not limited to the Research Governance Framework, Trust and Research Office policies and procedures and any subsequent amendments.

Non-Compliance

The sponsor will assess the non-compliances and action a timeframe in which they need to be dealt with. Each action will be given a different timeframe dependant on the severity. If the actions are not dealt with accordingly, the R&D Office will agree an appropriate action, including an on-site audit.

10. Finance and Publication Policy

10.1 Funding

The research is funded by the National Institute of Health Research (NIHR) HS&DR Award (13/54/54), with a total value of £872,756.

National Institute of Health Research
University of Southampton
Alpha House
Enterprise Road
Southampton
SO16 7NS

The funds will be managed centrally by the research team at Guy's Hospital and distributed to collaborating units according to a collaboration agreement, which has been negotiated by the relevant academic, financial and legal departments within the collaborating universities and hospitals.

10.2 Outputs

There will be several specific outputs in addition to published manuscripts and conference presentations:

- a) A report to NHSBT and the BTS, summarising the findings of the study
- b) National guidelines, produced in conjunction with NHSBT and the BTS
- c) A protocol for management of those presenting for unspecified donation
- d) A report to the Renal Transplant Clinical Reference Group, which reports to NHS England (which commissions transplant services in England), and to the Scottish, Welsh and Northern Irish Departments of Health.

The process for developing these outputs (beyond the first, which will be written by the study team) is as follows:

National Guidelines

The transplant community is small, and there is a widespread desire for guidance on unspecified donation. Existing guidelines on living donation are extensively used by donor teams, and these have been important in changing culture. We recognize that guidelines are not, however, necessarily effective by themselves at changing practice- in this regard, the close liaison that one team member (LB) has with donor co-ordinators at all transplant centres, and the living donor forum which she organizes, will be vital.

The support of the BTS Clinical Trials Committee for this study (attached) is indicative of the close involvement and support of the BTS. There is an existing process for developing guidelines by the BTS, through the BTS Standards Committee. We will convene a small group, including NHSBT and BTS representatives, as well as service users, to draft a guideline, which can be sent to the BTS Standards Committee for consideration. Typically, this is opened for public consultation via the BTS website for a short period, revised and then disseminated to all units. The leads for this work will be Prof N Mamode and Ms L Burnapp.

Commissioners' report

The Chief Investigator is a member of the Renal Transplant Clinical Reference Group (CRG) and has been involved in drafting Service Specifications for transplantation. He will send a report, which will be drafted with the help of the study team, including service users, to the CRG for discussion and dissemination to NHSE and counterparts in other constituent countries.

References

1. Organ Donation and Transplantation Activity Report 2013-14.
http://www.organdonation.nhs.uk/statistics/transplant_activity_report
2. Are 'anonymous' and 'non-directed' prerequisites for living altruistic donation? The views of transplant physicians from France and Québec. Fortin MC1, Dion-Labrie M, Hébert MJ, Achille M, Doucet H. Soc Sci Med. 2008 Jul;67(1):147-51.
3. Understanding donation experiences of unspecified (altruistic) kidney donors. Clarke A, Mitchell A, Abraham C. Br J Health Psychology 2014 May;19(2):393-408
4. A Testing Time? Exploration of the Psychiatric Assessment of Non-directed Altruistic Kidney Donors. Murray, A., Mitchell, A., & Clarke, A. Unpublished Declin Manuscript.
5. A Duty of Care towards the 'Good Samaritan': Experiences of Living Donor Co-ordinator Nurses working in Non-directed Altruistic Kidney Donation. Obradovic, I., Mitchell, A., Burnapp, L., & Clarke, A. Unpublished Declin Manuscript.
6. Motivations, Outcomes and Characteristics of Unspecified (Non-Directed Altruistic) Kidney Donors in the United Kingdom. Maple H, Chilcot J, Burnapp L, Gibbs P, Santhouse A, Norton S, Weinman J, Mamode N. Transplantation 2014 *In Press*.
7. Altruistic kidney donation: donor characteristics and outcomes from 11 years' experience. Griffin JL, Morton J. Br J Surg. 2011;98:147
8. Good samaritan kidney donation. Morrissey PE, Dube C, Gohh R, Yango A, Gautam A, Monaco AP. Transplantation. 2005;80(10):1369-73.
9. Twenty-two nondirected kidney donors: An update on a single center's experience. Jacobs CL, Roman D, Garvey C, Kahn J, Matas AJ American Journal of Transplantation. 2004;4(7):1110-6.
10. Kidney transplant chains amplify benefit of nondirected donors. Melcher ML, Veale JL, Javaid B, Leiser DB, Davis CL, Hil G, Milner JE. JAMA Surg. 2013 Feb;148(2):165-9.
11. UK Living Donor Strategy to 2020. NHSBT. www.organdonation.nhs.uk
12. The living anonymous kidney donor: Lunatic or saint? Henderson AJZ, Landolt MA, McDonald MF, Barrable WM, Soos JG, Gourlay W, et al. American Journal of Transplantation. 2003;3(2):203-13.
13. Qualitative research practice: A guide for social science students and researchers. Ritchie, J., & Lewis, J. (Eds.). (2003). Sage
14. Sample sizes for clinical trials with normal data. Julious SA. Statist. Med. 2004; 23:1921-1986
15. Costing psychiatric interventions. Beecham, J. K. & Knapp, M. R. J. In Measuring Mental Health Needs 1992 (eds G. Thornicroft, C. Brewin & J. K. Wing). London: Gaskell.
16. Unit Costs of Health & Social Care 2012. Curtis L (2012) Report on behalf of Personal Social Services Research Unit. University of Kent, Canterbury. ISBN: 978-1-902671-82-6
17. Design and analysis of pilot studies: recommendations for good practice. Lancaster GA, Dodd S, Williamson PR. J Eval Clin Practice 2004;10:307-312

References for Validated Questionnaires:

The Multidimensional Scale of Perceived Social Support. Zimet GD, Dahlem NW, Zimet SG & Farley GK. Journal of Personality Assessment 1988; 52: 30-41.

Society and the adolescent self-image. Rosenberg, M (1965). Princeton, NJ: Princeton University Press

A brief measure for assessing generalized anxiety disorder: the GAD-7. Spitzer RL, Kroenke K, Williams JB, et al; Arch Intern Med. 2006 May 22;166 (10):1092-7

The PHQ-9: validity of a brief depression severity measure. Kroenke K (1), Spitzer RL, Williams JB. J Gen Intern Med. 2001 Sep; 16 (9):606-13.

New measures of well-being: Flourishing and positive and negative feelings. Diener E, Wirtz D, Tov W, Kim-Prieto C, Choi D, Oishi S, & Biswas-Diener R. (2009). Social Indicators Research, 39, 247-266.

User Manual- Decision Regret Scale (document on the Internet). O'Connor AM. Ottawa: Ottawa Hospital Research Institute. 1996 (modified 2003; cited 2016 04 10) Available from https://decisionaid.ohri.ca/docs/develop/User_Manuals/UM_Regret_Scale.pdf

A Very Brief Measure of the Big Five Personality Domains. Journal of Research in Personality. Gosling SD, Rentfrow PJ, & Swann WB Jr. (2003), 37, 504-528.

The Satisfaction with Life Scale. Diener E, Emmons R A, Larsen R J, & Griffin S. (1985) Journal of Personality Assessment, 49, 71-75.

SF-12v2™ Health Survey. 1992-2002 by Health Assessment Lab, Medical Outcomes Trust and QualityMetric Incorporated. All rights reserved. SF-12® is a registered trademark of Medical Outcomes Trust. (IQOLA SF-12v2 Standard, English (United Kingdom) 8/02)

Beecham J, Knapp M (2001) Costing psychiatric interventions. In Thornicroft G (ed) Measuring mental health needs. London, Gaskell

Information with regards to Safety Reporting in Non-CTIMP Research

	Who	When	How	To Whom
SAE	Chief Investigator	-Report to Sponsor within 24 hours of learning of the event -Report to the MREC within 15 days of learning of the event	SAE Report form for Non-CTIMPs, available from NRES website.	Sponsor and MREC
<u>Urgent Safety Measures</u>	Chief Investigator	Contact the Sponsor and MREC Immediately Within 3 days	By phone Substantial amendment form giving notice in writing setting out the reasons for the urgent safety measures and the plan for future action.	Main REC and Sponsor Main REC with a copy also sent to the sponsor. The MREC will acknowledge this within 30 days of receipt.
<u>Progress Reports</u>	Chief Investigator	Annually (starting 12 months after the date of favourable opinion)	Annual Progress Report Form (non-CTIMPs) available from the NRES website	Main REC
<u>Declaration of the conclusion or early termination of the study</u>	Chief Investigator	Within 90 days (conclusion) Within 15 days (early termination) <i>The end of study should be defined in the protocol</i>	End of Study Declaration form available from the NRES website	Main REC with a copy to be sent to the sponsor
<u>Summary of final Report</u>	Chief Investigator	Within one year of conclusion of the Research	No Standard Format However, the following Information should be included:- Where the study has met its objectives, the main findings and arrangements for publication or dissemination including feedback to participants	Main REC with a copy to be sent to the sponsor

Appendix 1

Appendix 1. Projected number of unspecified kidney donors to 2020




Figure 1. Number of living kidney donations in the UK by unspecified and specified kidney donors


Figure 1 shows the total number of living kidney donations in the UK per year, separated by unspecified and specified type. Over the six year period shown there has been an increase in the total number of living donations. Since 2010, the increase is driven by unspecified donations, with the number of unspecified donations actually falling.

Figure 2 and Table 1 below shows projections for the number of unspecified (altruistic) kidney donors to 2020 by two methods. The first method fits a linear rate of increase based on the past trend. The second method assumes a non-linear (quadratic) rate of change. The quadratic method fits the observed data best but this is no indication it provides a more accurate projection.

With recruitment between March 2015 and Feb 2018 we can expect between 279 (linear = $83+93+103$) and 493 (non-linear = $131+163+199$) donors based on the projections. The expected sample size will be based on the more conservative linear estimate. An even more conservative estimate would assume rates staying stable at the 2012 figures. This would mean the expected sample size of 180 ($60+60+60$).


Appendix 2a

Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations




Appendix 2b

Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations



Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations


Unspecified Donors



Appendix 2d

Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations


Unspecified Donors



Appendix 2e

Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations


Unspecified Donors



Appendix 2f

Appendix 2: Diagram to demonstrate how unspecified donors can prime a chain of donations

Unspecified Donors



Appendix 3

BOUnD

Appendix 3: Topic list: Qualitative interviews

Appendix 4:

	Data (source)	Phase 1: Potential donors				Phase 2: Outcomes			
		Recruitment		Pre donation		Donation		Follow-Up	
		Pre-enrolment	Enrolment	Baseline	Pre-op (1 week)	Surgery	Post-op	Follow-up 3 months	Follow-up 12 months
Enrolment	Initial contact (Centre)	X							
	Referral (Centre)	X							
	Eligibility screen (Trial manager)		X						
	Informed Consent (Trial manager)		X						
Assessments	Baseline demographic data			A					
	Personality: TIPI (Questionnaire)			A					
	Social support: MSPSS (Questionnaire)			A	A			A	A
	Peri-operative physical outcome data (NHSBT)				A		D		D
	Surgical procedure data (NHSBT)					D			
	Withdrawal (Questionnaire & Centre data)			W	W				
	Rosenberg (Questionnaire)			A	A			A	A
	SWLS (Questionnaire)			A	A			A	A
	PHQ9 & GAD7 (Questionnaire)			A	A			A	A
	Flourishing Scale			A	A			A	A
	Decision Regret Scale							A	A
	SF12 (Questionnaire)			A	A			A	A
	Client Service Receipt Inventory (Questionnaire)			A	A			A	A
	In-House Questionnaire			A	A			A	A

X= recruitment and centre level data (pre-consent)

A= Data from all potential donors

D= Data for donors only

W= Data for withdrawn only