

**Developing a framework for a novel multi-disciplinary,
multi-agency intervention(s), to improve medication
management in older people on complex medication
regimens resident in the community (MEMORABLE) – a
realist synthesis**

IRAS Number: 211031

PROSPERO Number: CRD42016043506

Sponsor Aston University

FUNDER NIHR Health Services and Delivery Research.

FUNDERS Number: 15_137_01

Version Control

<u>Version number and date</u>	<u>Author</u>	<u>Next Steps</u>
V0.1 – 7/11/2016	IM	Sent to co-applicants.
V1 – 9/11/2016	IM	Submitted to HS&DR
V2 – 03/07/17	SL	Submit to Aston (AP) for review
V3 – 11/07/17	SL	Minor revisions as part of review
V4 – 08/11/17	IM	Minor revisions to recruitment
V5 – 30/01/18	IM	Added Hadar Zaman as co-applicant
V6 – 7/02/18	IM	Increased the sample size for carers and added new recruitment sites (page 9).

Table of Contents

1.	Summary	5
2.	Background	6
3.	Aims/Objectives	6
4.	WP1: Literature Review	7
5.	WP2: Realist Interviews	9
6.	WP3: Developing the Framework for an Intervention(s)	9
7.	Project Management and Governance	10

1. Summary

Background:

Medication related adverse events have been estimated to be responsible for 5,700 deaths and cost the United Kingdom £750 million annually. This burden falls disproportionately on older people. Outcomes from interventions to optimise medication management are caused by multiple context sensitive mechanisms. The MEdication Management in Older people: REalist Approaches BAseD on Literature and Evaluation (MEMORABLE) project uses realist synthesis to understand how, why, for whom and in what contexts interventions, to improve medication management in older people on complex medication regimes residing in the community, work.

Method

This realist synthesis uses primary data from interviews to develop the programme theory. A realist logic of analysis will synthesise data both within and across the two data sources to inform the design of a complex intervention(s) (see flow diagram: Appendix 1) to evidence a framework to help improve medication management in older people.

1. Literature Review

The review (using realist synthesis) contains five stages to develop an initial programme theory to understand why processes are more or less successful and under which situations: Focussing of the research question; developing the initial programme theory; developing the search strategy; selection and appraisal based on relevance and rigour; and data analysis/synthesis to develop and refine the programme theory and context, intervention, mechanisms configurations.

2. Realist Interviews / focus group

Realist interviews will explore and refine our understanding of the programme theory developed from the realist synthesis. Up to ~~4530~~ older people and their informal carers (15 older people with multi-morbidity, ~~2540~~ informal carers and 5 older people with mild dementia), and 20 health and care practitioners will be identified by promoting the research through link organisations. They will be consented and interviewed by the RA. Data will be held securely, confidentially and anonymised. Later in the research, if a focus group appears to be a suitable means of continuing people's engagement, one may be set up, subject the agreement of participants.

3. Developing framework for the Intervention(s)

Data from the realist synthesis and interviews / focus group will be used to refine the programme theory for the intervention(s) to identify: the mechanisms that need to be 'triggered', and the contexts related to these mechanisms. Intervention strategies that change the contexts so the mechanisms are triggered to produce desired outcomes will be developed. Feedback on these strategies will be obtained.

2. Background

Between 2010 and 2050 the number of older people living in the United Kingdom (UK) will nearly double from around 10 to 19 million (1). Half of people aged 75 or more are living with two or more long-term conditions (2). The Francis Report emphasised that patients should be protected from avoidable harm (3). Yet, medication related adverse events have been estimated to be responsible for 5,700 deaths, at a cost of £750 million to the UK health service, every year (4). Furthermore, between 5 to 8% of unplanned hospital admissions in the UK, are related to medication issues (5).

Safe and effective medication management is particularly challenging in older people with chronic or long-term conditions on multiple medications with some degree of frailty and/or cognitive impairment (6)(7)(8)(9)(10)(11)(12). Older people, particularly those with dementia, may be more susceptible to medication related adverse events and less able to identify when a medication error has taken place (6) (8) (13) (14).

Complex care pathways delivered by a diverse range of care staff are needed to support medication management (2) (15). For instance, older people often attend multiple clinics and interact with multiple health and social care professionals including GPs, community pharmacists, district nurses and various secondary care clinicians, all of whom help to optimise their medication management (6) (15). In addition, for many older people, formal and informal carers have a key and pivotal role and provide front-line support for their medication management needs (16) (17). However, formal carers frequently lack the appropriate training to deliver this role and informal carers, who often have self-care difficulties themselves, struggle with the responsibility and find the role stressful and burdensome (17) (18) (19) (20) (21). Recent NICE guidance has identified the need for a collaborative approach to supporting older people with long-term conditions in managing their medication(s) (22).

Realist approaches aim to understand how a given context affects any mechanism, to generate either a positive or a negative outcome (23)(24). In other words, they explore the relationship between context, mechanism and outcome (CMO) configurations or CMOCs (23). Our MEducation Management in Older people: Realist Approaches Based on Literature and Evaluation (MEMORABLE) study uses a realist synthesis to address the three steps within the Medical Research Council framework for Developing Complex Interventions: identifying the evidence base, identifying/developing theory and modelling process and outcomes (25) (see flow diagram for overview: Appendix 1). Recognised guidelines for ensuring the quality of the review and how it is reported (the RAMESES guidelines) will be followed (23) (26) (27). In this study the realist programme theories will be developed through a combination of secondary (literature review, in the form of a realist synthesis: Work Package 1) and primary (realist interviews: Work Package 2) data collection methods. In Work Package (WP) 3 we will use data from within and across WP1 and WP2 to further develop and refine the realist programme theory. The data collected from each work package will be continually analysed to enable emerging findings from one package to inform the other. Finally, the resulting refined realist programme theory will be used to design the framework for the intervention(s) (WP3).

3. Aims/Objectives

Aim: To use realist synthesis including primary data collection to develop a framework for a novel multi-disciplinary, multi-agency intervention(s), to improve medication management in older people on complex medication regimens resident in the community.

Objectives

1. To understand how and why any potentially relevant interventions, to optimise medication management, work (or do not work) for particular groups of older people in certain circumstances.
2. To synthesize the findings from objective 1 into a realist programme theory of an intervention(s) to support older people living in the community manage their medication.
3. To use realist programme theory developed from objective 2 to inform the development of an intervention(s) to assist older people living in the community to manage their medication.

4. WP1: Literature Review

Method

This literature review (in the form of a realist synthesis) comprises five stages:

1. Focussing the Synthesis: the overall focus is to develop a realist programme theory for an intervention to support older people living in the community to manage their medication. The realist synthesis will be focussed on the most relevant strategies that would be needed to be used in the medicine management intervention developed.

2. Developing Initial Programme Theory: a programme theory sets out how and why outcomes occur within an intervention (23) (28).

We will iteratively consult with our stakeholder group, and informally search the literature to locate current theories (29). This informal searching is more exploratory than the formal searching in step 3 and aims to rapidly identify a range of possibly relevant explanatory theories. Iterative discussions in the project team will aim to understand and synthesise the various theories into an initial coherent programme theory. The stakeholder group (that includes patient and public involvement) will provide content expertise to help synthesis the initial programme theory and later refinement.

The initial programme theory (candidate theory) will be refined as the synthesis progresses with input from both the project team and stakeholder group. The theory will be mapped out as a series of outcome steps required for the final desired outcome, identifying intermediate outcomes that take place either sequentially or in parallel (26). For each step (where possible at this stage) the relevant and associated context and mechanism for each outcome will be developed from data identified from our informal searches (26) (30).

3. Developing a Search Strategy: we will use a CIMO (Context, Intervention, Mechanisms, Outcome) question framework (31) to construct an initial sampling frame for medication management in older people. The initial CIMO framework is:

Context: older people living in the community, medication complexity, ethnicity. Subgroup: people with dementia living in the community.

Intervention: based on previous work and literature reviews possible interventions include support from formal or informal carers, education, medication review, self-management, tools (including technology) to support adherence.

Mechanisms: the mechanism(s) triggered by the intervention will be identified by the programme theory.

Outcomes: quality of life, adherence, adverse events, carer burden, economic (care costs including residential care and hospitalisations).

We will then select distinct subsets of literature from within the sampling frame with which to test emerging theory (30) (32). At this point we anticipate that we will be able to use a comprehensive sampling approach. However, if the size of the literature proves unmanageable then we will employ a variety of appropriate sampling strategies (e.g. theoretical sampling, maximum variation sampling, extreme case sampling) to optimise the analytical value of the realist synthesis component, as specified by the methodology (33).

The initial sampling frame will be used as the starting point for selection of 'index papers' from which suggested conceptual or contextual explanations will be identified, developed and explored by following links out to wider bodies of relevant literature (30) (32). When retrieved data suggests certain mechanisms may be particularly important, the search techniques will be refined to identify data from other clinical environments where these mechanisms may also be in operation, so that we can better understand their behaviour under different contexts.

Realist synthesis uses iterative, purposive sampling from a wide range of evidence to develop, refine, confirm and refute theories about how and why an intervention works, for whom, and in what circumstances (33). Consequently, the search strategy will be developed iteratively and re-visited at

predetermined milestones, using different permutations and additional concepts (30) (32). Searching will be guided by the need to find data to develop the programme theory and hence additional searching may be needed.

We will subsequently use 'cluster searching' to identify 'clusters' of data from related publications. This approach will add to the conceptual richness and contextual thickness of studies initially identified within the sampling frame constructed through conventional topic-based searching (32). We will identify "sibling" (i.e. directly linked outputs from a single study) and "kinship" (i.e. associated papers with a shared contextual or conceptual pedigree) papers and reports to add richness of data while preserving both rigour and relevance (32). Active pursuit of citation networks, using Google Scholar and Web of Science will be used to link index papers to the wider literature. Searching will continue until sufficient data is found ('theoretical saturation') to conclude that a candidate programme theory is sufficiently coherent and plausible (30).

We will use the most up-to-date methodological literature when devising search strategies relating to older people or medication management (34). Based on previous work international guidance and discussions with information specialists, sources will include: Scopus, Web of Science, EMBASE/PubMed/MEDLINE, Cochrane, CINAHL, PSYCINFO, ProQuest, Sociological Abstracts, Google Scholar, BASE (Bielefeld Academic Search Engine)/ETHOS (British Library Electronic Thesis Online)/ProQuest Dissertations and Thesis, Grey Literature in Europe (<http://www.opengrey.eu/>) and NHS Evidence and equivalent, external experts and charities/user groups and reference lists of relevant papers (30) (35).

4. Selection and Appraisal: Inclusion and exclusion are based on (30) (36):

- Does the document contain any data that can contribute to developing or testing theory (**relevance**)?
- Are the methods (if any) utilised to generate the relevant data trustworthy and credible (**rigour**)?

Selection and appraisal is a two-step process:

1). Potentially relevant documents will initially be screened by title, abstract and key words (30) (32) by the Research Associate (RA). A 10% random sample will be checked by AB and GW for consistency (any disagreements will be resolved with the input of IM);

2). The full texts of this set of documents will be obtained and screened by the RA. Again a 10% random sample will be checked by AB and GW for consistency using the same approach as outlined above.

The full texts of all relevant documents will be imported into NVivo (a qualitative data analysis software tool). Relevant data from included documents will be coded into NVivo. Some of the codes will come from the programme theory (i.e. deductive coding). Others will come from the data (i.e. inductive coding). These codes will cover concepts that are judged to be important and potentially relevant to the programme theory. When coding, where it is possible to make such inferences, data will be coded as context, mechanism or outcome. Any data that informs the relationship of data within Context-Mechanism-Outcome-Configurations (CMOCs) or between the CMOCs (contained within the programme theory) will also be coded (23) (36).

5. Data Analysis and synthesis: will configure the coded data to develop the CMOCs within the programme theory by piecing together data from different sources (23). Relevant data will be interpreted as being about context, mechanism and/or outcome within the CMOCs contained within our overarching programme theory. These data may come from a range of included documents or from the interview transcripts (see WP2 below). The configurations will be presented and discussed and debated within our regular project team meetings. We will ask a number of questions about the data informed by the approach used in Wong et al. (see Box 1 in the reference) (29). These questions will cover: relevance, interpretation of meaning, judgement regarding the CMOCs, judgements about the programme theory and rigour. Part of the process of data analysis will involve making inferences about the relationships of the CMOCs to the programme theory (23). Where needed, we will attempt

to link our findings to substantive theories to further refine our theoretical understanding; for example social cognitive theory is frequently referenced in connection with medication self-management and may help to explain some aspects of the programme theory (37) (38).

6. WP2: Realist Interviews

Interviews are seen as the main method to gather stories about people's experiences of medication management. Later in the research, if a focus group appears to be a suitable means of continuing people's engagement, one may be set up, subject to the agreement of participants.

Realist interviews are a type of qualitative interview where the researcher does a 'show and tell' with the participant (39). The participant is initially asked a series of general questions about the topic area (i.e. is 'eased in') and then questioned in the most neutral way possible about aspects of the programme theory. Interview Schedules for older people, carers and practitioners have been developed (see separate documents). Realist interviews will gather additional data to confirm, refute or refine aspects of the programme theory developed from the literature review work package (WP1) (40). The intention is that these interviews will be used to further develop and refine aspects of the programme theory that remain unclear based on the analyses of data from the realist synthesis. Or they may surface potentially relevant data about aspects of the programme theory that have not been found in the literature.

Method: Ethical approvals will be sought prior to the start of this WP. Focus groups and/or interviews with older people, informal carers, and care staff will be recorded and transcribed verbatim. We will ascertain from the participants whether focus groups or one-to-one interviews are most appropriate. We are mindful of the sensitive issues which may be discussed and some people may prefer a one-to-one interview although some may find the group environment empowering. Based on our practical experience one-to-one interviews are likely to be needed for care staff.

Sample size: Up to ~~4530~~ older people and their informal carers (15 older people with multi-morbidity, ~~25 40~~ informal carers and 5 older people with dementia), and 20 health and social care staff (including formal carers). Previous experience indicates that this will generate sufficient data for an in-depth analysis from multiple perspectives (41) (42). However, interviews will stop when theoretical saturation is reached. The sampling strategy will be informed by the data collected from WP1 and be directed by the need to find data relevant to explore and refine aspects of the programme theory developed in WP1. Sampling of participants will be as follows:

a. Interviews with older people and informal carers: Participants will be purposively sampled to ensure diversity in potentially conceptually-relevant characteristics (for example) including: age; ethnicity; gender; number of co-morbidities; presence of dementia; level of support from carer.

b. One-to-one interviews with health and social care professionals (including GPs, social workers, formal carers, nurses, community pharmacists, secondary care consultants) who support older people in medication management. Participants will be purposively sampled to ensure diversity in potentially conceptually-relevant characteristics (for example) including: locality (rural vs. urban) and the index of deprivation in the area that they work.

We may recruit patients, carers and care professionals, who become aware of the study via our PPI (patient and public involvement) engagement activities e.g. blogs, social media activity, radio/TV/newspaper articles, patient group meetings etc. and voluntarily offer to participate in the study.

[We will also recruit participants via Extracare Village and Retirement Homes and Staffordshire and Stoke on Trent Partnership NHS Trust.](#)

Data Analysis: NVivo software will be used to organise the qualitative data. The process of coding the data from the transcripts of the interviews / focus groups will be similar to that outlined above in WP1.

Data coding and analysis will initially be conducted by the RA. One member of the team (GW) will independently check 10% of interviews for consistency in coding. Data analysis will take place after each interview / focus group and use a realist logic of analysis. Through discussion and disputation

the project team will make inferences with respect to how the programme theory should be further refined (or not) based on the additional data from the realist interviews. In other words, asking the question how and why do these findings inform (if at all) the programme theory from WP1 and what refinements (if any) do we need to make to it? As quality control processes: a) transcripts will be shared with participants and feedback elicited as to their veracity and; b) a 10% sample of transcripts will be checked for consistencies in coding, interpretations and inferences made by GW. Any disagreements will be resolved by discussion between the RA, GW and IM, with IM being the final arbiter.

6. WP3: Developing the Framework for an Intervention(s)

The data from the realist synthesis (literature review) and realist interviews will be used to further refine realist programme theory of an intervention(s) to support older people, including people with dementia, living in the community manage their medication. If needed, at this stage we will undertake additional searching and/or interviews to find additional relevant data to refine the programme theory. However, we anticipate that by this stage we will have found enough data for us to reach theoretical saturation. We will be able to combine the data from these two sources as we will be using the same logic of analysis for both the interview data and that from the realist synthesis (33).

To move from programme theory to intervention(s) design we will:

- a) use the data from the realist synthesis and realist interviews to identify the most important mechanisms within the programme theory that need to be 'triggered' to get desired outcomes;
- b) identify which contexts are related to these 'key' mechanisms (i.e. which CMOCs are the mechanisms found in);
- c) draw on data from the realist synthesis that provides information of the intervention strategies that can change the contexts in the relevant 'key' CMOCs. In other words, for this last stage we will seek information on which intervention strategies we might be able to use to change the contexts in such a way that 'key' mechanisms are triggered to produce desired outcomes (23). Ultimately the analysis will provide information on the required intervention strategies (26) (40) (43).

These strategies will be presented to a project event involving 30 care staff plus 10 PPI reps to obtain detailed feedback on our proposed intervention strategies. This will include discussion on the plausibility, feasibility and relevance to patients and the NHS. The data from this event will be analysed and the outputs presented to the stakeholder group providing information on the required intervention strategies and thus the framework to formulate an intervention(s) for future feasibility testing.

7. Project Management and Governance

Project Management Structure

Project Management will be undertaken by Ian Maidment with support from the Research Associate and AHRIC (Aston Health Research and Innovation Cluster). The groups will support this project.

The Project Team (PT) / Project Group.

This will contain all co-applicants, the Research Associate and the Director of AHRIC (NS). The core team will be responsible for running all aspects of the project and dissemination of results.

Membership:

Ian Maidment (Senior Lecturer in Clinical Pharmacy, Aston University): Chief Investigator.

Nichola Seare (Director, Aston Health Research and Innovative Cluster): project management support.

Sally Lawson, Aston University: Research Associate.

Dr Andrew Booth, University of Sheffield: lead on realist review (WP1).

Associate Professor Judy Mullan, Wollongong University, Pharmacist: international perspective.

Dr Jane McKeown, University of Sheffield: advice on qualitative methods and PPI strategy.

Sylvia Bailey, Age UK: PPI Lead.

Dr Geoffrey Wong, University of Oxford: senior supervisory role on realist methods.

Hadar Zaman, University of Bradford: focused on challenges in the BME community.

Stakeholder Group (SG)

The SG will provide advice to the project team and provide feedback on the veracity of our programme theory as it is developed during the project. The group will also monitor progress against milestones, provide advice, promote the project, and communicate with stakeholders. The SG will advise on study documentation, and help us to refine our dissemination strategy and outputs to maximise impact and knowledge mobilisation.

The SG will have an independent chair and other members will include PPI reps, the Research Associate, IM and NS and health and social care professionals.

Sponsorship


Aston University will act as Sponsor for the study.

Research Ethics Committee (REC) Review

The study received a Favourable Opinion will be sought from East of England - Cambridgeshire and Hertfordshire Research Ethics Committee (Ref: 17/EE/0357).

Appendix 1:

Flow Chart: Developing a framework for a novel multi-disciplinary, multi-agency intervention(s), to improve medication management in older people on complex medication regimens resident in the community



References

1. Parliament UK. The ageing population: key issues for the 2010 Parliament [Internet]. [cited 2016 Oct 17]. Available from: <http://www.parliament.uk/business/publications/research/key-issues-for-the-new-parliament/value-for-money-in-public-services/the-ageing-population/>
2. Hobbs FDR, Baker M, Davies DSC. Morbidity matters: challenges for research. *Br J Gen Pract* [Internet]. 2015;65(633):e215--e216. Available from: <http://bjgp.org/content/65/633/e215>
3. Francis R. Report of the Mid Staffordshire NHS Foundation Trust Public Inquiry Executive summary Report of the Mid Staffordshire NHS Foundation Trust Public Inquiry [Internet]. 2013. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/279124/0947.pdf
4. National Patient Safety Agency. Safety in doses: medication safety incidents in the NHS [Internet]. 2007. Available from: <http://www.nrls.npsa.nhs.uk/resources/?EntryId45=59822>
5. Frontier Economics. Exploring the costs of unsafe care in the NHS [Internet]. 2014. Available from: <http://www.frontier-economics.com/documents/2014/10/exploring-the-costs-of-unsafe-care-in-the-nhs-frontier-report-2-2-2-2.pdf>
6. NICE. Medicines optimisation: the safe and effective use of medicines to enable the best possible outcomes. NICE Guidel [Internet]. 2015;(March). Available from: www.nice.org.uk/guidance/ng5/resources/medicines-optimisation-the-safe-and-effective-use-of-medicines-to-enable-the-best-possible-outcomes-51041805253
7. Mallet L, Spinewine A, Huang A. Prescribing In Elderly People 2 The challenge of managing drug interactions in elderly people. *Lancet*. 2007;370:185–91.
8. The British Geriatrics Society. Fit for Frailty: Consensus best practice guidance for the care of older people living with frailty in community and outpatient settings. 2014. 26 p.
9. Fox C, Richardson K, Maidment ID, Savva GM, Matthews FE, Smithard D, et al. Anticholinergic Medication Use and Cognitive Impairment in the Older Population: The Medical Research Council Cognitive Function and Ageing Study. *J Am Geriatr Soc* [Internet]. 2011;59(8):1477–83. Available from: <http://dx.doi.org/10.1111/j.1532-5415.2011.03491.x>
10. Jiang H-Y, Chen H-Z, Hu X-J, Yu Z-H, Yang W, Deng M, et al. Use of Selective Serotonin Reuptake Inhibitors and Risk of Upper Gastrointestinal Bleeding: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol* [Internet]. 2015;13(1):42–50.e3. Available from: <http://www.sciencedirect.com/science/article/pii/S1542356514009276>
11. Richardson K, Bennett K, Maidment ID, Fox C, Smithard D, Kenny RA. Use of Medications with Anticholinergic Activity and Self-Reported Injurious Falls in Older Community-Dwelling Adults. *J Am Geriatr Soc* [Internet]. 2015;63(8):1561–9. Available from: <http://dx.doi.org/10.1111/jgs.13543>
12. Peklar J, O'Halloran AM, Maidment ID, Henman MC, Kenny RA, Kos M. Sedative Load and Frailty Among Community-Dwelling Population Aged ≥65 Years. *J Am Med Dir Assoc* [Internet]. 2015;16(4):282–9. Available from: <http://www.sciencedirect.com/science/article/pii/S1525861014006720>
13. Maidment ID, Haw C, Stubbs J, Fox C, Katona C, Franklin BD. Medication errors in older people with mental health problems: a review. *Int J Geriatr Psychiatry* [Internet]. 2008;23(6):564–73. Available from: <http://dx.doi.org/10.1002/gps.1943>
14. Maidment I. Medication -related adverse events in older people with dementia causes and possible solutions [Internet]. 2013. Available from: <http://eprints.aston.ac.uk/21396/1/Studentthesis-2013.pdf>
15. SERCO/HSJ. Commission on hospital care for frail older people. Main Report. 2014;(November):1–69. Available from: http://www.hsj.co.uk/Journals/2014/11/18/l/q/r/HSJ141121_FRAILOLDERPEOPLE_LO-RES.pdf
16. National Voices. Supporting self-management. 2014;(September):27. Available from: http://www.nationalvoices.org.uk/sites/www.nationalvoices.org.uk/files/supporting_self-management.pdf
17. Local Democracy Think Tank. Key to Care - Report of the Burstow Commission on the future of the home care workforce [Internet]. 2014. Available from: <http://www.lgiu.org.uk/wp-content/uploads/2014/12/KeyToCare.pdf>
18. Strategic Society Centre and Independent Age. The bigger picture: understanding disability and care in England's older population [Internet]. 2014. Available from:

- <http://strategicsociety.org.uk/wp-content/uploads/2014/11/The-Bigger-Picture-Understanding-disability-and-care-in-Englands-older-population.pdf>
http://www.independentage.org/media/847199/biggerpicture_policyfinal.pdf
19. Poland F, Mapes S, Pinnock H, Katona C, Sorensen S, Fox C, et al. Perspectives of carers on medication management in dementia: lessons from collaboratively developing a research proposal. *BMC Res Notes* [Internet]. 2014;7(1):1–10. Available from: <http://dx.doi.org/10.1186/1756-0500-7-463>
 20. Lieto JM, Schmidt KS. Reduced ability to self-administer medication is associated with assisted living placement in a continuing care retirement community. *J Am Med Dir Assoc* [Internet]. 2005;6. Available from: <http://dx.doi.org/10.1016/j.jamda.2005.04.002>
 21. Gillespie R, Mullan J, Harrison L. Managing medications: the role of informal caregivers of older adults and people living with dementia. A review of the literature. *J Clin Nurs* [Internet]. 2014;23(23–24):3296–308. Available from: <http://dx.doi.org/10.1111/jocn.12519>
 22. National Institute for Health and Care Excellence. Older people with social care needs and multiple long-term conditions. NICE Guidel [Internet]. 2015;(November). Available from: <https://www.nice.org.uk/guidance/ng22>
 23. Wong G, Greenhalgh T, Westhorp G, Buckingham J, Pawson R. RAMESES publication standards: realist syntheses. *BMC Med* [Internet]. 2013;11(1):1–14. Available from: <http://dx.doi.org/10.1186/1741-7015-11-21>
 24. Greenhalgh T, Wong G, Westhorp G, Pawson R. Protocol - realist and meta-narrative evidence synthesis: Evolving Standards (RAMESES). *BMC Med Res Methodol* [Internet]. 2011;11(1):1–10. Available from: <http://dx.doi.org/10.1186/1471-2288-11-115>
 25. MRC. Developing and evaluating complex interventions: new guidance [Internet]. Vol. 337, Medical Research Council. 2008. Available from: <https://www.mrc.ac.uk/documents/pdf/complex-interventions-guidance/> (accessed 28th October 2016)
 26. Goodman C, Denning T, Gordon AL, Davies SL, Meyer J, Martin FC, et al. Effective health care for older people living and dying in care homes: a realist review. *BMC Health Serv Res* [Internet]. 2016;16(1):1–14. Available from: <http://dx.doi.org/10.1186/s12913-016-1493-4>
 27. Greenhalgh T, Wong G, Jagosh J, Greenhalgh J, Manzano A, Westhorp G, et al. Protocol—the RAMESES II study: developing guidance and reporting standards for realist evaluation. *BMJ Open* [Internet]. 2015 Aug 1;5(8). Available from: <http://bmjopen.bmj.com/content/5/8/e008567.abstract>
 28. Gordon AL, Goodman C, Denning T, Davies S, Gladman JRF, Bell BG, et al. The Optimal Study: Describing the Key Components of Optimal Health Care Delivery to {UK} Care Home Residents: A Research Protocol. *J Am Med Dir Assoc* [Internet]. 2014;15(9):681–6. Available from: <http://www.sciencedirect.com/science/article/pii/S1525861014003910>
 29. Wong G, Brennan N, Mattick K, Pearson M, Briscoe S, Papoutsis C. Interventions to improve antimicrobial prescribing of doctors in training: the IMPACT (IMProving Antimicrobial presCribing of doctors in Training) realist review. *BMJ Open* [Internet]. 2015 Oct 1;5(10). Available from: <http://bmjopen.bmj.com/content/5/10/e009059.abstract>
 30. Wong G, Westhorp G, Pawson R, Greenhalgh T. Realist Synthesis. RAMESES Training Materials. RAMESES Proj [Internet]. 2013;(July 2013):55. Available from: http://www.ramesesproject.org/media/Realist_reviews_training_materials.pdf
 31. Denyer D, Tranfield D, Van Aken JE. Developing Design Propositions through Research Synthesis. *Organ Stud*. 2008;29(3):393–413.
 32. Booth A, Harris J, Croot E, Springett J, Campbell F, Wilkins E. Towards a methodology for cluster searching to provide conceptual and contextual “richness” for systematic reviews of complex interventions: case study (CLUSTER). *BMC Med Res Methodol* [Internet]. 2013;13(1):1–14. Available from: <http://dx.doi.org/10.1186/1471-2288-13-118>
 33. Funnell S, Rogers P. Purposeful program theory: effective use of theories of change and logic models. San Francisco: Jossey-Bass; 2011.
 34. Stevenson M, Taylor BJ, Knox J. Risk in dementia care: searching for the evidence. *Health Risk Soc* [Internet]. 2015;8575(February):1–17. Available from: <http://www.tandfonline.com/doi/full/10.1080/13698575.2015.1119256>
 35. McMahon T, Ward PR. HIV among immigrants living in high-income countries: a realist review of evidence to guide targeted approaches to behavioural HIV prevention. *Syst Rev* [Internet]. 2012;1(1):56. Available from: <http://dx.doi.org/10.1186/2046-4053-1-56>
 36. Pawson R, Greenhalgh T, Harvey G, Walshe K. Realist review - a new method of systematic review designed for complex policy interventions. *J Heal Serv Res Policy* [Internet]. 2005;10.

- Available from: <http://dx.doi.org/10.1258/1355819054308530>
37. Heisler M, Hofer TP, Klamerus ML, Schmittdiel J, Selby J, Hogan MM, et al. Study protocol: The Adherence and Intensification of Medications (AIM) study - a cluster randomized controlled effectiveness study. *Trials* [Internet]. 2010;11(1):95. Available from: <http://dx.doi.org/10.1186/1745-6215-11-95>
 38. Rogers ER, King SR. Intervention based on social cognitive theory: Evaluating adolescents' knowledge of {OTC} pain medications. *J Am Pharm Assoc* [Internet]. 2013;53(1):30–8. Available from: <http://www.sciencedirect.com/science/article/pii/S1544319115302843>
 39. Pawson R tilley n. *Realistic Evaluation*. 1997.
 40. Salter KL, Kothari A. Using realist evaluation to open the black box of knowledge translation: a state-of-the-art review. *Implement Sci* [Internet]. 2014;9(1):115. Available from: <http://dx.doi.org/10.1186/s13012-014-0115-y>
 41. Gillespie RJ, Harrison L, Mullan J. Medication management concerns of ethnic minority family caregivers of people living with dementia. *Dementia* [Internet]. 2015 Jan 1;14(1):47–62. Available from: <http://dem.sagepub.com/content/14/1/47.abstract>
 42. Zegwaard MI, Aartsen MJ, Grypdonck MHF, Cuijpers P. Differences in impact of long term caregiving for mentally ill older adults on the daily life of informal caregivers: a qualitative study. *BMC Psychiatry* [Internet]. 2013;13(1):1–9. Available from: <http://dx.doi.org/10.1186/1471-244X-13-103>
 43. Rycroft-Malone J, McCormack B, Hutchinson AM, DeCorby K, Bucknall TK, Kent B, et al. Realist synthesis: illustrating the method for implementation research. *Implement Sci* [Internet]. 2012;7(1):33. Available from: <http://dx.doi.org/10.1186/1748-5908-7-33>