

Effectiveness and Cost-effectiveness of 'Usual Care' versus 'Specialist Integrated Care': A Comparative Study of Hospital Discharge Arrangements for Homeless People in England



THE LONDON SCHOOL
OF ECONOMICS AND
POLITICAL SCIENCE



Barts and The London
School of Medicine and Dentistry

RESEARCH PROTOCOL

SUMMARY OF RESEARCH

The overall aim of this two year study is to establish firstly the ways in which specialist integrated homeless health and care (SIHHC) services are being developed and implemented to facilitate hospital discharge in England and, secondly, the impact this is having on quality of care and organisational outcomes such as the prevention of readmission to hospital.

The study will evaluate the principal models of SIHHC developed as part of the Department of Health's Section 64 (voluntary sector led) '*ten million pound cash boost*' (DH, 2013) to improve hospital discharge for homeless people, alongside some of the more longer established approaches such as the Pathway model. The main purpose of the study is not to make recommendations as regard one particular model being more effective than the other, but to understand the different components of each model in terms of what works to secure safe, timely and compassionate discharge from hospital.

In terms of what is already known about SIHHC, it is recognised that there is a fine balance between promoting access by providing specialist services and creating the conditions for further marginalisation and reinforcement of stereotypes. In some areas it is possible that a well developed hospital discharge protocol is able to secure timely discharge for homeless people without the need for SIHHC. To enable exploration of this, hospitals with access to SIHHC will also be compared to those without access to such schemes.

This is an important consideration as a key implication of the Francis (2013) inquiry into the abuse and neglect at Mid Staffordshire NHS Foundation Trust is that commissioners and managers will need to pay close attention to the scope for challenge and innovation in routine 'standard care'. By educating hospital staff about the needs of homeless people, SIHHC may be a means of achieving this but little is known about its effectiveness in this area.

Following a literature review, in-depth fieldwork employing realist evaluation will be carried out in six case study sites across England. For the purposes of this study, sites are understood as 'whole systems' of care in which the hospital is the focal point. Reflecting the principal typological approaches, sites will be purposively selected to include hospitals that have access to clinically led models of SIHHC (n=2), those that have access to housing link worker schemes with no clinical in-pat (n=2) and those which rely on standard provision (n=2). In selecting case studies, particular attention will be paid to different aspirations for integration encompassed in service specifications and discharge protocols, showing in the final analysis if and how this impacts on a wide range of outcomes.

In each site, fieldwork will comprise: interviews with commissioners and service managers from SIHHC and a range of other agencies ([minimum] n=5 per site); interviews with discharge coordinators, case managers and key workers from SIHHC and a range of other agencies ([minimum] n=5 per site); and interviews with service users (n=10 per site) shortly after discharge from hospital and at three months follow-up. In addition to capturing information on service use, the interviews with service users will ascertain experiences of the hospital discharge process, if 'joined-up' care was delivered and overall reflections on the health and well-being outcomes that have been achieved. This qualitative data will then be complimented with quantitative evidence on the costs and outcomes of SIHHC: In each of the six fieldwork sites we will collect and analyse EQ5D data for 63 service users (378 in total). Extending the study to a further 14 sites (making 20 sites in total) we will link information about 'service users' (e.g. NHS number) to centrally held health and public service data. This will reveal outcomes linked to hospital readmissions and use of emergency services

Key outcomes of the research will include in-depth understanding of the how different mechanisms are working to address poor discharge, with a particular focus on integration. From this, the study will provide new insights that policymakers and commissioners will be able to utilise to make recommendations and effect changes in practice. Finally, we will produce a toolkit for commissioners about developing SIHHC if the findings support this.

AIMS AND OBJECTIVES

The overall aim of the study is to explore how SIHHC is working to secure safe, timely and compassionate discharge from hospital. The objectives of the study are:

1. To situate what is already known about delayed discharge among homeless people in terms of the broader literature on hospital discharge, intermediate care and reablement.
2. To explore how different models of SIHHC are being developed and implemented across England to facilitate effective hospital discharge
3. To explore how SIHHC impacts on outcomes and different patterns of service use across the whole system (e.g. the use of unplanned care) and the associated cost implications of this.
4. To explore homeless people's views and experiences of whether, and if so how, SIHHC works to improve experiences of hospital discharge and to deliver improved health and well-being outcomes
5. To produce a 'toolkit' for commissioners on developing SIHHC (if the findings support this)

BACKGROUND AND RATIONALE

What is already known about hospital discharge and homelessness?

The research literature on hospital discharge goes back at least 40 years and there is remarkable consistency in its findings, which continue to report the frequent breakdown of routine discharge arrangements (DH, 2003). Older people and people with multiple and complex needs (long-term conditions) including homeless people are particularly affected by poor discharge arrangements, often leading to poor health and well-being outcomes and high readmission rates (Tansley and Gray, 2009).

For older people, policy initiatives around hospital discharge have focused mainly on the problem of delayed discharge, whereby people are deemed 'medically fit' but unable to leave hospital because they are waiting for a community care assessment or placement. In England, this problem was addressed through two key strategies. First, the development of intermediate care, which provides integrated support to facilitate timely discharge and a period of rehabilitation to prevent readmission. Second, a reimbursement policy, which allows the NHS to impose a financial penalty on local authorities for inappropriate occupation of NHS beds. The number of delayed discharges (across all service user groups) reduced dramatically from 2001 onwards. However, evaluations found it difficult to untangle the impact of these different initiatives. Key learning from the extant literature on delayed discharge suggests that there is a need to extend the policy, resource and service lens from a primary

focus on acute hospital entry and exit points to the wider health and social care economy in order to develop a broader repertoire of rehabilitation, intermediate and transitional care provision as well as long-term community infrastructure. According to Godfrey et al. (2008 p10), this requires 'whole system working and joint investments, encompassing not only marginal additional growth money but also shared decision-making around mainstream budgets'.

For homeless people, length of stay in hospital is reported to be three times longer than for other service users (Hewett, Halligan and Boyce, 2012). However, this is often attributed to the complexity of homeless people's presenting health needs (multimorbidity) rather than to delayed discharge. In other words, 'homeless service users' stay three times as long as they are [three] times as sick' (Hewett and Halligan, 2010, p306). Despite this, there is strong evidence that a key problem facing homeless people in hospital is 'poor care' which often leads to difficulties accessing appropriate treatment, early 'self-discharge' and being discharged prior to being 'medically fit'. A review of hospital admission and discharge arrangements by Homeless Link (2012) identified 'countless examples' of the mention of homelessness on admission triggering prejudice among hospital staff. According to one homeless interviewee, 'As soon as [the nursing staff] find out you're homeless they want you out of the door' (p7). The review reported that more than 70% of homeless people were 'discharged straight back onto the streets, further damaging their health and all but guaranteeing their readmission' (p6). The main recommendation of the Homeless Link report is for cultural change in the way homeless people are viewed and treated in the NHS.

Other themes highlighted in the Homeless Link review and the literature more broadly include: hospital staff not asking people about their housing situation on admission and not addressing needs relating to drug and alcohol use (e.g. ensuring access to methadone); hospital staff not alerting accommodation providers to the fact that a person is in hospital leading to tenancies being terminated; hospital staff having a poor understanding of the services available to homeless people, including overestimating the level of care that can be provided in hostel settings leading to unsafe discharges; a lack of practical support, such as no replacement of soiled clothing; uncertainty regarding where the involvement of one sector should start and end, including little agreement about how far hospital staff should be expected to proactively seek accommodation; a lack of appropriate accommodation options raising the question of whether 'signposting' is an adequate response if this potentially leads to discharge back onto the streets

The case for specialist health and care services for homeless people is often justified on the grounds that their needs are so complex as to be beyond the scope of routine services (Gould, 2009). Long-term homelessness is characterised by 'tri-morbidity' (the combination of mental ill health, physical ill-health, and drug and alcohol misuse) (Hewett, Halligan and Boyce, 2012) and compounded through multiple exclusion (McDonagh, 2011). In terms of tackling health inequalities, the Marmot review proposes a system of 'proportionate universalism'; that is, helping all sections of society but with a scale and intensity that is proportionate to the level of disadvantage. For homeless people, this provides an opportunity to improve outcomes by targeted investment, with the very real prospect of reducing unscheduled expenditure in secondary care (Hewett and Halligan, 2010).

Specialist schemes are not, however, without limitations. Daly (2007) argues that there is a fine balance to be drawn between promoting access by providing specialist services and creating the conditions for further marginalisation and reinforcement of stereotypes. Recently, the Francis (2013) report into abuse and neglect at Mid Staffordshire NHS Foundation Trust leads us to question the value of 'specialist care' where it works to compensate for poor (discharge) practices in routine 'standard' care. Indeed, one of the key recommendations of the Francis inquiry is that managers and staff should challenge poor practice at the time they see it. Responding to Francis, Dixon-Woods et al. (2013) emphasize the importance of innovation in routine 'standard care' as the key to safe compassionate care and of providing staff with the means to achieve this. They see the need for a shift from 'rules and regulations' to cultures of reflection and constant learning.

'If the NHS can work well for homeless people, then it will certainly work well for the rest of us' (Hewett, 2013, p3).

Current picture of 'standard' versus 'specialist' provision in hospital discharge arrangements for homeless people

DH guidance stipulates that lead managers for hospital discharge in acute hospitals and local authority adult services should ensure that their hospital discharge policy includes guidance for staff dealing with individuals who are homeless and aged both under and over 65 years. Moreover, the protocol should ensure that homeless people are identified on admission and their pending discharge notified to relevant primary health care and homeless service providers (DH, 2003 p74-75). Evidence suggests that this is far from being implemented across England with only 39% of local authorities reporting that they have formal protocols in place for the admission and discharge of people who are homeless (Homeless Link, 2010).

In terms of conceptualising 'standard' versus 'specialist' discharge arrangements, the following typology - developed by Homeless Link as part of their aforementioned review - is a helpful starting point (2012, p5):

- *Specialist* - formal protocol involving a specialist post or service, funded either by the local authority or co-funded with the NHS
- *Standard* - formal protocol, without specialist provision, but a clear process in place with the hospital, often using agreed referral forms
- *Standard (sub-optimal)* - no formal policy in place and no specialist provision

Where protocols for homeless people's admission and discharge are in place Homeless Link (2012) report that these vary from area to area. In some areas, local authority staff visit service users on the ward to make an assessment, start the referral process to an accommodation provider and offer advice. Nonetheless, the extent to which hospital staff stay engaged in this process varies. Some alert housing agencies and then deem that to be the end of their role. Others stay involved to organise a case conference with the potential of delivering a more integrated or coordinated response. Importantly, 'good practice' or 'bright spots' were not seen as being restricted to hospitals with 'specialist provision' (p20). Even in areas with no discharge protocol, some good practice was recorded whereby community based staff were advocating successfully on the part of homeless people to support them through the hospital 'journey'.

Where formal protocols (without specialist provision) seemed to have driven improvements, a number of factors appear relevant. These include effective joint working at both operational and strategic levels, good communication between agencies, the provision of training (especially induction) and considerable efforts to keep protocols 'live' (e.g. regular meetings, phone calls and personal contacts between partners) (Homeless Link 2012 p10). In a number of areas visited by Homeless Link, 'specialist provision' appeared to have evolved from protocols that had started out as cross-agency referral agreements. In York, for example, hospital staff would refer homeless service users to the Salvation Army who would then provide support and advocacy, such as organising the housing support related side of the discharge including ensuring that hostel staff were briefed on the person's health needs. This bridging role led the Salvation Army to develop a visible presence in the hospital ('in-reach') and to provide training to hospital staff on the needs of homeless people (reported in Homeless Link, 2012 p8).

In London and inner city areas with higher populations of homeless people a number of specialist schemes have been developed which are GP or nurse (clinically) led. A number of different models fall within this typology. For example, in the Royal Liverpool University Hospital (RLUH) a nurse-led team is responsible for the discharge of service users who are homeless. The team comprises a 'link nurse', a substance misuse nurse, an adult safeguarding lead, and three hospital outreach workers providing 27/4 housing and welfare benefits support to homeless people. The RLUH team then link homeless people into the vulnerable persons' team at a specialist primary care centre (Brownlow) which includes access to a GP, two homelessness nurses, an alcohol outreach nurse and a Hepatitis C nurse.

Another high profile (award winning) 'clinically led' model is the Pathway approach. This approach was pioneered at University College Hospitals London (UCH) and has since been adopted across a further three London hospital sites as well as in Bradford, Leeds, Manchester and Brighton. The

Pathway approach comprises a hospital based team which is GP and nurse led. For example, the newly established Pathway Team, which works across the King's Health Partnership (KHP) hospitals (Guys and St Thomas, King's College Hospital and [to soon include] South London and the Maudsley) involves: six Homeless Health Practitioners (two general nurses and four Mental Health Practitioners); ten GP sessions per week, provided by two GP's contracted for 52 weeks per year from a local practice or urgent care centre provider; a full time social worker (employed by the hospital not the local authority); a full time housing specialist contracted from a third sector provider; a half time, band 4 secretary for administrative support; and specialist legal support from the KHP in-house legal team as necessary.

Peer support from Care Navigators is also a key component of the Pathway approach. The hospital based teams co-ordinate care for service users from hospital admission through to 'independence' (defined as 'putting service users in touch with follow-on services'). Key steps include nurse-led assessments, weekly homeless 'paper based' ward rounds and case conferences leading to multi-agency care plans. The Pathway service is structured to provide support to Accident and Emergency department staff in managing homelessness and 'frequent attenders'. Specialist intermediate care or 'medical respite' ('step down') facilities are also being developed to provide opportunities for convalescence and recuperative care.

The total annual cost of the King's Health Partners Pathway service is estimated to be £675,480. This is financed through both NHS and charitable sources. It is acknowledged that the Pathway approach is expensive and unlikely to be cost effective where there are less than 200 homeless or 'tri-morbid' admissions per annum (Pathway Annual Review, 2013, p4). As noted above, a number of areas have begun to implement the model (what has been termed 'Pathway Lite') adapting it to fit with local circumstances and available resources.

In 2013, the Department of Health announced a £10 million grant fund as part of a 'Homeless Health Discharge' initiative targeted at further developing partnerships between the voluntary sector, the NHS and local government. Fifty-two voluntary sector led schemes¹ have since been funded. An initial evaluation of the start-up phase by Homeless Link (2015) suggests the majority of the schemes developed with this funding are 'Housing Link Worker' projects which do not include any direct clinical in-put.

Link workers with a housing background aim to bridge the gap between hospitals and homelessness accommodation, and may have referral rights into local hostels or temporary accommodation. Where possible, they assess a client while in hospital and then refer the client to appropriate accommodation and other support services. Building up relationships with relevant organisations, accessing housing and signposting clients are key aspects of this role...

(Homeless Link, 2015, p12)

A much smaller number of the schemes are 'clinically led' employing Nurse Link Workers who can provide some direct health in-put or are 'mixed' employing both housing and nursing link workers. The principal typologies identified by Homeless Link (reflecting how the DH funding was spent) are:

- Housing Link Workers (22 Schemes)
- Accommodation with Housing Link Workers (8 Schemes)
- Pathway Model (funding for navigators) (6 Schemes)
- Accommodation (including intermediate care and respite) (7 schemes)
- Housing and Nursing Link Workers (4 projects)
- Nursing Link Workers (2 projects)

An important feature to have emerged in some of the inner city areas is the use of this Section 64 funding to add extra capacity to already established schemes. Some areas thus have a **nest** of different schemes and typologies ideally working closely and seamlessly together:

¹ Section 64 grants can only be made to non-statutory organisations.

'Following a Needs Assessment delivered in 2012, Lambeth and Southwark CCGs kindly commissioned an initial pilot of the 'Kings Health Partners Pathway Homeless Team'. The Pathway Homeless Team is affiliated to, and forms part of the Pathway network of 'Homeless Ward Rounds' in acute care settings nationally.... The pilot team was multi-professional, and multi-agency, and involved 14 core staff. In addition to the team members directly funded by the CCGs, the team also benefited from working in close partnership with two hospital discharge projects funded by short-term Department of Health money. Therefore, 4 out of 5 of the involved Housing Support Workers were actually employed by St Mungo's Broadway and the Passage. The team also benefited from the input of a Groundswell Peer Advocate and Groundswell volunteers. This input was also made possible by short-term Department of Health funding'.

Extract from the King's Health Partners
Pathway Homeless Team Pilot Project Report, March 2014

What is already known about the effectiveness and cost effectiveness of standard versus specialist hospital discharge provision?

Notably, few of the sites visited as part of Homeless Link's (2012) review were recording any formal data on the impact of the various approaches that had been introduced to improve hospital admission and discharge for homeless people. To address this oversight, Homeless Link funded an economic evaluation of a hospital discharge project in Wirral which employed a 'link worker' to train staff and undertake some direct case work to support homeless service users' discharge (White, 2011). The evaluation included a comparative analysis of the data for homeless service users, using Hospital Episode Statistics, in the year prior to and during the introduction of the link worker. This showed very positive indicative results in terms of cost savings (a reduction of around £26,500 in the total cost of episodes linked to service users with 'No Fixed Abode'). There was also a fall of one third in the number of episodes resulting in emergency readmissions within less than 28 days.

A randomized controlled clinical trial of the costs and benefits of the UCH and Brighton Pathway services has recently been completed by members of this research team [Hewett & Foster] (Hewett et al., submitted). In the trial, homeless service users admitted to hospital were randomly allocated 'standard' care or 'enhanced' care with input from the Pathway team (as described above). Service users assigned 'standard care' were visited once by the Pathway homeless health nurse. They and their named ward nurse were provided with an information leaflet describing local services for people who were homeless, but the Pathway team played no further role in their management. Health care usage was determined using hospital system data whilst quality of life and accommodation were assessed by questionnaires at admission and six weeks after discharge.

Results indicated that the Pathway approach does not alter length of stay but improves the quality of life of homeless people (with fewer service users returning to the streets) at a cost deemed cost effective under current guidelines.² In part, this was an unexpected finding since an earlier review had suggested the UCH Pathway was able to achieve a reduction of 1,000 bed days (30% reduction) in the first full year of service delivery (Hewett, Haligan and Boyce, 2012). The trial findings were, however, able to allay concerns that the introduction of the Pathway approach might increase overall length of stay by delaying discharges. An additional important finding was that improvement in housing status occurring during the trial did not manifest as a reduction in the use of health care services; although longer durations of follow up may change this picture. It was further noted that quality of life benefits are hard to demonstrate in an intervention where the short duration of follow up limits the accrual of benefits.

Integration

A consistent theme in the Coalition government's policy on homelessness is a belief that support services could be improved if agencies worked together more effectively (DCLG, 2011). This is also implicit in the DH's (2013) 'Homeless Health Discharge' initiative which aims to develop partnership working between the voluntary sector, the NHS and local government. A particular concern of the Commissioning Brief (NIHR 13/156) is to understand how these Section 64 schemes are working,

² £26,000 per Quality Adjusted Life Year (QALY).

with a particular focus on integration.

The evidence base for partnership working and integration across all sectors is weak and it has been argued that government exhortations to reduce public funding because joint working/integration can improve the effectiveness of services while also delivering cost savings may not be justified (Petch, 2012). In a recent systematic review, Cameron and Lart (2013) have argued that the evidence base for integrated care is underdeveloped because few studies have adopted a comparative design, so differences between 'standard' care and 'integrated' care are not assessed.

Drawing comparisons between 'integrated' and 'standard' care is methodologically fraught not least because there is a lack of conceptual clarity regarding what constitutes integrated care (Leathard, 2003). Additionally, there are many reports of integrated services (e.g. a co-located multi-disciplinary team) failing to deliver integrated care (e.g. where the service user continues to have multiple assessments and support plans in spite of co-location and multi-disciplinary team working) (Cornes et al., 2001).

Petch (2012) suggested that one strategy for overcoming this is to consider partnership working as the process and integration as a potential outcome. Partnership working might also be considered in terms of a continuum incorporating relative autonomy, co-ordination, joint appointments, enhanced partnership and structural integration. In considering partnership process, it is also important to consider the number and reach of partners, resources and depth of activity, from sharing information at one end to formal merger on the other. For example, co-ordination is evident in the St Petroc's scheme and more structural approaches exist in the Pathway model where there is both vertical integration (of primary and secondary care) and horizontal integration (across health, housing and social care).

Finding 'pure' comparator sites (i.e. hospital sites with no integration) may, in fact, be practically impossible given the raft of initiatives targeted at 'standard' discharge practices, most notably the mainstreaming of intermediate care and later reablement. In consequence, it seems more helpful to construct the comparative focus of this study around the differences between 'specialist integrated care' versus 'standard integrated care', recognising that different degrees of partnership working and integration may be 'designed for' and achieved across each of these typologies with associated differences in outcomes. Outcomes linked to continuity of care are perhaps the most important marker of effective integration. Thus, it will be important to assess the full range of possible 'transitions' in any given locality and how homeless service users move across sectors (health, housing and social care) and between 'specialist' and 'standard' care, and indeed, between different forms of 'specialist' care.

As noted above, voluntary sector projects in the vicinity of both Liverpool and King's hospitals have received some of the aforementioned DH Section 64 funding to develop 'specialist discharge schemes'. These are envisaged as working in partnership with the schemes that are already established. Unless these relationships and transitions are carefully managed, there is a danger that homeless people could experience a thicket of un-coordinated short term support (multiple 'sequential handovers') in which everybody has snippets of the individual but no one agency co-ordinates the overall picture. In their study of integrated working around multiple exclusion homelessness, Cornes et al., (2010) concluded that while short term specialist support does have a role to play, it should not detract attention from the need to develop higher level strategic measures which deliver personalised and integrated case management over the longer term for all adults (e.g. a 'Common Assessment Framework').

Summary

Our review of current UK provision (based on a preliminary literature search and desk top review) suggests that SIHHC discharge schemes comprise approaches which are either clinically led (such as the Pathway approach) or those that rely on housing link workers with no direct clinical input. Specialist schemes have access to many different types of resources and disciplinary expertise and are underpinned by different protocols for partnership and integrated working. Nearly all of the schemes encompass some form of housing 'in-reach' (statutory or voluntary sector support to secure accommodation on discharge). There is also a growing trend towards the development of

intermediate/reablement care which provides access to either 'temporary accommodation' or 'medical respite'.

In line with the HS&DR call we wish to generate new evidence about how these schemes are working on the ground. While there is some evidence about the scope and effectiveness of particular schemes, this brief review highlights that there are also many unknowns for commissioners. It has been suggested, for example, that 'GPs are the only health professionals trained to deal with physical ill-health, mental ill health and drug and alcohol issues in a social and psychological context' (Hewett, 2010, p6). This raises the question of how specialist discharge schemes which are clinically led compare to those that are not.

We would further argue that the comparison between 'specialist' integrated care and 'standard' integrated care is critical because both '*projects*' and '*protocols*' are potentially key components and mechanisms for improving care (discharge arrangements). As noted above, Homeless Link (2012) did not see 'good practice' as being restricted to areas with specialist provision. Post Francis, understanding what might best support innovation and integration in routine mainstream care is seen as an urgent priority for the NHS and its partners. It is important to know if SIHHC supports or detracts from this.

RESEARCH PLAN & METHODOLOGY

Overview of Research Plan

This research comprises two inter-linked work packages. The first is a realistic evaluation to be carried out across six case study sites across England. It is designed to explore: (Objective 2) how different models of SIHHC are being developed and implemented to facilitate effective hospital discharge and; (Objective 4) homeless people's views and experiences of whether, and if so how, SIHHC works to improve experiences of hospital discharge and to deliver improved health and well-being outcomes. The second is a data linkage and health economic analysis work package targeting twenty sites across England with specialist hospital discharge schemes. This is designed to explore (Objective 3) how SIHHC impacts on outcomes and different patterns of service use across the whole system (e.g. the use of unplanned care) and the associated cost implications of this. Both work packages will be underpinned by a literature review (Objective 1) and will inform the development of a commissioning 'tool kit' (Objective 5).

Research Design (Theoretical and Conceptual Framework)

The design of this study has been informed by guidance on developing and evaluating complex interventions in both health and social care (Evans et al, 2013). The guidance takes the researcher through a staged process of developing, evaluating and implementing a complex intervention in which multiple structures, people and processes are involved, notably the delivery of a service. Best practice is to develop interventions systematically, using the best available evidence and appropriate theory, then to test them using a carefully phased approach, starting with a series of pilot studies targeted at each of the key uncertainties in the design, and moving on to an exploratory and then definitive evaluation. As outlined above, the Pathway model has been developed in a carefully planned and staged way leading to definitive evaluation (RCT). In terms of the 'Continuum of Evidence' (p8) framework developed as part of the complex intervention guidance the next stage is to determine if others can replicate the intervention and results in uncontrolled settings. That is, 'To roll out and monitor results of wide implementation' (p14)

However, this linear view of service development does not always capture the complexities of local commissioning where an evidence based approach may be lacking (Huxley et al., 2010). Acknowledging the consistent failure to apply effective interventions in practice, the complex

intervention guidance notes the way in which evaluation may need 'to move back and forth between the development and feasibility phases' (p16). In terms of the design of this study, the roll out of 'Pathway Lite', the proliferation of new voluntary sector led predominantly housing link support schemes along with seeming inaction in many areas (where protocols for the discharge of homeless people have yet to be developed) points to the need to 'move back' and to provide a more nuanced understanding of each of these developments.

In making sense of the proliferation and diversity of provision which now surrounds hospital discharge arrangements for homeless people in England, a key concern in designing this study, is that we should not position 'models' comparatively so that they are in competition with one another. An important lesson to emerge from the national evaluations of intermediate care was that it proved impossible to arrive at recommendations as regards any particular model or service configuration being more or less effective and cost effective than the other (Barton et al., 2006).

Rather we see more practicable value for commissioning, in shifting the evaluative 'gaze' away from potentially replicable *whole* 'models' or 'interventions' towards a greater understanding of underpinning processes and practices. As in the case of Pathway Lite, commissioners will often 'mix and match' components of different interventions to ensure fit with local circumstances and resources. Indeed, while the findings of the Pathway evaluation provide what is often considered to be definitive ('gold standard') evidence, it is not clear from the trial which components and mechanisms delivered which results. For example, did the approach work because of 'integration' or because it was 'GP led'? Or were there other mechanisms at play that were not specifically 'designed for', such as workers having the opportunity to 'step out' of the mainstream services and to work in an unusually well-resourced team with good levels of management and charismatic leadership? Arguably, the RCT is also limited by the fact that comparisons were made against what might be assessed as 'sub-optimal' or 'poor care' (i.e. just giving the service user a leaflet) rather than against 'standard care' where there are fully developed discharge protocols in line with DH guidance.

In research design terms, 'realistic evaluation' (Work Package 1) has proved effective in generating evidence about how different components and underlying mechanisms can generate impact in different circumstances (Byng, Norman and Refern, 2005; Wilson and McCormack, 2006; Manzano-Santaella, 2011; and Pearson et al., 2015). The fundamental question that researchers working with a critical realist perspective ask, is how do certain causal mechanisms (e.g. GP led advocacy), operating in particular circumstances (e.g. a SIHHC scheme), create certain changes or outcomes (e.g. improved experiences of hospital discharge)? The overall aim is to generate a 'road map' or 'tool kit' for local commissioners and providers on how to translate these mechanisms effectively into their own contexts and systems of care.

The data linkage and economic evaluation (Work Package 2) is then designed to compliment this emergent and nuanced picture of ground level practice developments with more quantitative generalizable evidence on the costs and outcomes linked to the provision of 'specialist' arrangements for hospital discharge. Here, a cohort of homeless people who have used a specialist hospital discharge scheme will be compared to a cohort of homeless people who have not used such provision.

Plan of Investigation and Methodology for Delivering Case Studies

The study will comprise the following stages: [1] Preparation (research team building; devising project delivery plan; establishing advisory groups; training peer researchers; hosting an expert seminar and securing ethical permissions); [2] Literature Review; [3a] Delivery of Work Package One [3b] Delivery of Work Package Two; [4] Dissemination (preparing conference papers/articles) and developing a 'tool kit' for commissioners).

Stage 1: Preparation (including PPI)

Ethical approval and research governance will be sought from the National Research Ethics Service, and R&D from local NHS Foundations Trusts/Clinical Commissioning Groups and local authorities

(where appropriate). Please see below and in main application form for further discussion of ethical considerations.

The preparation phase will also involve an establishing an Advisory Group. The research has already secured support from a wide range of key stakeholders who are listed as collaborators. This includes representatives from research, practice, and stakeholder bodies such as Homeless Link and Revolving Doors.

'Experts by Experience' (people with direct personal experience of homelessness) have already been involved in all stages of developing the study proposal and will be central to the successful delivery of the research if funded. The idea for the project was first discussed at a meeting involving Revolving Doors' Peer Research Network. Two 'experts by experience' have since joined the team as collaborators. Both Alan and Natalie have peer research experience, have contributed extensively to electronic and face-to-face discussions during the preparation of the full application for this study and will be members of the main Advisory Group if the project is funded. Further details of PPI (the 'peer research' component) are outlined later in this document.

As part of the preliminary stage, we also plan to host an 'expert seminar' to ascertain a picture of the latest developments and ideas surrounding SIHHC, and as noted above, as a way of securing expressions of interest in becoming our research partners.

Stage 2: Literature Review

A narrative literature review will be undertaken to assess what is already known about hospital discharge and homelessness (Objective 1). This will be contextualised in terms of findings from key systematic reviews of (generic) delayed discharge, intermediate care and reablement literature. We will conduct an electronic literature search for peer-reviewed articles published in English from 2000 - 2014 in the CINAHL, NHR CRN Portfolio, PsycINFO and MEDLINE databases. We will use keywords such as: homeless, hospital discharge, integrated care (full range of relevant terminology multidisciplinary, interprofessional etc.). Potentially relevant articles will be analyzed by two experienced narrative reviewers. Further to this, we will also undertake a grey literature search through relevant websites (e.g. Department of Health; Homeless Link), as well as through the internet using the Google search engine.

The literature review will generate a more complete typology of the 'resource components' which have been used or could be used within SIHHC, along with any evidence for effectiveness. Qualitative studies will be used to generate preliminary theories about how underlying mechanisms within the resource components generate benefit.

Stage 3a: Delivery of Work Package One - Realistic Evaluation

Overview

For the purposes of the realistic evaluation we are proposing to generate six case studies. Four of the host sites will have access to a 'specialist' hospital discharge scheme and two of the sites will have 'standard' provision only (i.e. no access to specialist support around the discharge of homeless people from hospital).

Site Selection

Sites with 'specialist provision' will be targeted purposively to reflect the principle typologies described above and the potentially important distinctions that are to be made between those schemes (1) which have direct access to clinical input and leadership those which do not; and (2) those schemes which have direct access to accommodation and those which do not. As shown in Figure One below sites will be selected to reflect:

(Specialist Discharge Schemes with Direct Clinical Leadership and In-put)

Site One – Pathway (GP led with clinical in-put)

Site Two – Nursing Link Worker (nurse led with clinical in-put)

(Specialist Discharge Schemes with no Direct Clinical In-put)

Site Three – Housing Link Worker (no clinical in-put)

Site Four – Accommodation and Housing Link Worker (no clinical in-put)

(Standard Provision)

Site Five – No specialist hospital discharge scheme – discharge protocol only

Site Six – No specialist hospital discharge scheme – no discharge protocol

Figure 1: Definition of Specialist/Standard/Sub-optimal Hospital Discharge Arrangements for Homeless People (Homeless Link, 2012)	Site 1 Inner London	Site 2 Inner City	Site 3 Metro/ Rural	Site 4 Metro/ Rural	Site5	Site 6
Specialist (Enhanced DH Policy Guidance): Formal protocol around discharge arrangements for homeless people involving specialist post or service – funded either by the DH (Section 64) local authority or co-funded with the NHS	GP Led	Nurse Led	Housing Link Worker	Accommodation with Housing Link Worker		
Standard (DH Policy Guidance Implemented): Formal protocol around discharge arrangements for homeless people, without specialist provision					standard	
Standard (Sub-optimal) (DH Policy Guidance Not Implemented): No formal protocol in place as regards discharge arrangements for homeless people						standard

Typologies linked to '*mixed*' (combined housing and nursing link worker) and intermediate respite care will be considered in Sites 1 and 2 which will purposively targeted to additionally reflect the '*nesting*' of multiple discharge schemes. These will most likely be located in the inner London and inner city areas to reflect the potentially higher level of resource allocation needed to meet the needs of larger populations of homeless people. Sites Three and Four will target areas with a single discharge scheme located in metropolitan/rural areas. It is perhaps worth stressing here, that the qualitative phase of this (realist) evaluation does not aim to compare these different typologies 'like for like' but to generate multiple '*context, mechanism and outcome*' (CMOs) configurations to inform discussion around what works best for whom and in what circumstances. So for example, exploration of different CMOs may help to answer key commissioning questions such as in a rural area with limited resources and a small number of homeless people is it more advisable to have a 'housing link worker' or a 'nursing link worker' and what is the value of direct access to accommodation? For illustrative purposes (based on the preliminary typological review outlined above) Figure 2 charts some of the different resources and impact generating mechanisms we might expect to find attached to each purposively selected category of scheme.

Figure 2: Resources and Potential Impact Generating Mechanisms Attached to SIHHC Schemes	Site 1 e.g. London Pathway	Site 2 e.g. Liverpool Royal Nurse led	Site 3	Site 4	Site5	Site 6
Provision of specialist training for hospital staff on the needs of homeless people		specialist			N/A	N/A
Specialist case management with professionally qualified case manager (e.g. GP/nurse/social worker)	specialist	specialist			N/A	N/A
Specialist key working (navigation and brokerage) led by non-professionally qualified staff (or peer mentors)	specialist	specialist	Specialist Housing link worker	Specialist Housing link worker	N/A	N/A
Arrangements for single shared assessment/information sharing/integrated support planning.	specialist				N/A	N/A
Weekly multi-disciplinary team meetings/case conferences	specialist				N/A	N/A
Arrangements for review/follow-up post discharge (e.g. support to make sure care continues after discharge)	specialist	specialist			N/A	N/A
Availability of intermediate care/medical respite beds (dedicated homeless service, beds)	planned				N/A	N/A
Access to emergency/temporary accommodation on discharge	specialist - the passage	specialist		specialist	N/A	N/A

The final consideration in selecting the case studies is that we wish to be able to study different aspirations for integration within SIHHC, showing in the final analysis if and how this impacts on a wide range of outcomes. This will be guided by Figure 3. Again, for illustrative purposes, Site 1 is shown as the inner London Pathway model where a high degree of integration is known to have been achieved, with lesser but substantial integration being achieved in the Liverpool model. In Sites 3 and 4 we anticipate 'Housing Link Work' to involve primarily enhanced partnership working between the hospital and local homeless services and agencies.

Figure 3: Protocols for Partnership and Integration as Envisaged in Commissioning/Service Delivery Specifications (Anticipated Degree of Integration)	Site 1 e.g. London Pathway	Site 2 e.g. Liverpool Royal Nurse led	Site 3	Site 4	Site5	Site 6
(Structural) Vertical Integration (e.g. integration of primary/secondary Care)	specialist					
(Structural) Horizontal Integration (e.g. integration across health, housing, social care)(co-location/multi-disciplinary team based in hospital)	specialist	specialist				
Enhanced Partnership (e.g. arrangements for in-reach/outreach)	specialist	specialist	specialist	specialist		
Joint Appointments (e.g. social worker seconded to GP led scheme)	specialist					
Co-ordination (e.g. shared 'referral forms')					standard	
Autonomy						standard
<i>Extent of horizontal integration – partnership arrangements between three or more disciplines are in operation (e.g. GPs, nurses, social workers, housing support workers)</i>	specialist	specialist				

Site Recruitment

In order to recruit the case study sites, we will identify and write to all schemes fitting the criteria outlined above informing them about the study and seeking expressions of interest to become our research partners. To boost potential interest and to prevent against research fatigue we will also host an expert seminar in the first month of the research. We will host this in partnership with Homeless Link and we will invite representatives from the longer established and newer Section 64 Hospital Discharge Schemes. We will use the event to feedback some of the early findings from other studies and Homeless Link's evaluations and then to invite interest in the new study. Once we have ascertained which schemes would be interested in becoming a research partners for the in-depth qualitative phase of this study, we will then carry out some preliminary information gathering to fully describe these projects (e.g. costs, staffing, difference in practice approaches and the different resource and component parts). Figures 2 and 3 will act as a guide to inform final site selection so that the case studies present the opportunity to study many different context, mechanism, outcome (CMO) configurations. As regards Sites 5 and 6 research will needed to establish which areas have no protocol and no specialist provision (these sites are known to Homeless Link regional officers).

To date we have held preliminary discussions with a number of schemes who have already expressed a keen interest to take part. We have also met with the National Housing Federation as regards the Housing Link Worker schemes.

Fieldwork³ in Six Case Study Sites Across England

Each case study might be usefully thought of as a 'mini-ethnography'. Sites are understood as 'whole systems' of care in which the hospital is the focal point. The aims are to situate each of the SIHHC projects (or protocols or practices in standard care) in their wider historical and geographical **contexts**; understanding how they fit as a 'jigsaw piece' in a wider whole system of care; exploring their theoretical underpinnings and 'programme logic' (in particular, unearthing what projects define as their 'Key Practice Principles [KPPs]) and testing out these '*uncertainties*' or hypotheses by way of locally tailored evaluations which investigate the **outcomes** that are being achieved.

In the language of realistic evaluation, KPPs are '**mechanisms**', but we chose to use these terms as we find they are more user friendly when carrying out fieldwork. Already, KPPs can be discerned from our background review of SIHHC, for example, the idea that training provided through specialist homeless workers can deliver more compassion in mainstream care. Another example, taken from the Pathway model, suggested that 'advocates' need to be of sufficiently 'high professional standing' (GP led) because of the enormity of the barriers facing homeless people in accessing social care and housing related resources. In terms of KPPs a particular focus of this study will be on integration and how the projects are designed around this. For example, Pathway multi-disciplinary meetings are in keeping with a 'community of practice' ethos where, for example, even if a practitioner does not have direct involvement in a case he or she will still be asked for their opinion. The 'hypothesis' is that this will engender better service user care through more collegiate and interdisciplinary team working.

So as not to lose sight of what is the overarching research question (that is, what is the value of 'specialist integrated care' as compared to 'standard care') we have framed the proposed research conceptually as shown in Figure 5 (overleaf). This is to illustrate how the different Context-Mechanism-Outcomes (CMO) are linked to each site and to facilitate comparisons and contrasts. In this context, comparative analysis works principally in the anthropological sense to generate critical insight and reflexivity by enabling participants to know something of what is happening 'over the hill' (i.e. how other schemes are operating, their challenges and if and how they have been overcome). In terms of understanding scope for innovation, we are especially interested to know if KPPs 'cluster' significantly more in specialist projects rather than in standard care.

While a 'specialist hospital discharge scheme' might be conceptualised as mechanism for change in its own right, we situate it as 'context' within this particular interpretative framework to maintain the integrity of our underpinning philosophy. This is about the need to move beyond '**evidence based**' guidelines about single interventions or models towards a more '**evidence informed**' approach which is concerned more with encouraging reflection on and in practice and supporting staff and commissioners to challenge themselves and each other (Wilson and McCormack, 2006). As detailed below, we also include methods for engaging 'experts by experience' in this iterative process drawing on the work of Beresford (2007) who sees participatory research as having particular benefits in developing new knowledge about health inequalities.

³ The fieldwork will be carried out by Martin Whiteford (100% FTE) who will be responsible for four sites and Michelle Cornes (50% FTE) who will be responsible for two sites.

Figure 5: Overarching Research Framework for Understanding CMOs Across Specialist and Standard Care		
Context - Including an understanding of the wider whole system in which a specialist scheme or hospital protocol is situated	Specialist Integrated Care (X4) <i>(Note that a separate column would be generated for each site)</i>	Standard Care (X2) <i>(Note that a separate column would be generated for each site)</i>
Mechanism (s) including resources for facilitating safe discharge of homeless people from hospital including identification of any Key Practice Principles (KPP) encompassed within the broader programme logic. Underlying or hidden mechanisms are also important (e.g. practitioners “going the extra mile”)	KPPs: E.g. GP led advocacy overcomes the barriers facing homeless people in accessing community services. KPPs (Integration) E.g. Regular multi-disciplinary meetings using community of practice methodology	KPPs: E.g. To keep protocols live there is a need for robust induction processes for new staff . This obviates the need for specialist schemes and leads to more seamless services KPPs (Integration) E.g. Integration – special arrangements are in place with local supporting housing providers so that people in hospital are ‘fast tracked’ in the referral process
Outcomes – Understanding what it is that the ‘projects’ or ‘protocol’ is trying to achieve and if and how it has been achieved	Examples... <ul style="list-style-type: none"> Improved experiences of hospital discharge Improved health & well-being Compassionate Care Social inclusion benefits Cost of care – increased or decreased Cost of integrated care – opportunity costs Other (e.g. reducing hospital readmission rates) 	Examples... <ul style="list-style-type: none"> Improved experiences of hospital discharge Improved health and well-being Compassionate Care Social inclusion benefits Cost of care – increased or decreased Cost of integrated care – opportunity costs Other (e.g. reducing hospital readmission rates)

Because of the need for locally tailored evaluation in each case study site, some flexibility in the data collection strategy will be required. For example, focusing on mechanisms linked to GP led advocacy may require a different interview mix as compared to exploring mechanisms linked to specialist workers providing training to hospital staff. However, as a minimum the following data collection and analysis will be carried out in/for each site:

[3:1] Documentary Analysis: This stage involves collecting a wide range of background information about the sites (population, local homelessness provision, etc.). Service specifications (for specialist schemes) or ‘protocols’ for standard care will be collected and reviewed, as will any previous local evaluations/inspections.

[3:2] Interviews with commissioners and service managers from SIHHC and a range of agencies (minimum 5 per site). The focus of these interviews will be on the history and current conceptualisation of the ‘problem’ of homeless people’s discharge from hospital in that particular site; understanding what solutions have been developed locally (identifying any KPP), how they have been implemented, the costs associated with them and perceptions of how they are working. A standard typology pro-forma will be used to gather information on the different resource components that are available across the six sites.

[3:3] Observations of Practice and Interviews with discharge coordinators, case managers and key workers from SIHHC and other agencies (minimum 5 per site). Time will be spent observing the practices of the SIHHC schemes and standard hospital discharge practices working with homeless people. Focussing more on a practice rather than strategic perspective, these interviews will also cover the history and current conceptualisation of the ‘problem’ of homeless people’s discharge from hospital in that particular site; understanding what solutions have been developed

locally (identifying any KPPs), how they have been implemented and perceptions of how they are working.

[3:4] Interviews with service users (homeless people) (n=10 per site).

These interviews will be carried out shortly after discharge from hospital and then again in three months time. Where there is a period of 'step down' care a further interview will be carried out at the point of discharge. The purpose of these interviews is to find out about people's experiences of being admitted to (i.e. the events leading up to the admission) and discharged from hospital and whether SIHHC (and the mechanisms therein) or the protocols of standard hospital care had the intended impact. The interview guide will be tailored to ascertain feedback on any local KPPs but will be informed more broadly by Pearson's et al.'s (2006) realist systematic review of intermediate care. This provides a conceptualisation of what might be termed good practice or compassionate care in hospital discharge practice and provides an organising structure for a depth conversation. In this view, intermediate care is conceptualised as a short-term service arrangement which responds to a person's 'health crises' or acute hospital admission with⁴: (1) the objectives of care and place of care being negotiated between the service user and health and social care professionals; (2) health and social care professionals fostering the self-care skills of service users and shaping the social and physical environment to 're-enable' (promote recovery); and (3) service users, carers, health and social care professionals and voluntary services contributing actively to decision making and the delivery of that care is meaningfully integrated.

The main focus of the second follow-up interview will be on: (i) Patterns of service use in the intervening three months period, including any readmission to hospital (What support was received? Did the support materialize as planned? Did the support meet self-perceived needs?); (ii) What was liked and disliked about the support? (iii) How was the support perceived to have helped? What goals were achieved? What goals were still aspired to and what additional support might help to achieve those goals?

The timing of the second interview at three months is intended to give sufficient time to reveal potential for continuity (i.e. support beyond the 6-8 weeks cut off point often associated with intermediate care support) but not so far into the future as the connections with the initial hospital and discharge planning episode will likely be lost to memory.

We will also seek homeless people's written consent to access support plans and personal information which may be held by a range of providers.

The inclusion criterion for service users will be that they are experiencing 'multiple exclusion homelessness' (MEH). This criterion is chosen as it encompasses multimorbidity (a range of health issues) and compounding factors potentially impacting on well-being which are linked to experiences of 'deep social exclusion'. People experience MEH if they are 'homeless' (broadly defined, to include temporary/unsuitable accommodation as well as sleeping rough), and have also experienced one or more of the following domains of deep social exclusion –'institutional care' (prison, local authority care, stays in mental health hospitals or wards); 'substance misuse' (drug problems, alcohol problems, abuse of solvents, glue or gas); participation in street culture activities (begging, street drinking, 'survival' shop lifting or sex work) (Fitzpatrick et al. 2011 p.502).

Service users will be recruited on the basis that they are 'key informants' with regard to different experiences of MEH which might highlight different patterns of service user (including barriers). In each site, we will purposively recruit service users referred either to specialist schemes or in (standard care) to discharge coordinators/ward managers. Recruitment will aim to target the following groups/demographics and whenever possible we will approach people in the order to which they referred to the service:

⁴ Note that the definition of intermediate care used in the review is broad in scope and would encompass SIHHC not just medical respite/step-down accommodation

- Young Homeless People (1 male/ 1 female) (Age 18-24)
- Men in the middle age range (4 male) (Age 25-49) [Known to have the most complex needs (McDonagh, 2011)]
- Women in the middle age range (25-49)(2 Female)
- Older Homelessness (50+) (1 male/1 female)

If a service user (key informant) decides to drop out of the research or becomes unavailable we will recruit another participant to take their place. Participants will receive a £10 voucher (per interview) for sharing their expertise and giving-up their time. We will also employ 'peer researchers' (see below) to ensure engagement and support for the research locally among homeless people.

Peer Research

The Commissioning Brief suggests that this population poses particular problems for research in terms of participation and consent. This team has extensive experience of undertaking participatory longitudinal research with people experiencing MEH. Our experience is that the involvement of 'peer researchers' helps address some of these challenges. The gains of involving service users in this way include: research that is better informed by the concerns and needs of homeless people; methods which are more sensitive to and better enable effective engagement of research participants; and likely to elicit fuller and more open responses from them because people with similar experience have been involved in formulating them (see Beresford, 2007).

A team of six 'peer researchers' will be recruited and trained with a view to one 'peer researcher' being attached to each site (ideally they will be local to the research site in which they will work so as to have good knowledge of services and local community). Many of the SIHC projects already include 'peer mentors' (it maybe that the peer mentors would be willing to take on the additional role of peer researcher). A two day training course will be provided to allow the peer researchers to meet as a team and latterly as part of a specialist service user advisory group. In the fieldwork sites, the peer researchers and university researchers will work closely together (sharing expertise and offering mutual support and supervision).

The 'peer researchers' will be involved in all duties commensurate with being a full member of the research team (i.e. treated equally and inclusively). The 'peer researchers' main role will be to accompany and/or carry out joint interviews with the university researchers (where appropriate⁵) and to stay 'in touch' with key informants during the three month intervening period (mainly through telephone contact and informal contacts, for example, at day centres and other venues).

Two experienced 'peer researchers' are collaborating at the design stage and have made the following recommendations: *"I remember from my time in the Hostel the Buddy system we had in place, I became a buddy and the idea was that any other client who needed to talk to someone about any issues they had could talk to their designated buddy at any time of day. Buddies would have a smiley face on their room door with their name underneath. My idea is that peer researchers could be used to Buddy homeless people due for discharge at an early stage, as early intervention would be needed, and peer researchers could build a rapport with the service user and gain the trust needed to be able to follow them over a 3 month period as intended"*.

There is the potential here, that the peer research process could (while engendering the rapport and trust needed to prevent 'attrition') begin to cross over into the actual discharge process. This will need to be carefully managed and, if it occurs, made explicit in the analysis. Beresford (2007) would not however, see this as inherently problematic or risky but as part of the inherent nature of service user research with its commitment to change and challenging traditional research practices which have often been exclusionary. Perhaps the most important consideration is that if the research does build relationships and in interest in peer research among the homeless people taking part in the study, then there are opportunities in place for that to continue beyond the life of the programme. This is where our links with Homeless Link, the faculty of Inclusion Health and Revolving Doors are important because they already have these facilities in place.

⁵ If for example, the peer researcher is personally known to the interview participant then it may not be appropriate for them to be part of the interview as it may prevent full and open discussion.

Data Analysis

In relation to each individual site, data will be brought together in a matrix showing the different domains of interest. The domains will generate a series of codes which will be used to analyse documentary, interview and observational data. Key qualitative and quantitative (cost related) findings will then be summarised, ensuring the data from each system can be considered together (O’Cathain et al, 2010). Theory will then be developed for each site: initial KPPs (the provider view of the CMOs) will be further tested and challenged and added to within the qualitative interviews and initial analysis. This final stage of analysis for each case will provide the opportunity for further hypotheses to be developed as CMO configurations articulating how the research team believe each system, made up of its own particular combination of ‘resource components’ is generating outcomes.

Comparative (cross case) analysis will then be performed using further matrices in the predictor-outcome form (Miles & Huberman 1994) to examine patterns and predictors for a series of key outcomes. This combination will allow us through a process of modified analytic induction (Byng, Norman, & Redfern 2005) to develop hypotheses about the key influences on good quality discharge practices and a range of other outcomes. This process should also provide some initial findings about the likely cost differences between the different overarching models and specific ‘resource components’ for managing admissions, including insight into which hypothetical mechanisms of action or conducive contexts have the most significant resource implications (whether implying costs or savings).

These initial hypotheses will be tentative and will be tested by relooking across the cases for confirmation, rebuttal and ‘silence’. The results will not include definitive causal mechanisms and context dependencies, but will provide weightier evidence as to which mechanisms are likely to be effective in which contexts. This will allow us to develop guidance for commissioners, as well as develop provisional theories which can be tested further in more quantitative studies.

Preliminary findings will be fed back to sites for scrutiny and further analysis by means of a local feedback event (n=6). This is intended to guard against ascertainment bias where evaluators focus on outcomes which they have knowledge of and will investigate these outcomes more thoroughly.

Stage 3b: Delivery of Work Package Two – Data Linkage and Health Economic Analysis

Overview

The most comprehensive assessment to date of the health care usage of homeless populations in England was prepared and published by the Department of Health in March 2010. The report ‘Healthcare for Single Homeless People’ (Office of the Chief Analyst and Department of Health, 2010) analysed Hospital Episode Statistics data for people who gave their address as “No fixed abode” and also summarised data from six specialist GP practices who provided healthcare to homeless people. Extrapolating from these data they estimated that health care costs of homeless people are approximately eight times greater than those of similarly aged adults and this client group have almost three times the average length of stay when hospitalised with a cost of at least £85 million per year. However, there were several key problems with this analysis: the use of the “No fixed abode” category will miss many homeless people, for example those living in shelters, hostels or other temporary accommodation; the size of denominator populations to allow comparison of hospitalisation rates was highly uncertain and, those using specialist practices may be highly unrepresentative of the homeless population. To address these issues, WP2 will use data on homeless hospital admissions in order to determine the long term outcomes required to investigate the effectiveness and cost-effectiveness of different hospital discharge arrangements in England.

Data sets used for WP2

WP2 will extend the number of research sites to 20. This will include the 6 realist evaluation sites, plus a further 7 sites which have ‘clinically led’ SIHHC and 7 sites which have ‘housing link’ worker (non-clinically led) SIHHC. Appendix One provides a summary of the data which will be collected in each site and how for practical data collection purposes WP1 and WP2 will be integrated on the ground.

A total of five data sources will be used within this linkage study.

1. **Unconsented data:** SIHHC (hospital discharge projects) taking part in the study (Sites 1-2, 7-13 using clinically led services; Sites 3-4, 14-20 using non-clinically led services) will be asked to provide identifying and demographic data (forename, surname, aliases, date of birth, sex, current or previous postcodes, nationality, ethnicity, NHS number) for service users during the period November 2013 to a maximum of November 2016. However, note that for some services we will collect data that ends before 2016 as some will have been shut down before this time period and in some we may collect data at an earlier point, providing we receive the appropriate permissions and ethical and CAG approvals.
2. **Consented data from a subset of study fieldwork sites (n=6 sites):** We will collect data on 252 individuals across 4 sites with SIHHC (hospital discharge) schemes (two sites that are clinically led (Sites 1&2), two sites that are non-clinically led (Sites 3&4)) and on 126 individuals across two sites with standard /usual care (Sites 5&6), giving 378 individuals in total (63 per site x 6 sites). The data collection for this part of the work is contained in a baseline, exit and 3-month follow up data questionnaire and will collect information on demographic information, along with dates of contact with SIHHC services, past use of health and social care services, mental health services, smoking, alcohol, and drug taking history, prison and homeless history, employment, user satisfaction with service delivery, and data on quality of life (EQ5D). The voluntary sector hospital discharge projects will be asked to help collect this data on behalf of the research team as part of their routine outcomes monitoring and will be reimbursed for doing so at a rate of £20 per completion. The data flow chart shows how this work will be integrated with WP1 for practical purposes.
3. The data flow chart shows how this work will be integrated with WP1 for practical purposes.
4. **Find and Treat Service Unconsented Data:** In order to compare the outcomes of those individuals accessing SIHHC and standard usual care services to a wider homeless population, we will also use data from Find and Treat NHS clinical service that screens for tuberculosis in homeless hostels in London and people using day centres and the street population accessing soup kitchens and cold weather shelters. To identify hospital admissions during the introduction of SIHHC services (November 2013 to November 2016) we will collect all data from Find and Treat since 2008 as this will allow a period of follow up for this community group to have subsequent hospital admissions.
5. **Personal Demographics Service (PDS):** Where available, NHS number will be collected from the 20 sites. Where it is not available we will use the Personal Demographics Service (PDS) at Health & Social Care Information Centre (HSCIC) to undertake a 'list clean' in order to add an NHS number to as many records as possible.
6. **Linked Hospital Episode Statistics (HES) ONS mortality data:** Once these records have been 'list cleaned' they will then be linked by HSCIC to their linked Hospital Episode Statistics (HES) ONS mortality database to provide data on hospitalisation rates (emergency and planned hospital admissions). In addition to data linked from the 18 SIHHC services and Find and Treat, we will also request a geographically comparable and representative sample of lowest quintile of deprivation population in HES (based upon the index of multiple deprivation) equal in size to the Find and Treat dataset during the hospital admission study period of November 2008 to November 2016 and is therefore likely to be 60,000 records. Unlike the F&T group who are homeless, but not necessarily admitted to hospital, we will collect data on this comparison group to enable us to calculate rates for the primary and secondary outcomes in a group of individuals in the most deprived population that have been admitted to hospital and therefore comparable in terms of history of hospital admission to the homeless individuals at the 20 sites.

Comparative Effectiveness Analysis

Data linkage

We will provide data from the 20 sites and Find and Treat to the Health & Social Care Information Centre (HSCIC). These records will be list cleaned and then linked by HSCIC to the Hospital Episode Statistics (HES) database to provide data on hospitalisations and A&E attendances and deaths for individuals in the 20 sites and Find and Treat database. In order to be able to identify co-morbidities more accurately we will request data on all previous admissions for these groups for identification of

comorbidities but analyses of admission rates will be confined to the period November 2008 – November 2016.

Study design:

We will construct a virtual cohort that will take first date of admission as the entry point (earliest November 2008) or, for control groups an admission matched by month and year restricted to individuals over the age of 18. Exit from the virtual cohort will be at the end of November 2016 or, if earlier, date of death. We will conduct person-level analyses using Poisson regression to compare overall hospitalisation rates, emergency admission rates, all cause mortality and avoidable mortality rates in:

- 1) Homeless people using the different categories of SIHCC service (clinically led services - sites 1&2, 7-13; non-clinically led services - sites 3&4, 14-20)
- 2) Find and Treat Homeless population and sites 5&6 with standard usual care
- 3) Socially deprived non-homeless population.

We will also conduct analyses at the level of the hospital admission whereby through knowledge of the dates of SIHHC involvement and service user group we will be able to classify hospital admissions in the period November 2008- November 2016) into the following groups:

- 1) Admissions in a socially deprived non homeless population
- 2) Admissions in the general homeless population (Find and Treat data)
- 3) Admissions in homeless service users prior to their first engagement with SIHCC
- 4) Admissions in homeless service users where there is evidence of SIHCC support for that admission
- 5) Admissions in homeless service users that occur following SIHCC support but where there is no record of support for the subsequent admission.

We will use Cox regression survival models and Kaplan Meir plots (accounting for clustering at the level of the hospital and individual) to explore how admissions in these groups compare in terms of the following primary outcomes:

- 1) Time from discharge to next A&E attendance
- 2) 28 day emergency readmission rates (these are a commonly used indicator within the NHS)
- 3) Time from discharge to next hospital inpatient readmission

Secondary outcomes we will examine include:

- 1) Time from admission to death
- 2) Duration of inpatient admission
- 3) Time from admission to mortality from causes amenable to healthcare
- 4) Time from discharge to admission with ambulatory care sensitive (ACS) conditions
- 5) Time from discharge to next planned admission,
- 6) Overall readmission rates
- 7) Unscheduled readmission rates
- 8) All cause mortality
- 9) International Classification of Diseases 10th Revision chapter-specific mortality
- 10) In-patient costs using Health Resource Group

For those service users with quality of life data we will explore how this changes following involvement of services and how it compares to the measures of quality of life derived from studies within Find and Treat service users and within socially deprived non-homeless groups.

A variety of mortality groupings have been developed to identify causes of death that could be considered avoidable through better medical care depending on age at which they occur. The proposals for the Public Health Outcome Monitoring framework take a similar approach and focus on key areas of avoidable mortality namely: mortality from suicide, communicable diseases, excess

seasonal mortality and, in persons aged less than 75 years, cardiovascular disease, cancer, chronic liver disease and chronic respiratory diseases. All of these are useful indicators within homeless populations.

Sample size:

Our sample size is based upon our estimates of historical data from the two types of SIHHC services (non-clinically led SIHHC services vs. clinically led SIHHC). We estimate that each type will have data from approximately 2,208 individuals for the two-year duration of the study, based on the services seeing on average 92 people per month.

From previous health service evaluation data on homeless groups we expect hospitalisation rates to be 0.7 episodes per person-year for homeless individuals admitted to hospital. An important reduction in readmission rates would be 10%. To undertake a sample size calculation to determine the study size required to detect such a reduction in readmission rates, the following variables were defined:

μ_0 = mean readmission rate in the baseline group = 0.7 episodes per person-year

μ_1 = mean readmission rate in the SIHHC group = 0.6 episodes per person-year

$v=1.96$ =percentage point of the normal distribution corresponding to a 5% two-sided significance level

$u=0.84$ =one-sided percentage point of the normal distribution corresponding to (100%- type II error [FN / (TP + FN)]) at 80% power

Using the following equation we estimated minimum sample size (per group) required for comparison of two rates (readmissions per person-year):

$$\begin{aligned} & (u+v)^2 * (\mu_1 + \mu_0) / (\mu_1 - \mu_0)^2 \\ & (0.84+1.96)^2 * (0.7 + 0.6) / (0.7 - 0.6)^2 \\ & =1019 \text{ person-years per group} \end{aligned}$$

Adjustment for confounders as far as is possible within the analysis results in a doubling of sample size (additional 10% per confounder; likely 10 confounding variables). Therefore 2038 person-years are required per group. Assuming that the SIHHC clinically led and non-clinically led sites see on average 90 service users each month during the study period (which for the purposes of this calculation we have assumed this to be November 2013-November 2015 as many services may have been shut down before this time period and in some we may collect data at an earlier point) we estimate that this translates to a total 2,160 individuals at each of these two types of service.

Follow up: For the primary analysis, individuals will enter the cohort on date of first hospital admission (earliest November 2008) and will be censored at the end of the study period (November 2016) or death, whichever comes first.

Confounding variables: As this is a virtual cohort, we will not be able to collect a complete set of confounding variable data on individuals at entry to the study. Where data is provided by SIHHC services we will include age (year and month only), sex. As a result of the lack of baseline data will use historical HES data to determine past medical history from the age of 18 which will be used to create a variable to control for co-morbid conditions and historical hospital usage. Analyses will therefore control for age, gender, comorbidities (identified from HES data) and reason for admission, classified by Healthcare Resource Group (HRGs – groupings of treatment episodes which are similar in resource use and clinical response). Subgroup analyses will compare the different types of SIHCC services. Where provided by the SIHHC services, we will also control in subgroup analysis for medical symptoms, past use of health and social care services, mental health services, smoking, alcohol, and drug taking history, prison and homeless history, employment history.

Health Economics Analysis:

This analysis will cover three economic models. The primary analysis will be a cost effectiveness model (linked to effectiveness data described in the comparative effectiveness section). This model will take an NHS perspective, with costs based on the linked HES data analysis and outcomes based on morbidity and mortality data. Outcomes of homeless users of the two types of SIHHC service (clinically led and non-clinically led) will be compared to each other and homeless controls. Outcomes used for this analysis will include: 1. Overall hospital readmission rates; 2. Unscheduled readmission rates; 3. 28-day emergency readmission rates. Cost estimates will be calculated using national NHS

tariff data, and will include length of hospital stay, emergency admissions, and fixed and variable costs estimates for each of the SIHHC services.

A second cost utility analysis will also be performed. This model will also take an NHS perspective and use costs based on HES data as for the cost effectiveness model, but additionally will use outcomes based on QALY data (primary EQ5D data collected through the field work in a subgroup of SIHHC users at baseline and three month follow up). Data from the two types of service will also be compared to each other and homeless controls from a wider London based survey of 455 homeless individuals accessing Find and Treat screening service for tuberculosis. QALY loss due to morbidity and mortality averted will be examined. Cost estimates will be based on HES data and calculated at two time points: baseline, including retrospective data up to 3 month prior enrolment in the study, and follow up, including prospective data up to 3 month service follow up. Outcome data (QALY) will be calculated from EQ5D data collected at baseline and 3 month follow up.

Cost and effectiveness estimates to be calculated for one year follow up period after entering the study with simulations to 20 years and costs will be discounted at 3.5% per annum. A sensitivity analysis will be carried out in order to examine uncertainties within the cost effectiveness and utility model to examine the uncertainty within the parameters used.

A third cost utility analysis will also be performed using costs and health outcome (consented) data collected from the 6 case study sites. This model will take an NHS and also a wider public sector perspective (e.g. exploring costs for use of social care, prison, supported housing etc.).

The added value of this third economic analysis would be to estimate the cost utility of hospital discharge services from a larger public sector perspective whilst incorporating contextual information from the realist evaluation into the economic evaluation.

The objective of this economic evaluation will be to:

- Evaluate the effect of SIHHC on access to resources and their attached costs (health and mental health services, drug and alcohol services, social care services, housing situation) and on individual wellbeing, and how effect may vary across schemes in different sites
- Compare the effect of SIHHC on access to resources and individual wellbeing to sites relying on standard provision only (i.e. those that do not have access to SIHHC)
- Evaluate any variation in effect of SIHHC across time (dependent on opportunities for tracking participants over the longer term)

Where feasible, data on access to resources will be collected up to 12-month from discharge. This will allow capturing changes in yearly estimate before and after participating in SIHHC. To determine the effect of the SIHHC we will compare individual service use and health outcome after joining the study (based on 3-month follow-up) with our estimate of their service use had they not participated (one year prior to the enrolment). The specialist schemes will then be compared to the standard schemes which will be assessed in the same way. Cost of care will be estimated multiplying individual-specific health care resource utilization by unit cost using available tariff.

Sample size for economic evaluation:

Data from our 455 controls from Find & Treat provides a mean EQ5D visual analogue scale score of 65 and standard deviation of 25. An important improvement (or worsening) in the EQ5D visual analogue scale (0 to 100) as result of the provision of specialist services to this group would be 10. Using a power of 80% at the 0.05 level of significance (two sided) and a two sample *t* means test it is estimated that a sample size of 100 is required to detect such a change. Based on our previous work it is assumed that of all eligible service users identified, 80% of these will consent to take part in the study, and therefore 126 individuals will need to be approached to achieve this sample size. Therefore, 63 individuals will be approached at each of sites 1-6 (for a total of 378 individuals).

Sensitivity analysis and economic discounting: Cost and effectiveness estimates to be calculated for one year follow up period after entering the study with simulations to 20 years and costs will be discounted at 3.5% per annum. A sensitivity analysis will be carried out in order to examine uncertainties within the cost effectiveness and utility model to examine the uncertainty within the parameters used.

Additional ethical issues

The approach to this analysis takes into account the current political and public climate for approvals and linkage of personal identifiable data. We have therefore developed a two stage approach to our work: Firstly, to ensure a minimal dataset for this analysis we will seek individual consent from individuals using SIHHC services, to link their personal data to HES with appropriate NHS ethics approval; secondly, and to improve the power of the study by including a greater number of individuals, we will apply for approval under Section 251 of the NHS Act 2006 to obtain identifiable service user information from services to achieve linkage from an additional sixteen SIHHC services nationally. Our Section 251 request will comply with the Act by being "in the interests of improving service user care" and because there is no reasonable practicable alternative. The linked dataset will be stored in the Farr Safe Haven, a robust infrastructure certified according to international and national information security standards (ISO/IEC 270001 and NHS IG Toolkit). Data providers will securely upload encrypted identifiable data to a highly secure system. The UCL Identifiable Data Handling Solution (IDHS) has recent experience of gaining such approvals within 1 month of application.

The key principles are that:- Data security will be ensured through the a "Trusted Third Party"/"Safe Haven" model where UCL will provide a secure environment for processing identifiable personal data, where only approved researchers' can gain access to the data and where there are penalties for anyone who abuses personal data Research is needed to improve the health of vulnerable populations.

The main ethical issues are those that relate to the use of personal information without the explicit consent of those from whom the information is derived i.e. the second stage of data linkage proposed within this work. The recent Academy of Medical Sciences Report "A new pathway for the regulation and governance of health research" January 2011 highlights the importance of research use of health and social care data to improve understanding of disease and provision of services but state that "access to service user data for research is currently hampered by a fragmented legal framework, inconsistency in interpretation of the regulations, variable guidance and a lack of clarity among investigators, regulators, service users and the public". Despite these complexities we are confident of both the ethical and legal basis of our proposed research. The robust methodology described uses previously approved principles used by ourselves and other groups to link health and non-health data. We have allocated time within the grant to navigate these complexities and ensure relevant approvals. We also plan to host a number of Patient and Public Involvement (PPI) events to engage the full range of stakeholders in addressing these issues

Safeguarding

Researchers have a legal and moral duty to respect the rights of participants for their participation and data to remain confidential. However, there are some circumstances where a disclosure might be made by a participant, or evidence may come to light of a safeguarding issue. In such circumstances, the researcher may have a legal, professional or moral duty to breach the confidentiality of participants and to inform a third party outside of the research team. In such circumstances, the Chief Investigator will inform the Safeguarding Lead in the local authority and /or the NHS Trust, and the local authority Chief Executive and/or the clinical governance lead/Chief Executive in the NHS Trust or the police. The research team will receive training in safeguarding as part of their induction and will be made aware of these procedures. At the commencement of the fieldwork in each study site we will contact local safeguarding leads to alert them the research is taking place. Study participants will also be made aware on the Information Sheet that there may be limits to confidentiality, and that certain situations would mean that the researcher has a duty to inform a relevant authority outside of the research team.

Patient and Public Involvement (PPI)

In addition to the plans for PPI described in the main proposal, this work package will also include three one day engagement events in each of the Public Health Regions of England (North, Central and South). The events will be designed to look at all aspects of data linkage, the ethical issues that may arise, what would be required of services willing to take part in this study and the potential benefits, for example, improved outcomes data. The events will target PPI stakeholders involved in

all SIHHC hospital discharge schemes across England (including service providers and service users). On completion of the study, three further events will be hosted to feedback the findings of study. These events will be in addition to the initial expert seminar and the local feedback events planned for each fieldwork site in order to give sufficient time to look in depth at 'data linkage' (rather than findings around 'hospital discharge per se').

Stage 4: Dissemination and Outputs

In addition to a full project report for the NIHR's journal series and academic papers in learned journals, we will target key professional journals (e.g. Connect, Nursing Times, Community Care, etc.) and organisational/network newsletters and social media with articles and briefings to promote the project throughout and once findings are completed. The combination of these will bring a comprehensive description of the research and academic credibility to the evidence (i.e. the project report and academic articles) with accessible insights and recommendations targeted at specific audiences. Finally, if the findings support this, we will also produce a commissioner's guide (checklist for practice) in the form of toolkit.

In designing for impact we are aware that passive forms of dissemination do not always promote evidence informed practice. We have secured the expert support of Homeless Link who will provide consultancy and support throughout the life of the project to help us develop work in this area.

Gantt Chart

(24 months)	1-3 Q1	4-6 Q2	7-9 Q3	10-12 Q4	13-15 Q5	16-18 Q6	19-21 Q7	22-24 Q8
Preparation & Ethics								
Literature Review								
WP1 Fieldwork Site 1/2								
WP1 Fieldwork Site 3/4								
WP1 Fieldwork Site 5/6								
WP2 PPI Regional Engagement Events (n=3)								
WP2 Data linkage permissions and collection								
WP2 Set up, cleaning of data, and performing linkage								
WP2 Preparing cost data, EQ5D data entering and cleaning								
WP2 Statistical analysis for effectiveness study								
WP2 Cost effectiveness analysis								
Overarching Analysis & Report Writing								
Local Feedback (n=6), Regional PPI (n=3) and National Dissemination.								
Advisory Group Meetings	*	*	*	*	*	*	*	*

WP =Work Package

PROJECT TEAM

Dr Michelle Cornes (King's College London): CI overall management responsibility for delivery of contract and lead for Work Package 1. [50% FTE]

(Working Mainly Work Package 1)

Dr Martin Whiteford (Liverpool University: Main field researcher for Work Package 1 [100% FTE]

(Working Mainly on Work Package 2)

Professor Andrew Hayward, (UCL) will be the overall lead for Work Package 2 [5% FTE]

Dr Robert Aldridge (UCL) will oversee the data linkage, statistical analysis and health economics aspects of this project. [5% FTE]

Dr Spiros Denaxas (UCL) Research Fellow in Health Informatics based in the Farr Institute for Health Informatics Research at UCL Partners: Will oversee and advise on the storage and processing of data in the Farr Safe Haven using national information security standards (ISO/IEC 270001 and NHS IG Toolkit) based on his extensive experience in this area. [5% FTE]

Dr Michela Tinelli (LSE) is a Fellow at LSE Health and Social Care: Lead for economic evaluation [20% FTE] Her role will be to prepare cost data and EQ5D datasets (months 1-12) and undertake economic analysis and reporting (months 12-24).

Wurie Fatima (UCL) responsible for liaising with the services, training staff on how to consent service users and collect EQ5D questionnaires, and assist the preparation and collection of data required for the linkage. Fatima will also administer the necessary ethical section 251 applications relevant to the project. At the end of the project the researcher will develop individual reports for the services to feedback to them about the outcomes of their clients, and the results of the wider project. [50% FTE]

Bio-informatician (UCL): database administrator responsible for setting up the probabilistic linkage process, cleaning the data from the 20 SIHHC schemes, and administer and clean the national HES data that will be used in the linkage and finally to perform the probabilistic linkage. [50% FTE]

Statistics post-doctoral researcher (UCL) After the data has been cleaned and linked, statistical analysis for effectiveness study will be performed by a statistics post-doctoral role. This individual will perform the statistical modelling for the comparative effectiveness analysis, conduct the sensitivity analyses, and produce data for the study and individual feedback reports for services [100% FTE in Y2]

(Role of other co-applicants)

Professor Richard Byng (Plymouth University); Will add SDR credibility and role in synthesis theory building [5% FTE]

Dr Mike Clark (LSE): To work with commissioners to develop the findings for practice [3% FTE]

Professor Graham Foster (QMUL): To advise on clinical issues, methodological and quantitative aspects of the study [5% FTE]

Dr Nigel Hewett (Pathway) is medical director of Pathway and will advise on clinical issues/site access. [5% FTE]

Professor Jill Manthorpe (King's College London): Responsible for literature review, support for the PI, and overall financial management [5% FTE]

Dr Jo Neale (Institute of Psychiatry): To offer methodological support, particularly in relation to the qualitative components of the case studies and to advise on analyses and dissemination [5% FTE]

Collaborators: Raphael Wittenberg (LSE and Oxford University) Natalie Atkinson (Expert by Experience) and Alan Kilminster (Expert by Experience), Helen Mathie (Homeless Link), Sarah Anderson (Revolving Doors), Mike Seale (Newman University Birmingham), William Lee and Lynne Callaghan (Plymouth University) Professor Mark Gabbay, Dr Stuart Oultram, Dr. Paula Byrne (Liverpool University), Dr Diane Exley, Brownlow Health (Liverpool) Andrew Hayward and Rachel Hunter (UCL).

REFERENCES

- Backer, T., Howard, E., & Moran, G. (2007) The Role of Effective Discharge Planning in Preventing Homelessness, *Journal of Primary Prevention*, Vol. 28 No. (3/4) pp. 229-243.
- Barton, P., Bryan, S., Glasby, J., Hewitt, G., Jagger, C., Kaambwa, B., Martin, G., Nancarrow, S., Parker, H., Parker, S., Regan, E., Wilson, A. (2006) *A National Evaluation of the Costs and Outcome of Intermediate Care for Older People*, Birmingham: University of Birmingham and University of Leicester.
- Beresford, P. (2007) User involvement, research and health inequalities: developing new directions. *Health and Social Care in the Community*, 15(4), 306-312 doi: 10.1111/j.1365-2524.2007.00688.x
- Brandoli, M. & Cocchi, M. (2013) Protected Hospital Discharges for Anyone Leaving Hospital without a Home to go to, *Homelessness in Europe*. Winter 2013 pp. 20-22.
- Byng R, Norman I, & Redfern S, (2005), Using realistic evaluation to evaluate a practice level intervention to improve primary health care for service users with long-term mental illness. *Evaluation* 11[1], 69-93.
- Cameron, A. Lart, R. Bostock, L. and Coomber, C. (2013) Factors that promote and hinder joint and integrated working between health and social care services: a review of research literature. *Health and Social Care in the Community*, doi: 1111/hsc.12057
- Cockersell, P. (2011) Homelessness and Mental Health: Adding Clinical Mental Health Interventions to Existing Social Ones can Greatly Enhance Positive Outcomes, *Journal of Public Mental Health*, Vol. 10 No. 2 pp. 88-98.
- Cornes, M., Joly, L., Manthorpe, J., O'Halloran, S., and Smythe, R. (2011) Working Together to Address Multiple Exclusion Homelessness, *Social Policy and Society*, 10(4), 513-522
- Cornes, M. and Clough, R. (2001) The Continuum of Care: older people's experiences of intermediate care. *Education and Ageing*, 16(2)
- Crane, M. & Warnes, T. (2011) Single Homeless People's Access to Health-care Services in South Yorkshire (CLAHRC SY, Sheffield).
- Crisis (2011) *Homelessness: A Silent Killer - A Research Briefing on Mortality amongst Homeless People* (London, Crisis).
- Daly, B. (2007) *The Oral Health Needs of Homeless People Living in an Urban Setting in the UK*, unpublished PhD Thesis, London, King's College London.
- Department for Communities & Local Government (2011) *Vision to End Rough Sleeping: No Second Night Out*, DCLG, London.
- Department for Communities and Local Government (2012) Evidence Review of the Costs of Homelessness. London: DCLG.
- Department of Health (2003) *Discharge from Hospital: pathway, process and practice*. London: DH.

- Department of Health (2010) Healthy lives, healthy people: our strategy for public health in England - White Paper. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_121941 (accessed 1 Feb2011).
- Department of Health (2013) *Ten Million Cash Boost to Improve the Health of Homeless People*. Press Release.
- Dixon-Woods, M, Baker, R., Charles, K., Dawson, J., Jerzembek, G, Martin, G., McCarthy, I., McKee, L., Minion, J., Ozieranski, P., Willars, J., Wilkie, P., and West, M. (2013) Culture and behaviour in the English NHS: Overview of lessons from a large multi-method study, *BMJ Quality and Safety*. 0(1) doi: 10.1136/bmjqs-2013-001947
- European Observatory on Homelessness (2013) The Costs of Homelessness in Europe: An Assessment of the Current Evidence Base. Brussels: FEANTSA. Available at http://www.feantsaresearch.org/IMG/pdf/feantsa-studies_03_web.pdf
- Evans, J., Stone, K., Manthorpe, J. and Higginson, J. (2013) *MRC Guidance on Developing and Evaluating Complex Interventions: Application to Research on Palliative and End of Life Care*. Method Review 15, London NIHR/School for Social Care Research. Francis, R. (2013) *Report of the Mid Staffordshire NHS Foundation Trust Public Inquiry*. London: Stationary Office.
- Godfey, M., Townend, J., Cornes, M., et al. (2008) *Reimbursement in Practice*, Scotland: University of Stirling.
- Gould, M. (2009) No Fixed Address. *Health Service Journal*, 12th February p20-22
- Hewett, N., Buchman, P., Musariri, J., Sergeant, C., Johnson, P. Abeysekera, K. Eleftheriades, C., McCormick, B., Halligan, A., Marlin, N., Kerry, S., and Foster, G. (under review) *Randomised control trial of General Practitioner led in-hospital management of homeless people ('Pathway')*
- Hewett N, Halligan A, Boyce T. (2012) A general practitioner and nurse led approach to improving hospital care for homeless people. *BMJ* 2012; 345:e5999
- Hewett, N. & Halligan, A. (2010) Editorial: Homelessness is a Healthcare Issue, *Journal of the Royal Society of Medicine*, 103, pp. 306-307.
- Hewett, N., (2010) *Evaluation of the London Pathway for Homeless Service users*. London: UCL Hospitals.
- Hewett, N. and Dorney-Smith, S. (2013) *King's Health Partners and the Impact of Homelessness*. London: King's Health Partners.
- Homeless Link (2010) *The Health and Wellbeing of People who are Homeless: Evidence from a National Audit* (London, Homeless Link).
- Homeless Link & St Mungo's (2012) *Improving Hospital Admission and Discharge for People who are Homeless* (London, Homeless Link & St Mungo's).
- Homeless Link (2015) *Evaluation of the Homeless Hospital Discharge Fund*. London: Homeless Link
- Huxley, P., Maegusuku-Hewett, T., Evans, S., Cornes, M., Manthorpe, J., and Stevens, M., (2010) Better evidence for better commissioning: a study of the evidence base of generic social care commissioning guides in the UK. *Evidence and Policy*. 6(3), 291-308
- Manzano-Santaella, A., (2011) A realistic evaluation of fines for hospital discharges: incorporating the history of programme evaluations in the analysis. *Evaluation* 17(1) 21-36
- McDonagh, T. (2011) *Tackling homelessness and exclusion: Understanding complex lives*, Joseph Rowntree Foundation, York.
- Fitzpatrick, S., Johnsen, S., & White, M. (2011) Multiple exclusion homelessness in the UK: Key patterns and intersections. *Social Policy and Society* 10 (4), 501-502.
- Miles MB & Huberman AM, (1994), *Qualitative data analysis: an expanded sourcebook*. Second Edition. Sage, London.
- Newcombe HB. *Handbook of Record Linkage: Methods for Health and Statistical Studies, Administration and Business*. Oxford University Press 1988.
- O'Cathain A, Murphy E and Nicholl J, (2010), Three techniques for integrating data in mixed methods studies. *BMJ* 341:c4587.
- Office for National Statistics NSPD. Measuring premature and avoidable mortality. Off. Natl. Stat. 2008. <http://www.ons.gov.uk/ons/about-ons/consultations/closed-consultations/measuring-premature-and-avoidable-mortality/index.html> (accessed 16 Oct2014).
- Office of the Chief Analyst - Department of Health. *Healthcare for Single Homeless People*. London: : Department of Health 2010. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_114250 (accessed 14 Nov2012).
- Pathway (2013) *Pathway Annual Review*, London: Pathway.
- Pawson R and Tilley N, (1997), *Realistic Evaluation*. London: Sage.

- Pearson, M., Hunt, H., Cooper, C., Shepperd, S., Pawson, R., and Anderson, R. (2015) Providing effective and preferred care closer to home: a realist review of intermediate care. Health and Social Care in the Community. Doi: 10.1111/hsc.12183
- Pearson, M., Hunt, H., Cooper, C., Shepperd, S., Pawson, R and Anderson, R. (2013) *Intermediate Care: a realist review and conceptual framework*, London: NIHR
- Petch, A. (2012) Tectonic plates: aligning evidence, policy and practice in health and social care integration, *Journal of Integrated Care*, 20(2) 77-88.
- Potz N, Powell D. Manual for the probabilistic linkage of infection and death records. 2007.
- Story A, Aldridge RW, Gray T, *et al.* Influenza vaccination, inverse care and homelessness: cross-sectional survey of eligibility and uptake during the 2011/12 season in London. *BMC Public Health* 2014;**14**:44. doi:10.1186/1471-2458-14-44
- Tansley, K., and Gray, J., (2009) Ensuring safe and appropriate discharge for people who are homeless or in housing need. *Nursing Times*. 105 (40) 14-17
- Teddlie C& Tashakkori A (2003) 'Major Issues and Controversies in the Use of Mixed Methods in the Social and Behavioural Sciences' Chapter 1. In Tashakkori A & Teddlie C eds. (2003) *Handbook of Mixed Methods in Social and Behavioural Research*. Sage. Thousand Oaks. London.
- White, J. (2011) Homelessness and Hospital Discharge in Wirral: An investigation into the Hospital Discharge Project: NHS Brief No.2 (Centre for Health Service Economics and Organisation).
- Wilson, V. and McCormack, B. (2006) Critical realism as emancipatory action: the case for realistic evaluation in practice development. *Nursing Philosophy*, 7 45-57

APPENDIX ONE: FLOW CHARTS

Box 1: Summary Table of Data to be Collected in Each Study Site

	WP1 In-depth fieldwork Topic Guides 1-6	WP2 Consented EQ5D Baseline +Follow-Up Questionnaires	WP2 Unconsented
Realist Evaluation Case Study Sites			
Site 1 – SIHHC Clinically led	✓	✓	✓
Site 2- SIHHC Clinically led	✓	✓	✓
Site 3 – SIHHC Non-clinical (e.g housing link worker)	✓	✓	✓
Site 4 – SIHHC Non-clinical (e.g. housing link worker)	✓	✓	✓
Site 5 – Usual/Standard Care	✓	✓	✓
Site 6 – Usual Standard Care	✓	✓	✓
Other Research Sites			
Site 7 – SIHHC Clinically led			✓
Site 8 – SIHHC Clinically led			✓
Site 9 – SIHHC Clinically led			✓
Site 10 – SIHHC Clinically led			✓
Site 11 – SIHHC Clinically led			✓
Site 12 – SIHHC Clinically led			✓
Site 13 – SIHHC Clinically led			✓
Site 14 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 15 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 16 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 17 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 18 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 19 – SIHHC Non-clinical (e.g housing link worker)			✓
Site 20 – SIHHC Non-clinical (e.g housing link worker)			✓

Box 2: Arrangements for Data Collection in Fieldwork Sites 1-6*

Scheme Typologies Across Fieldwork Sites 1-6

Sites 1&2: In these sites the hospital has access to a specialist integrated hospital discharge scheme (SIHHC) for homeless people which is ‘clinically-led’

Sites 3&4: In these sites the hospital has access to a specialist integrated discharge scheme (SIHHC) for homeless people which is non-clinical (e.g. provides access to housing advice only)

Sites 5&6: In these sites the hospital does not have access to any specialist schemes for discharging homeless service users (i.e. ‘standard or usual care’)

Stakeholder Recruitment (n=60)

In each of these 6 sites, a minimum of 10 stakeholders (practitioners, managers and clinicians) will be recruited for semi-structured interview as part of Work Package 1 (Realist Evaluation). Prospective participants will be given an invitation letter and project information sheet. Informed consent will be ascertained prior to commencement of the interview.

Service user Recruitment (n=378)

In each of these 6 sites a total of 63 service users will be recruited:

In each site:

10 service users will be allocated to Group A

53 service users will be allocated to Group B

Group A (n=10 service users)

Inclusion Criteria: Service users will be recruited on a first come first served basis until the following demographic categories are filled:

Young Homeless People (1 male/ 1 female) (Age 18-24)

Men in the middle age range (4 male) (Age 25-49) [Known to have the most complex needs (McDonagh, 2011)]

Women in the middle age range (25-49)(2 Female)

Older Homelessness (50+) (1 male/1 female)

NB: Once these categories are filled, homeless service users will then be allocated to **Group B (n=53 service users)**

Group A

Data is to be collected for both Work Package 1 (Realist Evaluation) and Work Package 2 (Comparative Effectiveness and Economic Evaluation)

Homeless service user is on the hospital ward.

Step 1

Seek the support of hospital/scheme discharge coordinators to recruit 10 service users for Group A (n=60)

Step 2

The hospital/scheme discharge coordinator informs the service user about the study, provides an invite letter and a 'Service user Information Leaflet' (Form A). These will be talked through and permission sought to pass the service user's contact details to the research team.

Step 3

A few days after discharge from hospital, the researcher makes contact with the service user (where permission to contact has been given) to arrange a community visit. This will be made by the University Researcher and Peer Researcher working in pairs.

The researcher provides another copy of the 'Service user information Leaflet' (Form A), answers any questions and if the service user is willing to participate

takes informed consent (either written or verbal)(Form A).

Step 4

A semi-structured interview is carried out (Topic Guide 1 – Time 1)

A baseline questionnaire is completed (WP2/BL)

Step 5

Qualitative interview data is transcribed and stored in pass word protected computer at King's College London

Secure upload of consented (WP2BL) individual data to Farr data safe haven

Step 6

The Peer Researcher maintains telephone contact until Time 2 interview takes place at 3 month follow-up (Topic Guide 1 –Interim Telephone)

Notes are taken and passed to King's College London to be stored on pass word protected computer

Step 7

At 3 month follow-up, a semi-structured interview is carried out (Topic Guide 1 – Time 2)

A follow-up questionnaire is completed (WP2/FU)
(University and Peer researcher working in pairs)

Step 8

Qualitative interview data is transcribed and stored in pass word protected computer at King's College London

Secure upload of consented (WP2FU) individual data to Farr data safe haven (WP2/FU)

Group B

Data is to be collected for Work Package 2 (Comparative Effectiveness and Economic Evaluation) Only

Homeless service user is on the hospital ward.

Step 1

Seek the support of hospital/scheme discharge coordinators to recruit 53 service users for Group B at each site (n=318)

Step 2

The hospital/scheme discharge coordinator informs the service user about the study, provides an invite letter and a 'Service user Information Leaflet' (Form B). These will be talked through and permission sought to pass the service user's contact details to the research team.

Step 3

A few days after discharge from hospital, the researcher makes contact with the service user to arrange a community visit.

The researcher provides another copy of the 'Service user Information Leaflet' (Form B) answers any questions and if the service user is willing to participate takes informed consent (either written or verbal) (Form B)

Step 4

A baseline questionnaire is completed (WP2/BL)

Step 5

Secure upload of consented (WP2/BL) individual data to Farr data safe haven

Step 6

A follow-up questionnaire is completed on exist from the scheme and at three months follow up (WP2/FU)

Step 7

Secure upload of consented (WP2/FU) individual data to Farr data safe haven (WP2)

Box 3: Arrangements for Data Collection in Fieldwork Sites 7-20*

Scheme Typologies Across Sites 7-20

Sites 7-13: In these sites the hospital has access to a specialist integrated hospital discharge scheme for homeless people which is 'clinically-led'

Sites:14-20 In these sites the hospital has access to a specialist integrated discharge scheme for homeless people which is non-clinical (e.g. provides access to housing advice only)

Comparator Databases:

Find and Treat [comparator database]

Hospital Episode Statistics [comparator database representative sample of lowest quintile of deprivation equal in size to the Find and Treat dataset]

Step 1

CAG Approvals in Place

Step 2

Unique study ID created for each individual record and left on site

Step 3

Secure upload of operational service data (non-consented) to Farr data safe haven


Step 4

Transfer of sensitive data (first name, surname, dob, sex, current or previous postcodes, nationality, ethnicity, NHS Number, Hospital of admission, Study ID

Step 5:

Transfer of HES ID and HES data back to Farr data safe haven

Box 4: Detailed Data Flow for WP2:



Box 5: Data variables collected for WP2:**Linked HES / ONS data variables requested**

Field name (where known)	Description of contents
admidate	Date of admission
elecdate	Date of decision to admit
admimeth	Method of admission
admisorc	Source of admission
firstreg	First regular day or night admission
elecdu	Waiting time
disdate	Date of discharge
disdest	Destination on discharge
dismeth	Method of discharge
DISREADYDATE	Discharge ready date (delay in discharge)
ARRIVALDATE	Arrival date
AEATTENDCAT	Attendance category
AEATTENDDISP	Attendance disposal
TREAT2_N	treatment
APPTDATE	Appointment date
ATTENDED	Attended or did not attend
TRETSPEF	Treatment specialty
cause_of_death	Original underlying cause of death
cause_of_death_non-neonatal	Non-neonatal cause of death mentions
	Match rank (ranking of quality of match)
dod	Date of death

SIHHC data variables

Other than the demographic information, we expect few sites to contain all the data variables listed below, but where they are (and for example we know of several sites that collect EQ5D data on all service users) we will collect these unconsented data. The list below therefore provides a potential list of variables we would attempt to collect at each site.

Example field name (where known)	Description of contents
date_interviewed	Date of interview
first_name	First name
surname	Surname
middle_names	Middle names (s)
Alias	Any alias names
sex	Sex of individual
address1	Address line 1
address2	Address line 2
address 3	Address line 3
postcode	postcode

telephone	Home telephone number
mobile	Mobile telephone number
plbirth	Place of birth
ethnicity	Ethnicity
Date_admission	Date of hospital admission
Ad_diagnosis	Admitting diagnosis
Date_discharge	Date of hospital discharge
sympwloss	Currently has symptoms of weight loss for no reason
sympcough3w	Currently has been coughing for more than 3 weeks
sympfevns	Currently has fever or night sweats
symphaemopt	Currently is coughing up blood
sympsob	Currently has shortness of breath
symprone	Currently has none of these symptoms
gpyeslocal	Registered with Local GP
gpnotlocal	Registered with GP that is not local
noreggp	Not Registered to GP
ae_visits	Number of A&E visits in the last year
op_visits	Number of hospital outpatient user visits in the last year
hospitalisations	Number of hospitalizations in the last year
vaxhepb	Individual been vaccinated against Hepatitis B
vaxhepbtimes	Individual been vaccinated against Hepatitis B – how many times
vaxhepbnever	Individual has never vaccinated against Hepatitis B
vaxhepbdk	Does not know if individual has been vaccinated against hepatitis B
HIVtestyes~s	Individual has been tested positive for HIV
HIVtestyes~g	Individual has been tested negative for HIV
HIVtestno	Individual has never been tested for HIV
HIVtestdk	Individual does not know if he/she has been teste...
HIVrx	Individual has been tested negative for HIV and treated
HCVtestyes~s	Individual has been tested positive for HCV
HCVtestyes~g	Individual has been tested negative for HCV
HCVtestno	Individual has never been tested for HCV
HCVtestdk	Individual does not know if he/she has been teste...
HCVrx	Individual has been tested positive for HCV and treated
HBVtestyes~s	Individual has been tested positive for HBV
HBVtestyes~g	Individual has been tested negative for HBV
HBVtestno	Individual has never been tested for HBV
HBVtestdk	Individual does not know if he/she has been tested for HBV
HBVrx	Individual has been tested positive for HBV and treated
TBtestyes~s	Individual has been tested positive for TB

TBtestyes~g	Individual has been tested negative for TB
TBtestno	Individual has never been tested for TB
TBtestdk	Individual does not know if he/she has been tested for TB
TBrx	Individual has been tested positive for TB
benefits	Does individual get any state benefits
smokecigs	Does individual smoke cigarettes
smokecigsn	How many cigarettes per day
smokecigsyr	How many years smoking cigarettes
drink	Has individual or a health worker ever been concerned about your drinking or suggested you cut down?
prison	Individual has ever been in prison before
arrests	How many times has the person been arrested by police
nights_prison	How many nights has the person spent time in prison
magistrate	Magistrate court attendance
crown	Crown court attendance
nights_cust	Number of nights spent in police custody
mh_admisssions	Mental health hospital admissions
mh_community	Mental health community provision
la_care_home	Local authority funded care home for people with mental health problems
a_care_mh	Local authority funded day care for people with mental health problems
herinj	Individual has ever injected heroin and length in years
ccinj	Individual has ever injected crack/cocaine and length in years
hersmk	Individual has ever smoked heroin and length in years
ccsmk	Individual has ever smoked crack/cocaine and length in years
needles	Individual has ever shared needles
social_ass	A social care assessment
social_con	A consultation with a social worker
care_home	Care Home
sleptrough	Individual has slept rough in the last year and if so number of nights
hostel	Individual has ever slept in a hostel in last year and if so number of nights
Squat	Individual has ever slept in a squat in last year and if so number of nights
Social_ten	Individual has had on social tenancy in last year and if so number of nights
Priv_rent	Individual has had private tenancy in last year and if so number of nights
Share_rent	Individual has ever private sector rented property in last year and if so number of nights
Income_loss	Did individual lose any work income as a result of your admission to hospital
Income_lossa	How much income was lost as a result of admission

Time_missed	Individual missed time from paid or unpaid work as a result of your admission to hospital
Time_missed	How much time was missed from paid or unpaid work as a result of your admission to hospital
Satistiifed	Was individual satisfied with the way their discharge from hospital has been managed
EQ5_mobili~b	EQ-5D: I have no problems in walking about
EQ5D_mobil~a	EQ-5D: I have some problems in walking about
EQ5D_mobil~d	EQ-5D: I am confined to bed
EQ5D_selfc~c	EQ-5D: I have no problems with self-care
E~rwashdress	EQ-5D: I have some problems washing or dressing myself
E~twashdress	EQ-5D: I am unable to wash or dress myself
EQ5D_u~puact	EQ-5D: I have no problems with performing my usual activities
EQ5D_u~ruact	EQ-5D: I have some problems with performing my usual activities
EQ5D_u~tuact	EQ-5D: I am unable to perform my usual activities
EQ5D_p~opain	EQ-5D: I have no pain or discomfort
EQ5D_p~dpain	EQ-5D: I have moderate pain or discomfort
EQ5D_p~rpain	EQ-5D: I have extreme pain or discomfort
EQ5~otanxdep	EQ-5D: I am not anxious or depressed
EQ5D~danxdep	EQ-5D: I am moderately anxious or depressed
EQ5~xtanxdep	EQ-5D: I am extremely anxious or depressed
EQ5D_scale	EQ-5D: Health on a scale of-100

Find and treat variables

Example field name (where known)	Description of contents
date_screened	Date of screening
first_name	First name
surname	Surname
middle_names	Middle names (s)
Alias	Any alias names
sex	Sex of individual
address1	Address line 1
address2	Address line 2
address 3	Address line 3
postcode	postcode
telephone	Home telephone number
mobile	Mobile telephone number
Venue	Venue in which screening took place

Outcome	Outcome of screening – including referral to hospital
EQ5_mobili~b	EQ-5D: I have no problems in walking about
EQ5D_mobil~a	EQ-5D: I have some problems in walking about
EQ5D_mobil~d	EQ-5D: I am confined to bed
EQ5D_selfc~c	EQ-5D: I have no problems with self-care
E~rwashdress	EQ-5D: I have some problems washing or dressing myself
E~twashdress	EQ-5D: I am unable to wash or dress myself
EQ5D_u~puact	EQ-5D: I have no problems with performing my usual activities
EQ5D_u~ruact	EQ-5D: I have some problems with performing my usual activities
EQ5D_u~tuact	EQ-5D: I am unable to perform my usual activities
EQ5D_p~opain	EQ-5D: I have no pain or discomfort
EQ5D_p~dpain	EQ-5D: I have moderate pain or discomfort
EQ5D_p~rpain	EQ-5D: I have extreme pain or discomfort
EQ5~otanxdep	EQ-5D: I am not anxious or depressed
EQ5D~danxdep	EQ-5D: I am moderately anxious or depressed
EQ5~xtanxdep	EQ-5D: I am extremely anxious or depressed
EQ5D_scale	EQ-5D: Health on a scale of-100