

FFIT FOLLOW UP

Data Linkage Report

Version 3.1

Author: Ruiqi Zhang

Report Date: 12 April 2016

SAP Version 1.0 Date: 30 October 2015

Protocol Version 2.0 Date: 29 July 2013

Table of Contents

Table 1	Summary of subject disposition	4
Table 2.1	Overall Count of Hospitalisations	5
Table 2.2	Hospitalisations Per Participant	12
Table 3	Psychiatric Hospital Admissions	16
Table 4	Cancer Incidences	17
Table 5	Deaths	19
Table 6	Hospitalisations due to Stroke or Myocardial Infarction	20
Table 7	Prescriptions	21
Table 8	Analysis of Hospitalisations	149

Executive Summary

Title

FFIT FU: Football Fans in Training Follow Up - Data linkage

Description

FFIT FU is a longitudinal, follow-up study of men who were aged 35-65 with BMI at least 28kg/m² at the start of the FFIT randomised controlled trial. Follow up visit is approximately 3.5 years post randomisation in the RCT

Primary objective

To explore the utility and feasibility of using linkage to routinely collected NHS datasets.

Note

* beside p-values indicate Wilcoxon-Mann-Whitney tests for continuous variables and Fishers exact test for categorical variables, ** indicate Wilcoxon signed rank tests for non-parametric and McNemar for binary variables.

Table 1
Summary of Subject Disposition, Overall and by Group

	All	Intervention	Control
RCT Population			
Numbers randomised	747	374	373
In the data linkage consented population	648 (86.7%)	312 (83.4%)	336 (90.1%)
In the data linkage analysis population	645 (86.3%)	310 (82.9%)	335 (89.8%)
Linkage not possible	102 (13.7%)	64 (17.1%)	38 (10.2%)
Consented Population			
Numbers consented to linkage	648	312	336
In the data linkage analysis population	645 (99.5%)	310 (99.4%)	335 (99.7%)
Linkage not possible	3 (0.5%)	2 (0.6%)	1 (0.3%)

Table 2.1.1
Overall hospitalisations prior to FFIT Intervention, Overall and by Group

		All	Intervention	Control
Total Hospitalisations				
Prior to intervention comparison (IA vs CB)		92	49	43
Primary Diagnosis	n (missing)	92 (0)	49 (0)	43 (0)
	Endocrine, nutritional and metabolic diseases	1 (1.1%)	0 (0.0%)	1 (2.3%)
	Diseases of the nervous system	8 (8.7%)	7 (14.3%)	1 (2.3%)
	Diseases of the eye and adnexa	2 (2.2%)	1 (2.0%)	1 (2.3%)
	Diseases of the circulatory system	17 (18.5%)	10 (20.4%)	7 (16.3%)
	Diseases of the respiratory system	4 (4.3%)	3 (6.1%)	1 (2.3%)
	Diseases of the digestive system	12 (13.0%)	6 (12.2%)	6 (14.0%)
	Diseases of the skin and subcutaneous tissue	3 (3.3%)	1 (2.0%)	2 (4.7%)
	Diseases of the musculoskeletal system and connective tissue	16 (17.4%)	9 (18.4%)	7 (16.3%)
	Diseases of the genitourinary system	6 (6.5%)	4 (8.2%)	2 (4.7%)
	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	19 (20.7%)	6 (12.2%)	13 (30.2%)
	Injury, poisoning and certain other consequences of external causes	3 (3.3%)	2 (4.1%)	1 (2.3%)
	Factors influencing health status and contact with health services	1 (1.1%)	0 (0.0%)	1 (2.3%)

Table 2.1.2
Overall hospitalisations during intervention, Overall and by Group

		All	Intervention	Control
Within intervention comparison (IB vs CD)		92	45	47
Primary Diagnosis	n (missing)	92 (0)	45 (0)	47 (0)
	Certain infectious and parasitic diseases	1 (1.1%)	1 (2.2%)	0 (0.0%)
	Neoplasms	2 (2.2%)	1 (2.2%)	1 (2.1%)
	Diseases of the blood/blood-forming organs, certain disorders involving the immune mechanism	1 (1.1%)	0 (0.0%)	1 (2.1%)
	Endocrine, nutritional and metabolic diseases	2 (2.2%)	1 (2.2%)	1 (2.1%)
	Diseases of the nervous system	8 (8.7%)	5 (11.1%)	3 (6.4%)
	Diseases of the eye and adnexa	5 (5.4%)	1 (2.2%)	4 (8.5%)
	Diseases of the circulatory system	6 (6.5%)	2 (4.4%)	4 (8.5%)
	Diseases of the respiratory system	3 (3.3%)	2 (4.4%)	1 (2.1%)
	Diseases of the digestive system	17 (18.5%)	8 (17.8%)	9 (19.1%)
	Diseases of the skin and subcutaneous tissue	2 (2.2%)	1 (2.2%)	1 (2.1%)
	Diseases of the musculoskeletal system and connective tissue	11 (12.0%)	3 (6.7%)	8 (17.0%)
	Diseases of the genitourinary system	5 (5.4%)	3 (6.7%)	2 (4.3%)
	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	20 (21.7%)	12 (26.7%)	8 (17.0%)
	Injury, poisoning and certain other consequences of external causes	4 (4.3%)	3 (6.7%)	1 (2.1%)
	Factors influencing health status and contact with health services	5 (5.4%)	2 (4.4%)	3 (6.4%)

Table 2.1.3
Overall hospitalisations during long term follow-up, Overall and by Group

		All	Intervention	Control
Long term follow-up comparison (IC vs CC)		281	134	147
Primary Diagnosis	n (missing)	281 (0)	134 (0)	147 (0)
	Certain infectious and parasitic diseases	7 (2.5%)	6 (4.5%)	1 (0.7%)
	Neoplasms	30 (10.7%)	26 (19.4%)	4 (2.7%)
	Diseases of the blood/blood-forming organs, certain disorders involving the immune mechanism	2 (0.7%)	0 (0.0%)	2 (1.4%)
	Endocrine, nutritional and metabolic diseases	10 (3.6%)	0 (0.0%)	10 (6.8%)
	Mental, Behavioral and Neurodevelopmental disorders	2 (0.7%)	0 (0.0%)	2 (1.4%)
	Diseases of the nervous system	12 (4.3%)	4 (3.0%)	8 (5.4%)
	Diseases of the eye and adnexa	9 (3.2%)	4 (3.0%)	5 (3.4%)
	Diseases of the ear and mastoid process	1 (0.4%)	0 (0.0%)	1 (0.7%)
	Diseases of the circulatory system	26 (9.3%)	14 (10.4%)	12 (8.2%)
	Diseases of the respiratory system	10 (3.6%)	3 (2.2%)	7 (4.8%)
	Diseases of the digestive system	41 (14.6%)	17 (12.7%)	24 (16.3%)
	Diseases of the skin and subcutaneous tissue	8 (2.8%)	5 (3.7%)	3 (2.0%)
	Diseases of the musculoskeletal system and connective tissue	33 (11.7%)	13 (9.7%)	20 (13.6%)
	Diseases of the genitourinary system	20 (7.1%)	8 (6.0%)	12 (8.2%)
	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	43 (15.3%)	20 (14.9%)	23 (15.6%)
	Injury, poisoning and certain other consequences of external causes	12 (4.3%)	7 (5.2%)	5 (3.4%)
	Factors influencing health status and contact with health services	15 (5.3%)	7 (5.2%)	8 (5.4%)

Table 2.1.4
Overall hospitalisations during trial, Overall and by Group

		All	Intervention	Control
Within trial comparison (IB vs CB)		88	45	43
Primary Diagnosis	n (missing)	88 (0)	45 (0)	43 (0)
	Certain infectious and parasitic diseases	1 (1.1%)	1 (2.2%)	0 (0.0%)
	Neoplasms	1 (1.1%)	1 (2.2%)	0 (0.0%)
	Endocrine, nutritional and metabolic diseases	2 (2.3%)	1 (2.2%)	1 (2.3%)
	Diseases of the nervous system	6 (6.8%)	5 (11.1%)	1 (2.3%)
	Diseases of the eye and adnexa	2 (2.3%)	1 (2.2%)	1 (2.3%)
	Diseases of the circulatory system	9 (10.2%)	2 (4.4%)	7 (16.3%)
	Diseases of the respiratory system	3 (3.4%)	2 (4.4%)	1 (2.3%)
	Diseases of the digestive system	14 (15.9%)	8 (17.8%)	6 (14.0%)
	Diseases of the skin and subcutaneous tissue	3 (3.4%)	1 (2.2%)	2 (4.7%)
	Diseases of the musculoskeletal system and connective tissue	10 (11.4%)	3 (6.7%)	7 (16.3%)
	Diseases of the genitourinary system	5 (5.7%)	3 (6.7%)	2 (4.7%)
	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	25 (28.4%)	12 (26.7%)	13 (30.2%)
	Injury, poisoning and certain other consequences of external causes	4 (4.5%)	3 (6.7%)	1 (2.3%)
	Factors influencing health status and contact with health services	3 (3.4%)	2 (4.4%)	1 (2.3%)

Table 2.1.5
Summary of Subjects with Hospitalisations, Overall and by Group

	All	Intervention	Control
Number of Subjects with Hospitalisations			
Prior to intervention comparison (IA vs CB)	68	33	35
Within intervention comparison (IB vs CD)	72	34	38
Long term follow-up comparison (IC vs CC)	161	72	89
Within trial comparison (IB vs CB)	69	34	35

Table 2.1.6
Overall hospitalisations pre-FFIT intervention and during intervention, Overall and by Group

		All	Pre-FITT	FITT
Total Hospitalisations				
Pre-FITT vs FITT comparison		184	92	92
Primary Diagnosis	n (missing)	184 (0)	92 (0)	92 (0)
	Certain infectious and parasitic diseases	1 (0.5%)	0 (0.0%)	1 (1.1%)
	Neoplasms	2 (1.1%)	0 (0.0%)	2 (2.2%)
	Diseases of the blood/blood-forming organs, certain disorders involving the immune mechanism	1 (0.5%)	0 (0.0%)	1 (1.1%)
	Endocrine, nutritional and metabolic diseases	3 (1.6%)	1 (1.1%)	2 (2.2%)
	Diseases of the nervous system	16 (8.7%)	8 (8.7%)	8 (8.7%)
	Diseases of the eye and adnexa	7 (3.8%)	2 (2.2%)	5 (5.4%)
	Diseases of the circulatory system	23 (12.5%)	17 (18.5%)	6 (6.5%)
	Diseases of the respiratory system	7 (3.8%)	4 (4.3%)	3 (3.3%)
	Diseases of the digestive system	29 (15.8%)	12 (13.0%)	17 (18.5%)
	Diseases of the skin and subcutaneous tissue	5 (2.7%)	3 (3.3%)	2 (2.2%)
	Diseases of the musculoskeletal system and connective tissue	27 (14.7%)	16 (17.4%)	11 (12.0%)
	Diseases of the genitourinary system	11 (6.0%)	6 (6.5%)	5 (5.4%)
	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	39 (21.2%)	19 (20.7%)	20 (21.7%)
	Injury, poisoning and certain other consequences of external causes	7 (3.8%)	3 (3.3%)	4 (4.3%)
	Factors influencing health status and contact with health services	6 (3.3%)	1 (1.1%)	5 (5.4%)
Number of Subjects with Hospitalisations				
Pre-FITT vs FITT comparison		140	68	72

Table 2.2.1
Summary of Hospitalisations I, Overall and by Group

		All	Intervention	Control
Number of Hospitalisations Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	0	577 (89.5%)	277 (89.4%)	300 (89.6%)
	1	52 (8.1%)	24 (7.7%)	28 (8.4%)
	2	12 (1.9%)	6 (1.9%)	6 (1.8%)
	3	1 (0.2%)	0 (0.0%)	1 (0.3%)
	4	2 (0.3%)	2 (0.6%)	0 (0.0%)
	5	1 (0.2%)	1 (0.3%)	0 (0.0%)
	P-value			0.6592*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	0	573 (88.8%)	276 (89.0%)	297 (88.7%)
	1	57 (8.8%)	27 (8.7%)	30 (9.0%)
	2	12 (1.9%)	5 (1.6%)	7 (2.1%)
	3	2 (0.3%)	1 (0.3%)	1 (0.3%)
	5	1 (0.2%)	1 (0.3%)	0 (0.0%)
	P-value			0.9460*

Table 2.2.1
Summary of Hospitalisations I, Overall and by Group

		All	Intervention	Control
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	0	484 (75.0%)	238 (76.8%)	246 (73.4%)
	1	98 (15.2%)	48 (15.5%)	50 (14.9%)
	2	39 (6.0%)	13 (4.2%)	26 (7.8%)
	3	15 (2.3%)	5 (1.6%)	10 (3.0%)
	4	5 (0.8%)	3 (1.0%)	2 (0.6%)
	5	1 (0.2%)	1 (0.3%)	0 (0.0%)
	6	1 (0.2%)	1 (0.3%)	0 (0.0%)
	7	1 (0.2%)	0 (0.0%)	1 (0.3%)
	22	1 (0.2%)	1 (0.3%)	0 (0.0%)
	P-value			0.2463*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	0	576 (89.3%)	276 (89.0%)	300 (89.6%)
	1	55 (8.5%)	27 (8.7%)	28 (8.4%)
	2	11 (1.7%)	5 (1.6%)	6 (1.8%)
	3	2 (0.3%)	1 (0.3%)	1 (0.3%)
	5	1 (0.2%)	1 (0.3%)	0 (0.0%)
		P-value		

Table 2.2.2
Summary of Hospitalisations II, Overall and by Group

		All	Intervention	Control
Any Hospitalisations				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	68 (10.5%)	33 (10.6%)	35 (10.4%)
	No	577 (89.5%)	277 (89.4%)	300 (89.6%)
	P-value			0.9350
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	72 (11.2%)	34 (11.0%)	38 (11.3%)
	No	573 (88.8%)	276 (89.0%)	297 (88.7%)
	P-value			0.8797
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	161 (25.0%)	72 (23.2%)	89 (26.6%)
	No	484 (75.0%)	238 (76.8%)	246 (73.4%)
	P-value			0.3273
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	69 (10.7%)	34 (11.0%)	35 (10.4%)
	No	576 (89.3%)	276 (89.0%)	300 (89.6%)
	P-value			0.8310

Table 2.2.3
Summary of Hospitalisations III, Overall and by Group

		All	Pre-FITT	FITT
Number of Hospitalisations Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	0	1150 (89.1%)	577 (89.5%)	573 (88.8%)
	1	109 (8.4%)	52 (8.1%)	57 (8.8%)
	2	24 (1.9%)	12 (1.9%)	12 (1.9%)
	3	3 (0.2%)	1 (0.2%)	2 (0.3%)
	4	2 (0.2%)	2 (0.3%)	0 (0.0%)
	5	2 (0.2%)	1 (0.2%)	1 (0.2%)
	P-value			0.8241**
Any Hospitalisations				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	140 (10.9%)	68 (10.5%)	72 (11.2%)
	No	1150 (89.1%)	577 (89.5%)	573 (88.8%)
	P-value			0.7029**

Table 3
Psychiatric Hospital Admissions, Overall and by Group

		All	Intervention	Control
Total Psychiatirc Hospital Admissions				
Prior to intervention comparison (IA vs CB)		0	0	0
Within intervention comparison (IB vs CD)		0	0	0
Long term follow-up comparison (IC vs CC)		6	3	3
Primary Diagnosis	n (missing)	6	3	3
	F209 - Schizophrenia	2	0	2
	Z032 - Observation for suspected mental and behavioural disorders	1	1	0
	Z004 - General psychiatric examination, not elsewhere classified	1	1	0
	F4322 - Adjustment disorders	1	0	1
	F323 - Severe depressive episode with psychotic symptoms	1	1	0
Within trial comparison (IB vs CB)		0	0	0
Number of Subjects with Psychiatric Hospital Admissions				
Prior to intervention comparison (IA vs CB)		0	0	0
Within intervention comparison (IB vs CD)		0	0	0
Long term follow-up comparison (IC vs CC)		4	2	2
Within trial comparison (IB vs CB)		0	0	0

Table 4.1
Cancer Incidences, Overall and by Group

		All	Intervention	Control
Total Cancer Incidence				
Prior to intervention comparison (IA vs CB)		0	0	0
Within intervention comparison (IB vs CD)		2	0	2
Primary Diagnosis	n (missing)	2	0	2
	C924 - Malignant neoplasm: myeloid leukaemia	1	0	1
	C61X - Malignant neoplasm: prostate	1	0	1
Long term follow-up comparison (IC vs CC)		9	5	4
Primary Diagnosis	n (missing)	9	5	4
	C924 - Malignant neoplasm: myeloid leukaemia	1	0	1
	C445 - Malignant neoplasms: skin, trunk	1	0	1
	C20X - Malignant neoplasm: rectum	1	0	1
	C61X - Malignant neoplasm: prostate	2	1	1
	D075 - In situ neoplasm: carcinoma of prostate	1	1	0
	C435 - Malignant melanoma: skin, trunk	1	1	0
	C601 - Malignant neoplasm: glans penis	1	1	0
	D469 - Neoplasms of uncertain or unknown behaviour: myelodysplastic syndrome	1	1	0
Within trial comparison (IB vs CB)		0	0	0

Table 4.2
Cancer Incidences, Overall and by Group

	All	Intervention	Control
Number of Subjects with Cancer Incidences			
Prior to intervention comparison (IA vs CB)	0	0	0
Within intervention comparison (IB vs CD)	2	0	2
Long term follow-up comparison (IC vs CC)	9	5	4
Within trial comparison (IB vs CB)	0	0	0

Table 5
Deaths, Overall and by Group

		All	Intervention	Control
Number of Deaths		0	0	0
Prior to intervention comparison (IA vs CB)		0	0	0
Within intervention comparison (IB vs CD)		1	0	1
Primary Diagnosis	C924 - Acute promyelocytic leukemia	1	0	1
Long term follow-up comparison (IC vs CC)		1	0	1
Primary Diagnosis	C924 - Acute promyelocytic leukemia	1	0	1
Within trial comparison (IB vs CB)		0	0	0

Table 6
Stroke or MI During Period B and C, Overall and by Group

	All	Intervention	Control
Stroke or MI During Period B and C	4	3	1

Table 7.1.1
Overall Prescriptions I, Overall and by Group

		All	Intervention	Control
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		8047	3379	4668
BNF Chapter	n (missing)	8047 (0)	3379 (0)	4668 (0)
	1 Gastro-intestinal system	606 (7.5%)	291 (8.6%)	315 (6.7%)
	2 Cardiovascular system	2642 (32.8%)	1144 (33.9%)	1498 (32.1%)
	3 Respiratory system	617 (7.7%)	253 (7.5%)	364 (7.8%)
	4 Central nervous system	1436 (17.8%)	526 (15.6%)	910 (19.5%)
	5 Infections	348 (4.3%)	143 (4.2%)	205 (4.4%)
	6 Endocrine system	886 (11.0%)	393 (11.6%)	493 (10.6%)
	7 Obstetrics, gynaecology, and urinary-tract disorders	200 (2.5%)	78 (2.3%)	122 (2.6%)
	8 Malignant disease and immunosuppression	15 (0.2%)	3 (0.1%)	12 (0.3%)
	9 Nutrition and blood	65 (0.8%)	42 (1.2%)	23 (0.5%)
	10 Musculoskeletal and joint diseases	438 (5.4%)	186 (5.5%)	252 (5.4%)
	11 Eye	74 (0.9%)	26 (0.8%)	48 (1.0%)
	12 Ear, nose, and oropharynx	148 (1.8%)	49 (1.5%)	99 (2.1%)
	13 Skin	350 (4.3%)	164 (4.9%)	186 (4.0%)
	14 Immunological products and vaccines	20 (0.2%)	8 (0.2%)	12 (0.3%)
	15 Anaesthesia	5 (0.1%)	3 (0.1%)	2 (0.0%)
	Others	197 (2.4%)	70 (2.1%)	127 (2.7%)

Table 7.1.2
Overall Prescriptions II, Overall and by Group

		All	Intervention	Control
Within intervention comparison (IB vs CD)		8750	3692	5058
BNF Chapter	n (missing)	8750 (0)	3692 (0)	5058 (0)
	1 Gastro-intestinal system	637 (7.3%)	298 (8.1%)	339 (6.7%)
	2 Cardiovascular system	3043 (34.8%)	1358 (36.8%)	1685 (33.3%)
	3 Respiratory system	627 (7.2%)	220 (6.0%)	407 (8.0%)
	4 Central nervous system	1503 (17.2%)	605 (16.4%)	898 (17.8%)
	5 Infections	318 (3.6%)	128 (3.5%)	190 (3.8%)
	6 Endocrine system	957 (10.9%)	391 (10.6%)	566 (11.2%)
	7 Obstetrics, gynaecology, and urinary-tract disorders	263 (3.0%)	80 (2.2%)	183 (3.6%)
	8 Malignant disease and immunosuppression	28 (0.3%)	0 (0.0%)	28 (0.6%)
	9 Nutrition and blood	72 (0.8%)	54 (1.5%)	18 (0.4%)
	10 Musculoskeletal and joint diseases	510 (5.8%)	220 (6.0%)	290 (5.7%)
	11 Eye	97 (1.1%)	49 (1.3%)	48 (0.9%)
	12 Ear, nose, and oropharynx	130 (1.5%)	43 (1.2%)	87 (1.7%)
	13 Skin	361 (4.1%)	172 (4.7%)	189 (3.7%)
	14 Immunological products and vaccines	8 (0.1%)	4 (0.1%)	4 (0.1%)
	15 Anaesthesia	2 (0.0%)	1 (0.0%)	1 (0.0%)
	Others	194 (2.2%)	69 (1.9%)	125 (2.5%)

Table 7.1.3
Overall Prescriptions III, Overall and by Group

		All	Intervention	Control
Long term follow-up comparison (IC vs CC)		23965	10697	13268
BNF Chapter	n (missing)	23965 (0)	10697 (0)	13268 (0)
	1 Gastro-intestinal system	1750 (7.3%)	895 (8.4%)	855 (6.4%)
	2 Cardiovascular system	8174 (34.1%)	3714 (34.7%)	4460 (33.6%)
	3 Respiratory system	1705 (7.1%)	709 (6.6%)	996 (7.5%)
	4 Central nervous system	4079 (17.0%)	1632 (15.3%)	2447 (18.4%)
	5 Infections	863 (3.6%)	364 (3.4%)	499 (3.8%)
	6 Endocrine system	2611 (10.9%)	1139 (10.6%)	1472 (11.1%)
	7 Obstetrics, gynaecology, and urinary-tract disorders	904 (3.8%)	406 (3.8%)	498 (3.8%)
	8 Malignant disease and immunosuppression	88 (0.4%)	11 (0.1%)	77 (0.6%)
	9 Nutrition and blood	227 (0.9%)	157 (1.5%)	70 (0.5%)
	10 Musculoskeletal and joint diseases	1377 (5.7%)	647 (6.0%)	730 (5.5%)
	11 Eye	215 (0.9%)	101 (0.9%)	114 (0.9%)
	12 Ear, nose, and oropharynx	418 (1.7%)	145 (1.4%)	273 (2.1%)
	13 Skin	924 (3.9%)	475 (4.4%)	449 (3.4%)
	14 Immunological products and vaccines	27 (0.1%)	20 (0.2%)	7 (0.1%)
	15 Anaesthesia	18 (0.1%)	2 (0.0%)	16 (0.1%)
	Others	585 (2.4%)	280 (2.6%)	305 (2.3%)

Table 7.1.4
Overall Prescriptions IV, Overall and by Group

		All	Intervention	Control
Within trial comparison (IB vs CB)		8360	3692	4668
BNF Chapter	n (missing)	8360 (0)	3692 (0)	4668 (0)
	1 Gastro-intestinal system	613 (7.3%)	298 (8.1%)	315 (6.7%)
	2 Cardiovascular system	2856 (34.2%)	1358 (36.8%)	1498 (32.1%)
	3 Respiratory system	584 (7.0%)	220 (6.0%)	364 (7.8%)
	4 Central nervous system	1515 (18.1%)	605 (16.4%)	910 (19.5%)
	5 Infections	333 (4.0%)	128 (3.5%)	205 (4.4%)
	6 Endocrine system	884 (10.6%)	391 (10.6%)	493 (10.6%)
	7 Obstetrics, gynaecology, and urinary-tract disorders	202 (2.4%)	80 (2.2%)	122 (2.6%)
	8 Malignant disease and immunosuppression	12 (0.1%)	0 (0.0%)	12 (0.3%)
	9 Nutrition and blood	77 (0.9%)	54 (1.5%)	23 (0.5%)
	10 Musculoskeletal and joint diseases	472 (5.6%)	220 (6.0%)	252 (5.4%)
	11 Eye	97 (1.2%)	49 (1.3%)	48 (1.0%)
	12 Ear, nose, and oropharynx	142 (1.7%)	43 (1.2%)	99 (2.1%)
	13 Skin	358 (4.3%)	172 (4.7%)	186 (4.0%)
	14 Immunological products and vaccines	16 (0.2%)	4 (0.1%)	12 (0.3%)
	15 Anaesthesia	3 (0.0%)	1 (0.0%)	2 (0.0%)
	Others	196 (2.3%)	69 (1.9%)	127 (2.7%)

Table 7.1.5
Summary of Subjects with Prescriptions, Overall and by Group

	All	Intervention	Control
Number of Subjects with Prescriptions			
Prior to intervention comparison (IA vs CB)	522	241	281
Within intervention comparison (IB vs CD)	513	241	272
Long term follow-up comparison (IC vs CC)	574	269	305
Within trial comparison (IB vs CB)	522	241	281

Table 7.1.6
Overall Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Total Prescriptions				
Pre-FITT vs FITT comparison		16797	8047	8750
BNF Chapter	n (missing)	16797 (0)	8047 (0)	8750 (0)
	1 Gastro-intestinal system	1243 (7.4%)	606 (7.5%)	637 (7.3%)
	2 Cardiovascular system	5685 (33.8%)	2642 (32.8%)	3043 (34.8%)
	3 Respiratory system	1244 (7.4%)	617 (7.7%)	627 (7.2%)
	4 Central nervous system	2939 (17.5%)	1436 (17.8%)	1503 (17.2%)
	5 Infections	666 (4.0%)	348 (4.3%)	318 (3.6%)
	6 Endocrine system	1843 (11.0%)	886 (11.0%)	957 (10.9%)
	7 Obstetrics, gynaecology, and urinary-tract disorders	463 (2.8%)	200 (2.5%)	263 (3.0%)
	8 Malignant disease and immunosuppression	43 (0.3%)	15 (0.2%)	28 (0.3%)
	9 Nutrition and blood	137 (0.8%)	65 (0.8%)	72 (0.8%)
	10 Musculoskeletal and joint diseases	948 (5.6%)	438 (5.4%)	510 (5.8%)
	11 Eye	171 (1.0%)	74 (0.9%)	97 (1.1%)
	12 Ear, nose, and oropharynx	278 (1.7%)	148 (1.8%)	130 (1.5%)
	13 Skin	711 (4.2%)	350 (4.3%)	361 (4.1%)
	14 Immunological products and vaccines	28 (0.2%)	20 (0.2%)	8 (0.1%)
	15 Anaesthesia	7 (0.0%)	5 (0.1%)	2 (0.0%)
	Others	391 (2.3%)	197 (2.4%)	194 (2.2%)
Number of Subjects with Prescriptions				
Pre-FITT vs FITT comparison		1035	522	513

Table 7.2.1
Summary of All Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	12.5 (16.3)	10.9 (14.6)	13.9 (17.6)
	Median (IQR)	6 [1 , 18]	4 [1 , 15]	7 [1 , 21]
	Min - Max	0 , 83	0 , 79	0 , 83
	P-value			0.0200*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	13.6 (17.6)	11.9 (15.9)	15.1 (19.0)
	Median (IQR)	6 [1 , 19]	6 [1 , 18]	7 [2 , 23]
	Min - Max	0 , 96	0 , 77	0 , 96
	P-value			0.0313*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	37.2 (45.8)	34.5 (42.7)	39.6 (48.4)
	Median (IQR)	20 [4 , 54]	19 [3 , 52]	21 [5 , 56]
	Min - Max	0 , 270	0 , 270	0 , 234
	P-value			0.1524*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	13.0 (16.8)	11.9 (15.9)	13.9 (17.6)
	Median (IQR)	6 [1 , 19]	6 [1 , 18]	7 [1 , 21]
	Min - Max	0 , 83	0 , 77	0 , 83
	P-value			0.0803*

Table 7.2.2
Summary of All Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	522 (80.9%)	241 (77.7%)	281 (83.9%)
	No	123 (19.1%)	69 (22.3%)	54 (16.1%)
	P-value			0.0474
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	513 (79.5%)	241 (77.7%)	272 (81.2%)
	No	132 (20.5%)	69 (22.3%)	63 (18.8%)
	P-value			0.2776
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	574 (89.0%)	269 (86.8%)	305 (91.0%)
	No	71 (11.0%)	41 (13.2%)	30 (9.0%)
	P-value			0.0834
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	522 (80.9%)	241 (77.7%)	281 (83.9%)
	No	123 (19.1%)	69 (22.3%)	54 (16.1%)
	P-value			0.0474

Table 7.2.3
Summary of All Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	13.0 (17.0)	12.5 (16.3)	13.6 (17.6)
	Median (IQR)	6 [1 , 19]	6 [1 , 18]	6 [1 , 19]
	Min - Max	0 , 96	0 , 83	0 , 96
	P-value			0.0025**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	1035 (80.2%)	522 (80.9%)	513 (79.5%)
	No	255 (19.8%)	123 (19.1%)	132 (20.5%)
	P-value			0.4013**

Table 7.2.4
Summary of All Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient In Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	11.4 (15.2)	10.9 (14.6)	11.9 (15.9)
	Median (IQR)	5 [1 , 17]	4 [1 , 15]	6 [1 , 18]
	Min - Max	0 , 79	0 , 79	0 , 77
	P-value			0.0355**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	482 (77.7%)	241 (77.7%)	241 (77.7%)
	No	138 (22.3%)	69 (22.3%)	69 (22.3%)
	P-value			1.0000**

Table 7.2.5
Summary of All Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient In Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	14.5 (18.3)	13.9 (17.6)	15.1 (19.0)
	Median (IQR)	7 [1 , 21]	7 [1 , 21]	7 [2 , 23]
	Min - Max	0 , 96	0 , 83	0 , 96
	P-value			0.0265**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	553 (82.5%)	281 (83.9%)	272 (81.2%)
	No	117 (17.5%)	54 (16.1%)	63 (18.8%)
	P-value			0.2164**

Table 7.3.1
Summary of Chapter 1: Gastro-intestinal System Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (2.4)	0.9 (2.2)	0.9 (2.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 15	0 , 29
	P-value			0.9427*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (2.6)	1.0 (2.5)	1.0 (2.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 27	0 , 20	0 , 27
	P-value			0.6582*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.7 (6.6)	2.9 (6.8)	2.6 (6.4)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 2]
	Min - Max	0 , 66	0 , 66	0 , 58
	P-value			0.8758*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (2.6)	1.0 (2.5)	0.9 (2.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 20	0 , 29
	P-value			0.6796*

Table 7.3.2
Summary of Chapter 1: Gastro-intestinal System Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	146 (22.6%)	70 (22.6%)	76 (22.7%)
	No	499 (77.4%)	240 (77.4%)	259 (77.3%)
	P-value			0.9744
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	139 (21.6%)	64 (20.6%)	75 (22.4%)
	No	506 (78.4%)	246 (79.4%)	260 (77.6%)
	P-value			0.5907
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	222 (34.4%)	104 (33.5%)	118 (35.2%)
	No	423 (65.6%)	206 (66.5%)	217 (64.8%)
	P-value			0.6545
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	140 (21.7%)	64 (20.6%)	76 (22.7%)
	No	505 (78.3%)	246 (79.4%)	259 (77.3%)
	P-value			0.5298

Table 7.3.3
Summary of Chapter 1: Gastro-intestinal System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.0 (2.5)	0.9 (2.4)	1.0 (2.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 29	0 , 27
	P-value			0.3963**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	285 (22.1%)	146 (22.6%)	139 (21.6%)
	No	1005 (77.9%)	499 (77.4%)	506 (78.4%)
	P-value			0.4726**

Table 7.3.4
Summary of Chapter 1: Gastro-intestinal System Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	1.0 (2.3)	0.9 (2.2)	1.0 (2.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 15	0 , 20
	P-value			0.9021**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	134 (21.6%)	70 (22.6%)	64 (20.6%)
	No	486 (78.4%)	240 (77.4%)	246 (79.4%)
	P-value			0.3428**

Table 7.3.5
Summary of Chapter 1: Gastro-intestinal System Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	1.0 (2.7)	0.9 (2.6)	1.0 (2.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 29	0 , 27
	P-value			0.2879**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	151 (22.5%)	76 (22.7%)	75 (22.4%)
	No	519 (77.5%)	259 (77.3%)	260 (77.6%)
	P-value			0.8927**

Table 7.4.1
Summary of Chapter 2: Cardiovascular System Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	4.1 (8.2)	3.7 (7.5)	4.5 (8.9)
	Median (IQR)	0 [0 , 5]	0 [0 , 4]	0 [0 , 6]
	Min - Max	0 , 61	0 , 46	0 , 61
	P-value			0.4280*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	4.7 (9.5)	4.4 (9.2)	5.0 (9.7)
	Median (IQR)	0 [0 , 6]	0 [0 , 6]	0 [0 , 7]
	Min - Max	0 , 65	0 , 65	0 , 65
	P-value			0.2267*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	12.7 (22.8)	12.0 (20.5)	13.3 (24.8)
	Median (IQR)	0 [0 , 17]	0 [0 , 17]	0 [0 , 21]
	Min - Max	0 , 165	0 , 96	0 , 165
	P-value			0.9556*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	4.4 (9.0)	4.4 (9.2)	4.5 (8.9)
	Median (IQR)	0 [0 , 6]	0 [0 , 6]	0 [0 , 6]
	Min - Max	0 , 65	0 , 65	0 , 61
	P-value			0.5062*

Table 7.4.2
Summary of Chapter 2: Cardiovascular System Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	209 (32.4%)	97 (31.3%)	112 (33.4%)
	No	436 (67.6%)	213 (68.7%)	223 (66.6%)
	P-value			0.5613
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	211 (32.7%)	94 (30.3%)	117 (34.9%)
	No	434 (67.3%)	216 (69.7%)	218 (65.1%)
	P-value			0.2132
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	269 (41.7%)	130 (41.9%)	139 (41.5%)
	No	376 (58.3%)	180 (58.1%)	196 (58.5%)
	P-value			0.9092
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	206 (31.9%)	94 (30.3%)	112 (33.4%)
	No	439 (68.1%)	216 (69.7%)	223 (66.6%)
	P-value			0.3973

Table 7.4.3
Summary of Chapter 2: Cardiovascular System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	4.4 (8.9)	4.1 (8.2)	4.7 (9.5)
	Median (IQR)	0 [0 , 6]	0 [0 , 5]	0 [0 , 6]
	Min - Max	0 , 65	0 , 61	0 , 65
	P-value			<0.0001**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	420 (32.6%)	209 (32.4%)	211 (32.7%)
	No	870 (67.4%)	436 (67.6%)	434 (67.3%)
	P-value			0.7316**

Table 7.4.4
Summary of Chapter 2: Cardiovascular System Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	4.0 (8.4)	3.7 (7.5)	4.4 (9.2)
	Median (IQR)	0 [0 , 5]	0 [0 , 4]	0 [0 , 6]
	Min - Max	0 , 65	0 , 46	0 , 65
	P-value			0.0048**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	191 (30.8%)	97 (31.3%)	94 (30.3%)
	No	429 (69.2%)	213 (68.7%)	216 (69.7%)
	P-value			0.5127**

Table 7.4.5
Summary of Chapter 2: Cardiovascular System Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	4.8 (9.3)	4.5 (8.9)	5.0 (9.7)
	Median (IQR)	0 [0 , 7]	0 [0 , 6]	0 [0 , 7]
	Min - Max	0 , 65	0 , 61	0 , 65
	P-value			0.0053**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	229 (34.2%)	112 (33.4%)	117 (34.9%)
	No	441 (65.8%)	223 (66.6%)	218 (65.1%)
	P-value			0.1655**

Table 7.5.1
Summary of Chapter 3: Respiratory System Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (3.4)	0.8 (2.9)	1.1 (3.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 32	0 , 30	0 , 32
	P-value			0.1276*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (3.5)	0.7 (2.9)	1.2 (3.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 40	0 , 33
	P-value			0.1313*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.6 (8.9)	2.3 (7.8)	3.0 (9.7)
	Median (IQR)	0 [0 , 1]	0 [0 , 0]	0 [0 , 1]
	Min - Max	0 , 93	0 , 93	0 , 91
	P-value			0.1836*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (3.4)	0.7 (2.9)	1.1 (3.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 40	0 , 32
	P-value			0.1234*

Table 7.5.2
Summary of Chapter 3: Respiratory System Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	119 (18.4%)	49 (15.8%)	70 (20.9%)
	No	526 (81.6%)	261 (84.2%)	265 (79.1%)
	P-value			0.0960
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	118 (18.3%)	50 (16.1%)	68 (20.3%)
	No	527 (81.7%)	260 (83.9%)	267 (79.7%)
	P-value			0.1712
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	174 (27.0%)	76 (24.5%)	98 (29.3%)
	No	471 (73.0%)	234 (75.5%)	237 (70.7%)
	P-value			0.1756
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	120 (18.6%)	50 (16.1%)	70 (20.9%)
	No	525 (81.4%)	260 (83.9%)	265 (79.1%)
	P-value			0.1201

Table 7.5.3
Summary of Chapter 3: Respiratory System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.0 (3.4)	1.0 (3.4)	1.0 (3.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 32	0 , 40
	P-value			0.9003**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	237 (18.4%)	119 (18.4%)	118 (18.3%)
	No	1053 (81.6%)	526 (81.6%)	527 (81.7%)
	P-value			0.9013**

Table 7.5.4
Summary of Chapter 3: Respiratory System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.8 (2.9)	0.8 (2.9)	0.7 (2.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 30	0 , 40
	P-value			0.1283**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	99 (16.0%)	49 (15.8%)	50 (16.1%)
	No	521 (84.0%)	261 (84.2%)	260 (83.9%)
	P-value			0.8527**

Table 7.5.5
Summary of Chapter 3: Respiratory System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	1.2 (3.9)	1.1 (3.8)	1.2 (3.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 33	0 , 32	0 , 33
	P-value			0.2390**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	138 (20.6%)	70 (20.9%)	68 (20.3%)
	No	532 (79.4%)	265 (79.1%)	267 (79.7%)
	P-value			0.7389**

Table 7.6.1
Summary of Chapter 4: Central Nervous System Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.2 (5.8)	1.7 (4.5)	2.7 (6.7)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 2]
	Min - Max	0 , 60	0 , 40	0 , 60
	P-value			0.0235*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.3 (6.0)	2.0 (5.0)	2.7 (6.8)
	Median (IQR)	0 [0 , 2]	0 [0 , 1]	0 [0 , 2]
	Min - Max	0 , 66	0 , 40	0 , 66
	P-value			0.2195*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	6.3 (15.2)	5.3 (11.9)	7.3 (17.6)
	Median (IQR)	0 [0 , 4]	0 [0 , 3]	1 [0 , 6]
	Min - Max	0 , 165	0 , 77	0 , 165
	P-value			0.0281*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.3 (5.9)	2.0 (5.0)	2.7 (6.7)
	Median (IQR)	0 [0 , 2]	0 [0 , 1]	0 [0 , 2]
	Min - Max	0 , 60	0 , 40	0 , 60
	P-value			0.1388*

Table 7.6.2
Summary of Chapter 4: Central Nervous System Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	219 (34.0%)	92 (29.7%)	127 (37.9%)
	No	426 (66.0%)	218 (70.3%)	208 (62.1%)
	P-value			0.0274
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	222 (34.4%)	101 (32.6%)	121 (36.1%)
	No	423 (65.6%)	209 (67.4%)	214 (63.9%)
	P-value			0.3446
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	317 (49.1%)	139 (44.8%)	178 (53.1%)
	No	328 (50.9%)	171 (55.2%)	157 (46.9%)
	P-value			0.0352
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	228 (35.3%)	101 (32.6%)	127 (37.9%)
	No	417 (64.7%)	209 (67.4%)	208 (62.1%)
	P-value			0.1572

Table 7.6.3
Summary of Chapter 4: Central Nervous System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	2.3 (5.9)	2.2 (5.8)	2.3 (6.0)
	Median (IQR)	0 [0 , 2]	0 [0 , 1]	0 [0 , 2]
	Min - Max	0 , 66	0 , 60	0 , 66
	P-value			0.6219**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	441 (34.2%)	219 (34.0%)	222 (34.4%)
	No	849 (65.8%)	426 (66.0%)	423 (65.6%)
	P-value			0.7991**

Table 7.6.4
Summary of Chapter 4: Central Nervous System Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	1.8 (4.8)	1.7 (4.5)	2.0 (5.0)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 40	0 , 40	0 , 40
	P-value			0.1425**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	193 (31.1%)	92 (29.7%)	101 (32.6%)
	No	427 (68.9%)	218 (70.3%)	209 (67.4%)
	P-value			0.2715**

Table 7.6.5
Summary of Chapter 4: Central Nervous System Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	2.7 (6.7)	2.7 (6.7)	2.7 (6.8)
	Median (IQR)	0 [0 , 2]	0 [0 , 2]	0 [0 , 2]
	Min - Max	0 , 66	0 , 60	0 , 66
	P-value			0.5559**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	248 (37.0%)	127 (37.9%)	121 (36.1%)
	No	422 (63.0%)	208 (62.1%)	214 (63.9%)
	P-value			0.4795**

Table 7.7.1
Summary of Chapter 5: Infections Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.1)	0.5 (1.0)	0.6 (1.3)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 10	0 , 8	0 , 10
	P-value			0.3019*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.1)	0.4 (0.8)	0.6 (1.3)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 10	0 , 5	0 , 10
	P-value			0.4324*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.3 (2.4)	1.2 (1.9)	1.5 (2.7)
	Median (IQR)	0 [0 , 2]	0 [0 , 2]	1 [0 , 2]
	Min - Max	0 , 26	0 , 15	0 , 26
	P-value			0.3572*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.1)	0.4 (0.8)	0.6 (1.3)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 10	0 , 5	0 , 10
	P-value			0.0779*

Table 7.7.2
Summary of Chapter 5: Infections Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	202 (31.3%)	93 (30.0%)	109 (32.5%)
	No	443 (68.7%)	217 (70.0%)	226 (67.5%)
	P-value			0.4876
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	179 (27.8%)	83 (26.8%)	96 (28.7%)
	No	466 (72.2%)	227 (73.2%)	239 (71.3%)
	P-value			0.5937
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	319 (49.5%)	151 (48.7%)	168 (50.1%)
	No	326 (50.5%)	159 (51.3%)	167 (49.9%)
	P-value			0.7148
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	192 (29.8%)	83 (26.8%)	109 (32.5%)
	No	453 (70.2%)	227 (73.2%)	226 (67.5%)
	P-value			0.1097

Table 7.7.3
Summary of Chapter 5: Infections Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.5 (1.1)	0.5 (1.1)	0.5 (1.1)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 10	0 , 10	0 , 10
	P-value			0.2553**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	381 (29.5%)	202 (31.3%)	179 (27.8%)
	No	909 (70.5%)	443 (68.7%)	466 (72.2%)
	P-value			0.1201**

Table 7.7.4
Summary of Chapter 5: Infections Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.4 (0.9)	0.5 (1.0)	0.4 (0.8)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 8	0 , 8	0 , 5
	P-value			0.5539**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	176 (28.4%)	93 (30.0%)	83 (26.8%)
	No	444 (71.6%)	217 (70.0%)	227 (73.2%)
	P-value			0.3447**

Table 7.7.5
Summary of Chapter 5: Infections Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.6 (1.3)	0.6 (1.3)	0.6 (1.3)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 10	0 , 10	0 , 10
	P-value			0.3190**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	205 (30.6%)	109 (32.5%)	96 (28.7%)
	No	465 (69.4%)	226 (67.5%)	239 (71.3%)
	P-value			0.2088**

Table 7.8.1
Summary of Chapter 6: Endocrine System Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.4 (4.9)	1.3 (4.5)	1.5 (5.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 43	0 , 38	0 , 43
	P-value			0.3892*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.5 (5.3)	1.3 (4.3)	1.7 (6.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 53	0 , 28	0 , 53
	P-value			0.6210*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	4.0 (13.7)	3.7 (12.4)	4.4 (14.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 122	0 , 77	0 , 122
	P-value			0.4201*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.4 (4.8)	1.3 (4.3)	1.5 (5.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 43	0 , 28	0 , 43
	P-value			0.4645*

Table 7.8.2
Summary of Chapter 6: Endocrine System Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	83 (12.9%)	36 (11.6%)	47 (14.0%)
	No	562 (87.1%)	274 (88.4%)	288 (86.0%)
	P-value			0.3597
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	81 (12.6%)	37 (11.9%)	44 (13.1%)
	No	564 (87.4%)	273 (88.1%)	291 (86.9%)
	P-value			0.6462
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	116 (18.0%)	52 (16.8%)	64 (19.1%)
	No	529 (82.0%)	258 (83.2%)	271 (80.9%)
	P-value			0.4414
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	84 (13.0%)	37 (11.9%)	47 (14.0%)
	No	561 (87.0%)	273 (88.1%)	288 (86.0%)
	P-value			0.4298

Table 7.8.3
Summary of Chapter 6: Endocrine System Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.4 (5.1)	1.4 (4.9)	1.5 (5.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 53	0 , 43	0 , 53
	P-value			0.2095**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	164 (12.7%)	83 (12.9%)	81 (12.6%)
	No	1126 (87.3%)	562 (87.1%)	564 (87.4%)
	P-value			0.7150**

Table 7.8.4
Summary of Chapter 6: Endocrine System Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	1.3 (4.4)	1.3 (4.5)	1.3 (4.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 38	0 , 38	0 , 28
	P-value			0.8430**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	73 (11.8%)	36 (11.6%)	37 (11.9%)
	No	547 (88.2%)	274 (88.4%)	273 (88.1%)
	P-value			0.7815**

Table 7.8.5
Summary of Chapter 6: Endocrine System Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	1.6 (5.6)	1.5 (5.2)	1.7 (6.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 53	0 , 43	0 , 53
	P-value			0.1345**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	91 (13.6%)	47 (14.0%)	44 (13.1%)
	No	579 (86.4%)	288 (86.0%)	291 (86.9%)
	P-value			0.4669**

Table 7.9.1

Summary of Chapter 7: Obstetrics Gynaecology and Urinary-Tract Disorders Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.6)	0.3 (1.4)	0.4 (1.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 15	0 , 15
	P-value			0.2731*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.4 (1.9)	0.3 (1.4)	0.5 (2.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 14	0 , 18
	P-value			0.0648*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.4 (5.3)	1.3 (5.2)	1.5 (5.4)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 49	0 , 49	0 , 37
	P-value			0.3699*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.6)	0.3 (1.4)	0.4 (1.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 14	0 , 15
	P-value			0.4602*

Table 7.9.2
Summary of Chapter 7: Obstetrics Gynaecology and Urinary-Tract Disorders Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	38 (5.9%)	15 (4.8%)	23 (6.9%)
	No	607 (94.1%)	295 (95.2%)	312 (93.1%)
	P-value			0.2747
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	48 (7.4%)	17 (5.5%)	31 (9.3%)
	No	597 (92.6%)	293 (94.5%)	304 (90.7%)
	P-value			0.0684
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	81 (12.6%)	35 (11.3%)	46 (13.7%)
	No	564 (87.4%)	275 (88.7%)	289 (86.3%)
	P-value			0.3499
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	40 (6.2%)	17 (5.5%)	23 (6.9%)
	No	605 (93.8%)	293 (94.5%)	312 (93.1%)
	P-value			0.4672

Table 7.9.3

Summary of Chapter 7: Obstetrics Gynaecology and Urinary-Tract Disorders Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.4 (1.7)	0.3 (1.6)	0.4 (1.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 15	0 , 18
	P-value			0.0037**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	86 (6.7%)	38 (5.9%)	48 (7.4%)
	No	1204 (93.3%)	607 (94.1%)	597 (92.6%)
	P-value			0.0412**

Table 7.9.4

Summary of Chapter 7: Obstetrics Gynaecology and Urinary-Tract Disorders Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.3 (1.4)	0.3 (1.4)	0.3 (1.4)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 15	0 , 14
	P-value			0.6836**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	32 (5.2%)	15 (4.8%)	17 (5.5%)
	No	588 (94.8%)	295 (95.2%)	293 (94.5%)
	P-value			0.4795**

Table 7.9.5
Summary of Chapter 7: Obstetrics Gynaecology and Urinary-Tract Disorders Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.5 (2.0)	0.4 (1.7)	0.5 (2.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 15	0 , 18
	P-value			0.0013**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	54 (8.1%)	23 (6.9%)	31 (9.3%)
	No	616 (91.9%)	312 (93.1%)	304 (90.7%)
	P-value			0.0455**

Table 7.10.1

Summary of Chapter 8: Malignant Disease and Immunosuppression Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.5)	0.0 (0.2)	0.0 (0.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 3	0 , 12
	P-value			0.9597*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.8)	0.0 (0.0)	0.1 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 0	0 , 20
	P-value			0.1741*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (2.1)	0.0 (0.6)	0.2 (2.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 48	0 , 11	0 , 48
	P-value			0.3548*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.5)	0.0 (0.0)	0.0 (0.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 0	0 , 12
	P-value			0.3376*

Table 7.10.2

Summary of Chapter 8: Malignant Disease and Immunosuppression Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	2 (0.3%)	1 (0.3%)	1 (0.3%)
	No	643 (99.7%)	309 (99.7%)	334 (99.7%)
	P-value			0.9562
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	2 (0.3%)	0 (0.0%)	2 (0.6%)
	No	643 (99.7%)	310 (100.0%)	333 (99.4%)
	P-value			0.1730
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	4 (0.6%)	1 (0.3%)	3 (0.9%)
	No	641 (99.4%)	309 (99.7%)	332 (99.1%)
	P-value			0.3544
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	1 (0.2%)	0 (0.0%)	1 (0.3%)
	No	644 (99.8%)	310 (100.0%)	334 (99.7%)
	P-value			0.3357

Table 7.10.3

Summary of Chapter 8: Malignant Disease and Immunosuppression Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.0 (0.7)	0.0 (0.5)	0.0 (0.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 12	0 , 20
	P-value			0.5000**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	4 (0.3%)	2 (0.3%)	2 (0.3%)
	No	1286 (99.7%)	643 (99.7%)	643 (99.7%)
	P-value			1.0000**

Table 7.10.4

Summary of Chapter 8: Malignant Disease and Immunosuppression Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.2)	0.0 (0.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 3	0 , 3	0 , 0
	P-value			1.0000**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	1 (0.2%)	1 (0.3%)	0 (0.0%)
	No	619 (99.8%)	309 (99.7%)	310 (100.0%)

Table 7.10.5

Summary of Chapter 8: Malignant Disease and Immunosuppression Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.1 (1.0)	0.0 (0.7)	0.1 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 12	0 , 20
	P-value			0.5000**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	3 (0.4%)	1 (0.3%)	2 (0.6%)
	No	667 (99.6%)	334 (99.7%)	333 (99.4%)
	P-value			0.3173**

Table 7.11.1
Summary of Chapter 9: Nutrition and Blood Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.9)	0.1 (1.1)	0.1 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 16	0 , 16	0 , 8
	P-value			0.6749*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.8)	0.2 (1.1)	0.1 (0.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 9
	P-value			0.1185*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.4 (2.4)	0.5 (2.9)	0.2 (1.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 36	0 , 36	0 , 32
	P-value			0.1960*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.8)	0.2 (1.1)	0.1 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 8
	P-value			0.2807*

Table 7.11.2
Summary of Chapter 9: Nutrition and Blood Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	19 (2.9%)	10 (3.2%)	9 (2.7%)
	No	626 (97.1%)	300 (96.8%)	326 (97.3%)
	P-value			0.6857
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	20 (3.1%)	13 (4.2%)	7 (2.1%)
	No	625 (96.9%)	297 (95.8%)	328 (97.9%)
	P-value			0.1235
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	40 (6.2%)	23 (7.4%)	17 (5.1%)
	No	605 (93.8%)	287 (92.6%)	318 (94.9%)
	P-value			0.2174
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	22 (3.4%)	13 (4.2%)	9 (2.7%)
	No	623 (96.6%)	297 (95.8%)	326 (97.3%)
	P-value			0.2921

Table 7.11.3
Summary of Chapter 9: Nutrition and Blood Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.1 (0.8)	0.1 (0.9)	0.1 (0.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 16	0 , 16	0 , 13
	P-value			0.6015**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	39 (3.0%)	19 (2.9%)	20 (3.1%)
	No	1251 (97.0%)	626 (97.1%)	625 (96.9%)
	P-value			0.8084**

Table 7.11.4
Summary of Chapter 9: Nutrition and Blood Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.2 (1.1)	0.1 (1.1)	0.2 (1.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 16	0 , 16	0 , 13
	P-value			0.2081**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	23 (3.7%)	10 (3.2%)	13 (4.2%)
	No	597 (96.3%)	300 (96.8%)	297 (95.8%)
	P-value			0.3173**

Table 7.11.5
Summary of Chapter 9: Nutrition and Blood Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.1 (0.6)	0.1 (0.6)	0.1 (0.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 9	0 , 8	0 , 9
	P-value			0.4082**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	16 (2.4%)	9 (2.7%)	7 (2.1%)
	No	654 (97.6%)	326 (97.3%)	328 (97.9%)
	P-value			0.4795**

Table 7.12.1

Summary of Chapter 10: Musculoskeletal and Joint Diseases Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.7 (1.9)	0.6 (1.7)	0.8 (2.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 1]
	Min - Max	0 , 24	0 , 15	0 , 24
	P-value			0.1613*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.8 (2.0)	0.7 (1.7)	0.9 (2.2)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 19	0 , 11	0 , 19
	P-value			0.6098*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.1 (5.0)	2.1 (4.9)	2.2 (5.1)
	Median (IQR)	0 [0 , 2]	0 [0 , 2]	0 [0 , 2]
	Min - Max	0 , 51	0 , 40	0 , 51
	P-value			0.3558*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.7 (1.9)	0.7 (1.7)	0.8 (2.1)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 24	0 , 11	0 , 24
	P-value			0.7744*

Table 7.12.2
Summary of Chapter 10: Musculoskeletal and Joint Diseases Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	150 (23.3%)	64 (20.6%)	86 (25.7%)
	No	495 (76.7%)	246 (79.4%)	249 (74.3%)
	P-value			0.1311
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	177 (27.4%)	83 (26.8%)	94 (28.1%)
	No	468 (72.6%)	227 (73.2%)	241 (71.9%)
	P-value			0.7147
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	271 (42.0%)	124 (40.0%)	147 (43.9%)
	No	374 (58.0%)	186 (60.0%)	188 (56.1%)
	P-value			0.3185
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	169 (26.2%)	83 (26.8%)	86 (25.7%)
	No	476 (73.8%)	227 (73.2%)	249 (74.3%)
	P-value			0.7504

Table 7.12.3

Summary of Chapter 10: Musculoskeletal and Joint Diseases Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.7 (1.9)	0.7 (1.9)	0.8 (2.0)
	Median (IQR)	0 [0 , 1]	0 [0 , 0]	0 [0 , 1]
	Min - Max	0 , 24	0 , 24	0 , 19
	P-value			0.0253**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	327 (25.3%)	150 (23.3%)	177 (27.4%)
	No	963 (74.7%)	495 (76.7%)	468 (72.6%)
	P-value			0.0312**

Table 7.12.4
Summary of Chapter 10: Musculoskeletal and Joint Diseases Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.7 (1.7)	0.6 (1.7)	0.7 (1.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 1]
	Min - Max	0 , 15	0 , 15	0 , 11
	P-value			0.0959**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	147 (23.7%)	64 (20.6%)	83 (26.8%)
	No	473 (76.3%)	246 (79.4%)	227 (73.2%)
	P-value			0.0222**

Table 7.12.5
Summary of Chapter 10: Musculoskeletal and Joint Diseases Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.8 (2.1)	0.8 (2.1)	0.9 (2.2)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 24	0 , 24	0 , 19
	P-value			0.1282**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	180 (26.9%)	86 (25.7%)	94 (28.1%)
	No	490 (73.1%)	249 (74.3%)	241 (71.9%)
	P-value			0.3938**

Table 7.13.1
Summary of Chapter 11: Eye Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.9)	0.1 (0.4)	0.1 (1.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 5	0 , 15
	P-value			0.9132*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.2 (1.0)	0.2 (1.0)	0.1 (1.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 14	0 , 15
	P-value			0.5487*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (2.5)	0.3 (1.9)	0.3 (3.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 46	0 , 28	0 , 46
	P-value			0.2137*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.2 (1.0)	0.2 (1.0)	0.1 (1.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 14	0 , 15
	P-value			0.3657*

Table 7.13.2
Summary of Chapter 11: Eye Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	36 (5.6%)	17 (5.5%)	19 (5.7%)
	No	609 (94.4%)	293 (94.5%)	316 (94.3%)
	P-value			0.9173
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	44 (6.8%)	23 (7.4%)	21 (6.3%)
	No	601 (93.2%)	287 (92.6%)	314 (93.7%)
	P-value			0.5625
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	61 (9.5%)	34 (11.0%)	27 (8.1%)
	No	584 (90.5%)	276 (89.0%)	308 (91.9%)
	P-value			0.2073
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	42 (6.5%)	23 (7.4%)	19 (5.7%)
	No	603 (93.5%)	287 (92.6%)	316 (94.3%)
	P-value			0.3688

Table 7.13.3
Summary of Chapter 11: Eye Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.1 (0.9)	0.1 (0.9)	0.2 (1.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 15	0 , 15
	P-value			0.1697**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	80 (6.2%)	36 (5.6%)	44 (6.8%)
	No	1210 (93.8%)	609 (94.4%)	601 (93.2%)
	P-value			0.2935**

Table 7.13.4
Summary of Chapter 11: Eye Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.1 (0.7)	0.1 (0.4)	0.2 (1.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 14	0 , 5	0 , 14
	P-value			0.0439**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	40 (6.5%)	17 (5.5%)	23 (7.4%)
	No	580 (93.5%)	293 (94.5%)	287 (92.6%)
	P-value			0.2568**

Table 7.13.5
Summary of Chapter 11: Eye Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.1 (1.1)	0.1 (1.1)	0.1 (1.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 15	0 , 15	0 , 15
	P-value			0.8980**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	40 (6.0%)	19 (5.7%)	21 (6.3%)
	No	630 (94.0%)	316 (94.3%)	314 (93.7%)
	P-value			0.7150**

Table 7.14.1
Summary of Chapter 12: Ear Nose and Oropharynx Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.2 (1.0)	0.2 (0.7)	0.3 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 8	0 , 11
	P-value			0.2069*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.2 (1.0)	0.1 (0.7)	0.3 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 9	0 , 12
	P-value			0.3890*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.6 (2.3)	0.5 (1.3)	0.8 (2.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 1]
	Min - Max	0 , 29	0 , 9	0 , 29
	P-value			0.2300*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.2 (1.0)	0.1 (0.7)	0.3 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 9	0 , 11
	P-value			0.0411*

Table 7.14.2
Summary of Chapter 12: Ear Nose and Oropharynx Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	72 (11.2%)	30 (9.7%)	42 (12.5%)
	No	573 (88.8%)	280 (90.3%)	293 (87.5%)
	P-value			0.2492
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	56 (8.7%)	24 (7.7%)	32 (9.6%)
	No	589 (91.3%)	286 (92.3%)	303 (90.4%)
	P-value			0.4146
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	155 (24.0%)	68 (21.9%)	87 (26.0%)
	No	490 (76.0%)	242 (78.1%)	248 (74.0%)
	P-value			0.2308
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	66 (10.2%)	24 (7.7%)	42 (12.5%)
	No	579 (89.8%)	286 (92.3%)	293 (87.5%)
	P-value			0.0447

Table 7.14.3
Summary of Chapter 12: Ear Nose and Oropharynx Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.2 (1.0)	0.2 (1.0)	0.2 (1.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 11	0 , 12
	P-value			0.5675**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	128 (9.9%)	72 (11.2%)	56 (8.7%)
	No	1162 (90.1%)	573 (88.8%)	589 (91.3%)
	P-value			0.0558**

Table 7.14.4
Summary of Chapter 12: Ear Nose and Oropharynx Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.1 (0.7)	0.2 (0.7)	0.1 (0.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 9	0 , 8	0 , 9
	P-value			0.9367**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	54 (8.7%)	30 (9.7%)	24 (7.7%)
	No	566 (91.3%)	280 (90.3%)	286 (92.3%)
	P-value			0.3035**

Table 7.14.5
Summary of Chapter 12: Ear Nose and Oropharynx Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.3 (1.2)	0.3 (1.2)	0.3 (1.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 11	0 , 12
	P-value			0.4559**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	74 (11.0%)	42 (12.5%)	32 (9.6%)
	No	596 (89.0%)	293 (87.5%)	303 (90.4%)
	P-value			0.0956**

Table 7.15.1
Summary of Chapter 13: Skin Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.6)	0.5 (1.7)	0.6 (1.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 17	0 , 17	0 , 14
	P-value			0.2368*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.6 (1.9)	0.6 (1.6)	0.6 (2.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 17	0 , 29
	P-value			0.5054*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.4 (4.1)	1.5 (4.4)	1.3 (3.8)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 49	0 , 49	0 , 42
	P-value			0.9045*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.6 (1.6)	0.6 (1.6)	0.6 (1.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 17	0 , 17	0 , 14
	P-value			0.8792*

Table 7.15.2
Summary of Chapter 13: Skin Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	135 (20.9%)	58 (18.7%)	77 (23.0%)
	No	510 (79.1%)	252 (81.3%)	258 (77.0%)
	P-value			0.1824
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	136 (21.1%)	69 (22.3%)	67 (20.0%)
	No	509 (78.9%)	241 (77.7%)	268 (80.0%)
	P-value			0.4824
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	238 (36.9%)	115 (37.1%)	123 (36.7%)
	No	407 (63.1%)	195 (62.9%)	212 (63.3%)
	P-value			0.9203
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	146 (22.6%)	69 (22.3%)	77 (23.0%)
	No	499 (77.4%)	241 (77.7%)	258 (77.0%)
	P-value			0.8255

Table 7.15.3
Summary of Chapter 13: Skin Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.6 (1.7)	0.5 (1.6)	0.6 (1.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 17	0 , 29
	P-value			0.8786**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	271 (21.0%)	135 (20.9%)	136 (21.1%)
	No	1019 (79.0%)	510 (79.1%)	509 (78.9%)
	P-value			0.9298**

Table 7.15.4
Summary of Chapter 13: Skin Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.5 (1.7)	0.5 (1.7)	0.6 (1.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 17	0 , 17	0 , 17
	P-value			0.5348**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	127 (20.5%)	58 (18.7%)	69 (22.3%)
	No	493 (79.5%)	252 (81.3%)	241 (77.7%)
	P-value			0.1521**

Table 7.15.5
Summary of Chapter 13: Skin Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.6 (1.8)	0.6 (1.5)	0.6 (2.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 29	0 , 14	0 , 29
	P-value			0.7672**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	144 (21.5%)	77 (23.0%)	67 (20.0%)
	No	526 (78.5%)	258 (77.0%)	268 (80.0%)
	P-value			0.2320**

Table 7.16.1
Summary of Chapter 14: Immunological Products and Vaccines Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.3)	0.0 (0.2)	0.0 (0.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 4	0 , 2	0 , 4
	P-value			0.9001*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.1)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 2	0 , 2	0 , 2
	P-value			0.9250*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.3)	0.1 (0.4)	0.0 (0.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 3	0 , 3	0 , 2
	P-value			0.0589*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.2)	0.0 (0.1)	0.0 (0.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 4	0 , 2	0 , 4
	P-value			0.3713*

Table 7.16.2
Summary of Chapter 14: Immunological Products and Vaccines Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	12 (1.9%)	6 (1.9%)	6 (1.8%)
	No	633 (98.1%)	304 (98.1%)	329 (98.2%)
	P-value			0.8921
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	6 (0.9%)	3 (1.0%)	3 (0.9%)
	No	639 (99.1%)	307 (99.0%)	332 (99.1%)
	P-value			0.9240
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	17 (2.6%)	12 (3.9%)	5 (1.5%)
	No	628 (97.4%)	298 (96.1%)	330 (98.5%)
	P-value			0.0596
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	9 (1.4%)	3 (1.0%)	6 (1.8%)
	No	636 (98.6%)	307 (99.0%)	329 (98.2%)
	P-value			0.3731

Table 7.16.3
Summary of Chapter 14: Immunological Products and Vaccines Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.0 (0.2)	0.0 (0.3)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 4	0 , 4	0 , 2
	P-value			0.1513**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	18 (1.4%)	12 (1.9%)	6 (0.9%)
	No	1272 (98.6%)	633 (98.1%)	639 (99.1%)
	P-value			0.1573**

Table 7.16.4
Summary of Chapter 14: Immunological Products and Vaccines Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.0 (0.2)	0.0 (0.2)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 2	0 , 2	0 , 2
	P-value			0.4805**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	9 (1.5%)	6 (1.9%)	3 (1.0%)
	No	611 (98.5%)	304 (98.1%)	307 (99.0%)
	P-value			0.3173**

Table 7.16.5
Summary of Chapter 14: Immunological Products and Vaccines Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.0 (0.2)	0.0 (0.3)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 4	0 , 4	0 , 2
	P-value			0.2930**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	9 (1.3%)	6 (1.8%)	3 (0.9%)
	No	661 (98.7%)	329 (98.2%)	332 (99.1%)
	P-value			0.3173**

Table 7.17.1
Summary of Chapter 15: Anaesthesia Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.2)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 3	0 , 3	0 , 1
	P-value			0.6128*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.1)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 1	0 , 1	0 , 1
	P-value			0.9580*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.4)	0.0 (0.1)	0.0 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 10	0 , 2	0 , 10
	P-value			0.2077*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.1)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 1	0 , 1	0 , 1
	P-value			0.6103*

Table 7.17.2
Summary of Chapter 15: Anaesthesia Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	3 (0.5%)	1 (0.3%)	2 (0.6%)
	No	642 (99.5%)	309 (99.7%)	333 (99.4%)
	P-value			0.6088
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	2 (0.3%)	1 (0.3%)	1 (0.3%)
	No	643 (99.7%)	309 (99.7%)	334 (99.7%)
	P-value			0.9562
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	5 (0.8%)	1 (0.3%)	4 (1.2%)
	No	640 (99.2%)	309 (99.7%)	331 (98.8%)
	P-value			0.2074
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	3 (0.5%)	1 (0.3%)	2 (0.6%)
	No	642 (99.5%)	309 (99.7%)	333 (99.4%)
	P-value			0.6088

Table 7.17.3
Summary of Chapter 15: Anaesthesia Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.1)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 3	0 , 3	0 , 1
	P-value			0.5000**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	5 (0.4%)	3 (0.5%)	2 (0.3%)
	No	1285 (99.6%)	642 (99.5%)	643 (99.7%)
	P-value			0.5637**

Table 7.17.4
Summary of Chapter 15: Anaesthesia Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.2)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 3	0 , 3	0 , 1
	P-value			1.0000**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	2 (0.3%)	1 (0.3%)	1 (0.3%)
	No	618 (99.7%)	309 (99.7%)	309 (99.7%)
	P-value			<0.0001**

Table 7.17.5
Summary of Chapter 15: Anaesthesia Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.0 (0.1)	0.0 (0.1)	0.0 (0.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 1	0 , 1	0 , 1
	P-value			1.0000**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	3 (0.4%)	2 (0.6%)	1 (0.3%)
	No	667 (99.6%)	333 (99.4%)	334 (99.7%)
	P-value			0.5637**

Table 7.18.1
Summary of Other Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.5)	0.2 (1.1)	0.4 (1.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 16	0 , 9	0 , 16
	P-value			0.3165*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.6)	0.2 (1.3)	0.4 (1.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 18	0 , 15
	P-value			0.1140*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (4.2)	0.9 (4.2)	0.9 (4.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 51	0 , 51	0 , 39
	P-value			0.5030*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.6)	0.2 (1.3)	0.4 (1.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 18	0 , 16
	P-value			0.3746*

Table 7.18.2
Summary of Other Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	55 (8.5%)	23 (7.4%)	32 (9.6%)
	No	590 (91.5%)	287 (92.6%)	303 (90.4%)
	P-value			0.3325
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	62 (9.6%)	24 (7.7%)	38 (11.3%)
	No	583 (90.4%)	286 (92.3%)	297 (88.7%)
	P-value			0.1211
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	113 (17.5%)	51 (16.5%)	62 (18.5%)
	No	532 (82.5%)	259 (83.5%)	273 (81.5%)
	P-value			0.4926
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	56 (8.7%)	24 (7.7%)	32 (9.6%)
	No	589 (91.3%)	286 (92.3%)	303 (90.4%)
	P-value			0.4146

Table 7.18.3
Summary of Other Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.3 (1.5)	0.3 (1.5)	0.3 (1.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 16	0 , 18
	P-value			0.8548**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	117 (9.1%)	55 (8.5%)	62 (9.6%)
	No	1173 (90.9%)	590 (91.5%)	583 (90.4%)
	P-value			0.3621**

Table 7.18.4
Summary of Other Prescriptions IV, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Mean (SD)	0.2 (1.2)	0.2 (1.1)	0.2 (1.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 18	0 , 9	0 , 18
	P-value			0.8131**
Any Prescriptions in Intervention Group				
Pre-FITT vs FITT comparison	n (missing)	620 (0)	310 (0)	310 (0)
	Yes	47 (7.6%)	23 (7.4%)	24 (7.7%)
	No	573 (92.4%)	287 (92.6%)	286 (92.3%)
	P-value			0.8348**

Table 7.18.5
Summary of Other Prescriptions V, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Mean (SD)	0.4 (1.8)	0.4 (1.8)	0.4 (1.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 16	0 , 16	0 , 15
	P-value			0.9764**
Any Prescriptions in Control Group				
Pre-FITT vs FITT comparison	n (missing)	670 (0)	335 (0)	335 (0)
	Yes	70 (10.4%)	32 (9.6%)	38 (11.3%)
	No	600 (89.6%)	303 (90.4%)	297 (88.7%)
	P-value			0.3173**

Table 7.19.1
Number of BNF paragraphs I, Overall and by Group

		All	Intervention	Control
Total Number of BNF Paragraphs Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	3.4 (3.4)	3.1 (3.3)	3.6 (3.4)
	Median (IQR)	3 [1 , 5]	2 [1 , 4]	3 [1 , 5]
	Min - Max	0 , 22	0 , 20	0 , 22
	P-value			0.0218*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	3.5 (3.4)	3.3 (3.4)	3.7 (3.5)
	Median (IQR)	3 [1 , 5]	2 [1 , 5]	3 [1 , 5]
	Min - Max	0 , 21	0 , 21	0 , 18
	P-value			0.0939*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	5.9 (4.8)	5.7 (4.9)	6.1 (4.8)
	Median (IQR)	5 [2 , 8]	5 [2 , 8]	5 [3 , 8]
	Min - Max	0 , 31	0 , 31	0 , 26
	P-value			0.2221*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	3.5 (3.4)	3.3 (3.4)	3.6 (3.4)
	Median (IQR)	3 [1 , 5]	2 [1 , 5]	3 [1 , 5]
	Min - Max	0 , 22	0 , 21	0 , 22
	P-value			0.0819*

**Table 7.19.2
Number of BNF paragraphs II, Overall and by Group**

		All	Pre-FITT	FITT
Number of BNF Paragraphs Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	3.4 (3.4)	3.4 (3.4)	3.5 (3.4)
	Median (IQR)	3 [1 , 5]	3 [1 , 5]	3 [1 , 5]
	Min - Max	0 , 22	0 , 22	0 , 21
	P-value			0.7345*

Table 7.20.1
Summary of Antidepressants I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (2.8)	0.6 (2.2)	1.1 (3.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 28	0 , 19	0 , 28
	P-value			0.0256*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (2.8)	0.8 (2.8)	1.0 (2.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 20	0 , 20
	P-value			0.1650*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.4 (6.7)	2.0 (6.1)	2.7 (7.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 55	0 , 44	0 , 55
	P-value			0.0564*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (3.0)	0.8 (2.8)	1.1 (3.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 28	0 , 20	0 , 28
	P-value			0.0624*

Table 7.20.2
Summary of Antidepressants II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	96 (14.9%)	36 (11.6%)	60 (17.9%)
	No	549 (85.1%)	274 (88.4%)	275 (82.1%)
	P-value			0.0248
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	93 (14.4%)	38 (12.3%)	55 (16.4%)
	No	552 (85.6%)	272 (87.7%)	280 (83.6%)
	P-value			0.1329
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	122 (18.9%)	49 (15.8%)	73 (21.8%)
	No	523 (81.1%)	261 (84.2%)	262 (78.2%)
	P-value			0.0525
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	98 (15.2%)	38 (12.3%)	60 (17.9%)
	No	547 (84.8%)	272 (87.7%)	275 (82.1%)
	P-value			0.0457
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		564	201	363
Within intervention comparison (IB vs CD)		594	261	333
Long term follow-up comparison (IC vs CC)		1517	613	904
Within trial comparison (IB vs CB)		624	261	363

Table 7.20.3
Summary of Antidepressants III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.9 (2.8)	0.9 (2.8)	0.9 (2.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 28	0 , 28	0 , 20
	P-value			0.5958**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	189 (14.7%)	96 (14.9%)	93 (14.4%)
	No	1101 (85.3%)	549 (85.1%)	552 (85.6%)
	P-value			0.6547**
Total Prescriptions				
Pre-FITT vs FITT comparison		1158	564	594

Table 7.21.1
Summary of Pain Relief Prescriptions I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.6 (2.3)	0.5 (1.8)	0.8 (2.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 19	0 , 21
	P-value			0.1282*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.7 (2.4)	0.5 (1.7)	0.9 (2.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 24	0 , 16	0 , 24
	P-value			0.4671*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.2 (7.0)	2.0 (6.2)	2.3 (7.7)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 73	0 , 61	0 , 73
	P-value			0.5708*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.7 (2.3)	0.5 (1.7)	0.8 (2.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 16	0 , 21
	P-value			0.6411*

Table 7.21.2
Summary of Pain Relief Prescriptions II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	113 (17.5%)	47 (15.2%)	66 (19.7%)
	No	532 (82.5%)	263 (84.8%)	269 (80.3%)
	P-value			0.1296
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	125 (19.4%)	57 (18.4%)	68 (20.3%)
	No	520 (80.6%)	253 (81.6%)	267 (79.7%)
	P-value			0.5395
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	212 (32.9%)	98 (31.6%)	114 (34.0%)
	No	433 (67.1%)	212 (68.4%)	221 (66.0%)
	P-value			0.5138
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	123 (19.1%)	57 (18.4%)	66 (19.7%)
	No	522 (80.9%)	253 (81.6%)	269 (80.3%)
	P-value			0.6712
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		411	149	262
Within intervention comparison (IB vs CD)		452	163	289
Long term follow-up comparison (IC vs CC)		1388	609	779
Within trial comparison (IB vs CB)		425	163	262

Table 7.21.3
Summary of Pain Relief Prescriptions III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.7 (2.4)	0.6 (2.3)	0.7 (2.4)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 24	0 , 21	0 , 24
	P-value			0.1797**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	238 (18.4%)	113 (17.5%)	125 (19.4%)
	No	1052 (81.6%)	532 (82.5%)	520 (80.6%)
	P-value			0.2733**
Total Prescriptions				
Pre-FITT vs FITT comparison		863	411	452

Table 7.22.1
Summary of Statins and Other Cholesterol Lowering Drugs I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.9 (2.1)	0.8 (2.0)	1.0 (2.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 10	0 , 13
	P-value			0.4493*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (2.4)	1.0 (2.3)	1.1 (2.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 13
	P-value			0.7270*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.7 (5.9)	2.6 (5.5)	2.8 (6.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 33	0 , 30	0 , 33
	P-value			0.8413*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (2.3)	1.0 (2.3)	1.0 (2.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 13
	P-value			0.9556*

Table 7.22.2
Summary of Statins and Other Cholesterol Lowering Drugs II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	109 (16.9%)	49 (15.8%)	60 (17.9%)
	No	536 (83.1%)	261 (84.2%)	275 (82.1%)
	P-value			0.4762
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	115 (17.8%)	54 (17.4%)	61 (18.2%)
	No	530 (82.2%)	256 (82.6%)	274 (81.8%)
	P-value			0.7935
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	145 (22.5%)	72 (23.2%)	73 (21.8%)
	No	500 (77.5%)	238 (76.8%)	262 (78.2%)
	P-value			0.6628
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	114 (17.7%)	54 (17.4%)	60 (17.9%)
	No	531 (82.3%)	256 (82.6%)	275 (82.1%)
	P-value			0.8702
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		567	248	319
Within intervention comparison (IB vs CD)		674	308	366
Long term follow-up comparison (IC vs CC)		1764	811	953
Within trial comparison (IB vs CB)		627	308	319

Table 7.22.3
Summary of Statins and Other Cholesterol Lowering Drugs III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.0 (2.3)	0.9 (2.1)	1.0 (2.4)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 13
	P-value			<0.0001**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	224 (17.4%)	109 (16.9%)	115 (17.8%)
	No	1066 (82.6%)	536 (83.1%)	530 (82.2%)
	P-value			0.1088**
Total Prescriptions				
Pre-FITT vs FITT comparison		1241	567	674

Table 7.23.1
Summary of Diabetic Medication I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.1 (4.5)	0.9 (4.2)	1.2 (4.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 38	0 , 40
	P-value			0.3631*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.1 (4.6)	0.9 (3.9)	1.3 (5.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 49	0 , 28	0 , 49
	P-value			0.2647*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	3.0 (12.0)	2.6 (10.9)	3.3 (13.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 118	0 , 77	0 , 118
	P-value			0.2700*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.1 (4.3)	0.9 (3.9)	1.2 (4.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 40	0 , 28	0 , 40
	P-value			0.4528*

Table 7.23.2
Summary of Diabetic Medication II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	48 (7.4%)	20 (6.5%)	28 (8.4%)
	No	597 (92.6%)	290 (93.5%)	307 (91.6%)
	P-value			0.3566
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	52 (8.1%)	21 (6.8%)	31 (9.3%)
	No	593 (91.9%)	289 (93.2%)	304 (90.7%)
	P-value			0.2478
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	63 (9.8%)	26 (8.4%)	37 (11.0%)
	No	582 (90.2%)	284 (91.6%)	298 (89.0%)
	P-value			0.2560
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	49 (7.6%)	21 (6.8%)	28 (8.4%)
	No	596 (92.4%)	289 (93.2%)	307 (91.6%)
	P-value			0.4481
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		680	290	390
Within intervention comparison (IB vs CD)		713	288	425
Long term follow-up comparison (IC vs CC)		1915	814	1101
Within trial comparison (IB vs CB)		678	288	390

Table 7.23.3
Summary of Diabetic Medication III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.1 (4.5)	1.1 (4.5)	1.1 (4.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 49	0 , 40	0 , 49
	P-value			0.4005**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	100 (7.8%)	48 (7.4%)	52 (8.1%)
	No	1190 (92.2%)	597 (92.6%)	593 (91.9%)
	P-value			0.1025**
Total Prescriptions				
Pre-FITT vs FITT comparison		1393	680	713

Table 7.24.1
Summary of NSAIDS I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.8)	0.3 (1.1)	0.6 (2.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 11	0 , 21
	P-value			0.0435*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.7)	0.4 (1.3)	0.6 (2.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 19	0 , 12	0 , 19
	P-value			0.3533*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.6 (5.1)	1.5 (5.2)	1.6 (4.9)
	Median (IQR)	0 [0 , 1]	0 [0 , 1]	0 [0 , 1]
	Min - Max	0 , 61	0 , 61	0 , 50
	P-value			0.2058*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.8)	0.4 (1.3)	0.6 (2.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 12	0 , 21
	P-value			0.4917*

Table 7.24.2
Summary of NSAIDS II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	98 (15.2%)	38 (12.3%)	60 (17.9%)
	No	547 (84.8%)	272 (87.7%)	275 (82.1%)
	P-value			0.0457
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	111 (17.2%)	49 (15.8%)	62 (18.5%)
	No	534 (82.8%)	261 (84.2%)	273 (81.5%)
	P-value			0.3639
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	194 (30.1%)	85 (27.4%)	109 (32.5%)
	No	451 (69.9%)	225 (72.6%)	226 (67.5%)
	P-value			0.1567
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	109 (16.9%)	49 (15.8%)	60 (17.9%)
	No	536 (83.1%)	261 (84.2%)	275 (82.1%)
	P-value			0.4762
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		294	92	202
Within intervention comparison (IB vs CD)		323	126	197
Long term follow-up comparison (IC vs CC)		1000	470	530
Within trial comparison (IB vs CB)		328	126	202

Table 7.24.3
Summary of NSAIDS III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.5 (1.7)	0.5 (1.8)	0.5 (1.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 21	0 , 19
	P-value			0.0901**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	209 (16.2%)	98 (15.2%)	111 (17.2%)
	No	1081 (83.8%)	547 (84.8%)	534 (82.8%)
	P-value			0.2172**
Total Prescriptions				
Pre-FITT vs FITT comparison		617	294	323

Table 7.25.1
Summary of Aspirin I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.3)	0.3 (1.3)	0.3 (1.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 10	0 , 7	0 , 10
	P-value			0.6717*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.4 (1.5)	0.4 (1.7)	0.4 (1.4)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 12
	P-value			0.8239*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.0 (3.7)	1.0 (3.6)	1.0 (3.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 32	0 , 17	0 , 32
	P-value			0.7389*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.4 (1.5)	0.4 (1.7)	0.3 (1.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 13	0 , 10
	P-value			0.9479*

Table 7.25.2
Summary of Aspirin II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	47 (7.3%)	21 (6.8%)	26 (7.8%)
	No	598 (92.7%)	289 (93.2%)	309 (92.2%)
	P-value			0.6299
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	50 (7.8%)	23 (7.4%)	27 (8.1%)
	No	595 (92.2%)	287 (92.6%)	308 (91.9%)
	P-value			0.7612
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	62 (9.6%)	31 (10.0%)	31 (9.3%)
	No	583 (90.4%)	279 (90.0%)	304 (90.7%)
	P-value			0.7480
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	49 (7.6%)	23 (7.4%)	26 (7.8%)
	No	596 (92.4%)	287 (92.6%)	309 (92.2%)
	P-value			0.8700
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		207	105	102
Within intervention comparison (IB vs CD)		254	134	120
Long term follow-up comparison (IC vs CC)		644	322	322
Within trial comparison (IB vs CB)		236	134	102

Table 7.25.3
Summary of Aspirin III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.4 (1.4)	0.3 (1.3)	0.4 (1.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 13	0 , 10	0 , 13
	P-value			0.0338**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	97 (7.5%)	47 (7.3%)	50 (7.8%)
	No	1193 (92.5%)	598 (92.7%)	595 (92.2%)
	P-value			0.4669**
Total Prescriptions				
Pre-FITT vs FITT comparison		461	207	254

Table 7.26.1
Summary of Hypertensives I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.7)	0.0 (0.5)	0.1 (0.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 10	0 , 7	0 , 10
	P-value			0.4662*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.6)	0.0 (0.5)	0.1 (0.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 7	0 , 7	0 , 7
	P-value			0.5484*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (2.0)	0.2 (1.7)	0.3 (2.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 22	0 , 16	0 , 22
	P-value			0.2554*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.7)	0.0 (0.5)	0.1 (0.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 10	0 , 7	0 , 10
	P-value			0.7755*

Table 7.26.2
Summary of Hypertensives II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	6 (0.9%)	2 (0.6%)	4 (1.2%)
	No	639 (99.1%)	308 (99.4%)	331 (98.8%)
	P-value			0.4682
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	8 (1.2%)	3 (1.0%)	5 (1.5%)
	No	637 (98.8%)	307 (99.0%)	330 (98.5%)
	P-value			0.5474
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	16 (2.5%)	10 (3.2%)	6 (1.8%)
	No	629 (97.5%)	300 (96.8%)	329 (98.2%)
	P-value			0.2418
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	7 (1.1%)	3 (1.0%)	4 (1.2%)
	No	638 (98.9%)	307 (99.0%)	331 (98.8%)
	P-value			0.7817
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		44	13	31
Within intervention comparison (IB vs CD)		39	15	24
Long term follow-up comparison (IC vs CC)		165	74	91
Within trial comparison (IB vs CB)		46	15	31

Table 7.26.3
Summary of Hypertensives III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.1 (0.7)	0.1 (0.7)	0.1 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 10	0 , 10	0 , 7
	P-value			0.7656**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	14 (1.1%)	6 (0.9%)	8 (1.2%)
	No	1276 (98.9%)	639 (99.1%)	637 (98.8%)
	P-value			0.1573**
Total Prescriptions				
Pre-FITT vs FITT comparison		83	44	39

Table 7.27.1
Summary of All Hypertensives I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.3 (3.0)	1.0 (2.5)	1.5 (3.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 20	0 , 18
	P-value			0.1886*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.3 (2.9)	1.2 (2.7)	1.5 (3.1)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 19	0 , 21
	P-value			0.2087*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	3.8 (7.6)	3.5 (7.0)	4.1 (8.2)
	Median (IQR)	0 [0 , 1]	0 [0 , 2]	0 [0 , 1]
	Min - Max	0 , 48	0 , 48	0 , 48
	P-value			0.7245*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.3 (3.0)	1.2 (2.7)	1.5 (3.3)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 19	0 , 19	0 , 18
	P-value			0.3509*

Table 7.27.2
Summary of All Hypertensives II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	128 (19.8%)	56 (18.1%)	72 (21.5%)
	No	517 (80.2%)	254 (81.9%)	263 (78.5%)
	P-value			0.2754
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	134 (20.8%)	58 (18.7%)	76 (22.7%)
	No	511 (79.2%)	252 (81.3%)	259 (77.3%)
	P-value			0.2136
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	164 (25.4%)	78 (25.2%)	86 (25.7%)
	No	481 (74.6%)	232 (74.8%)	249 (74.3%)
	P-value			0.8818
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	130 (20.2%)	58 (18.7%)	72 (21.5%)
	No	515 (79.8%)	252 (81.3%)	263 (78.5%)
	P-value			0.3787
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		821	319	502
Within intervention comparison (IB vs CD)		865	365	500
Long term follow-up comparison (IC vs CC)		2461	1097	1364
Within trial comparison (IB vs CB)		867	365	502

Table 7.27.3
Summary of All Hypertensives III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	1.3 (2.9)	1.3 (3.0)	1.3 (2.9)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 20	0 , 21
	P-value			0.3578**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	262 (20.3%)	128 (19.8%)	134 (20.8%)
	No	1028 (79.7%)	517 (80.2%)	511 (79.2%)
	P-value			0.0833**
Total Prescriptions				
Pre-FITT vs FITT comparison		1686	821	865

Table 7.27.4
Summary of All Hypertensives with Baseline BPLetter Issued I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.8 (2.6)	0.5 (2.0)	1.0 (3.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 20	0 , 20	0 , 18
	P-value			0.1232*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.8 (2.5)	0.6 (2.2)	1.0 (2.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 19	0 , 21
	P-value			0.0775*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	2.3 (6.6)	1.9 (5.8)	2.7 (7.2)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 48	0 , 48	0 , 48
	P-value			0.2254*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.8 (2.7)	0.6 (2.2)	1.0 (3.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 19	0 , 19	0 , 18
	P-value			0.1730*

Subscription is counted in those who have been issued BPLetter at baseline

Table 7.27.5
Summary of All Hypertensives with Baseline BPLetter Issued II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	72 (11.2%)	29 (9.4%)	43 (12.8%)
	No	573 (88.8%)	281 (90.6%)	292 (87.2%)
	P-value			0.1607
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	77 (11.9%)	30 (9.7%)	47 (14.0%)
	No	568 (88.1%)	280 (90.3%)	288 (86.0%)
	P-value			0.0885
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	96 (14.9%)	41 (13.2%)	55 (16.4%)
	No	549 (85.1%)	269 (86.8%)	280 (83.6%)
	P-value			0.2551
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	73 (11.3%)	30 (9.7%)	43 (12.8%)
	No	572 (88.7%)	280 (90.3%)	292 (87.2%)
	P-value			0.2059
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		510	170	340
Within intervention comparison (IB vs CD)		523	189	334
Long term follow-up comparison (IC vs CC)		1503	601	902
Within trial comparison (IB vs CB)		529	189	340

Subscription is counted in those who have been issued BPLetter at baseline

Table 7.27.6
Summary of All Hypertensives with Baseline BPLetter Issued III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.8 (2.6)	0.8 (2.6)	0.8 (2.5)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 21	0 , 20	0 , 21
	P-value			0.9183**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	149 (11.6%)	72 (11.2%)	77 (11.9%)
	No	1141 (88.4%)	573 (88.8%)	568 (88.1%)
	P-value			0.0956**
Total Prescriptions				
Pre-FITT vs FITT comparison		1033	510	523

Subscription is counted in those who have been issued BPLetter at baseline

Table 7.27.7
Summary of All Hypertensives without Baseline BPLetter Issued I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.6)	0.5 (1.7)	0.5 (1.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 9
	P-value			0.9834*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.8)	0.6 (1.9)	0.5 (1.7)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 9
	P-value			0.8222*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	1.5 (4.7)	1.6 (4.6)	1.4 (4.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 28	0 , 24	0 , 28
	P-value			0.2944*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.5 (1.8)	0.6 (1.9)	0.5 (1.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 9
	P-value			0.8129*

Subscription is counted in those who have not been issued BPLetter at baseline

Table 7.27.8
Summary of All Hypertensives without Baseline BPLetter Issued II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	56 (8.7%)	27 (8.7%)	29 (8.7%)
	No	589 (91.3%)	283 (91.3%)	306 (91.3%)
	P-value			0.9810
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	57 (8.8%)	28 (9.0%)	29 (8.7%)
	No	588 (91.2%)	282 (91.0%)	306 (91.3%)
	P-value			0.8667
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	68 (10.5%)	37 (11.9%)	31 (9.3%)
	No	577 (89.5%)	273 (88.1%)	304 (90.7%)
	P-value			0.2678
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	57 (8.8%)	28 (9.0%)	29 (8.7%)
	No	588 (91.2%)	282 (91.0%)	306 (91.3%)
	P-value			0.8667
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		311	149	162
Within intervention comparison (IB vs CD)		342	176	166
Long term follow-up comparison (IC vs CC)		958	496	462
Within trial comparison (IB vs CB)		338	176	162

Subscription is counted in those who have not been issued BPLetter at baseline

Table 7.27.9
Summary of All Hypertensives without Baseline BPLetter Issued III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.5 (1.7)	0.5 (1.6)	0.5 (1.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 11
	P-value			0.1499**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	113 (8.8%)	56 (8.7%)	57 (8.8%)
	No	1177 (91.2%)	589 (91.3%)	588 (91.2%)
	P-value			0.5637**
Total Prescriptions				
Pre-FITT vs FITT comparison		653	311	342

Subscription is counted in those who have not been issued BPLetter at baseline

Table 7.28.1
Summary of Hypnotics & Anxyolitics I, Overall and by Group

		All	Intervention	Control
Number of Prescriptions Per Patient				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.8)	0.1 (0.8)	0.1 (0.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 12	0 , 9
	P-value			0.9233*
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.6)	0.1 (0.7)	0.1 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 5
	P-value			0.8414*
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.3 (1.9)	0.2 (1.8)	0.4 (2.0)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 31	0 , 31	0 , 28
	P-value			0.3356*
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Mean (SD)	0.1 (0.7)	0.1 (0.7)	0.1 (0.8)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 11	0 , 11	0 , 9
	P-value			0.6172*

Table 7.28.2
Summary of Hypnotics & Anxyolitics II, Overall and by Group

		All	Intervention	Control
Any Prescriptions				
Prior to intervention comparison (IA vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	36 (5.6%)	17 (5.5%)	19 (5.7%)
	No	609 (94.4%)	293 (94.5%)	316 (94.3%)
	P-value			0.9173
Within intervention comparison (IB vs CD)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	32 (5.0%)	15 (4.8%)	17 (5.1%)
	No	613 (95.0%)	295 (95.2%)	318 (94.9%)
	P-value			0.8904
Long term follow-up comparison (IC vs CC)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	63 (9.8%)	27 (8.7%)	36 (10.7%)
	No	582 (90.2%)	283 (91.3%)	299 (89.3%)
	P-value			0.3840
Within trial comparison (IB vs CB)	n (missing)	645 (0)	310 (0)	335 (0)
	Yes	34 (5.3%)	15 (4.8%)	19 (5.7%)
	No	611 (94.7%)	295 (95.2%)	316 (94.3%)
	P-value			0.6362
Total Prescriptions				
Prior to intervention comparison (IA vs CB)		81	37	44
Within intervention comparison (IB vs CD)		64	26	38
Long term follow-up comparison (IC vs CC)		203	75	128
Within trial comparison (IB vs CB)		70	26	44

Table 7.28.3
Summary of Hypnotics & Anxyolitics III, Overall and by Group

		All	Pre-FITT	FITT
Number of Prescriptions Per Patient				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Mean (SD)	0.1 (0.7)	0.1 (0.8)	0.1 (0.6)
	Median (IQR)	0 [0 , 0]	0 [0 , 0]	0 [0 , 0]
	Min - Max	0 , 12	0 , 12	0 , 11
	P-value			0.4131**
Any Prescriptions				
Pre-FITT vs FITT comparison	n (missing)	1290 (0)	645 (0)	645 (0)
	Yes	68 (5.3%)	36 (5.6%)	32 (5.0%)
	No	1222 (94.7%)	609 (94.4%)	613 (95.0%)
	P-value			0.5553**
Total Prescriptions				
Pre-FITT vs FITT comparison		145	81	64

Table 8.1
Relationship Between Any Hospitalisations During Time Period B and C and the Following:

	Odds Ratio (95% CI)	P-Value
Baseline Weight (kg)	0.9984 (0.9886, 1.0084)	0.753
Change in Weight from Baseline to 3.5 Years (kg)	1.0113 (0.9882, 1.0348)	0.340
Number of Longstanding Illnesses at 3.5 Years	1.6151 (1.3175, 1.9798)	<0.001

Each variable is analyzed separately. Models also adjusted for club as random effect and FFIT RCT group as fixed effect.

Table 8.2.1
Relationship Between Any Prescriptions During Time Period B and C and the Following:

	Odds Ratio (95% CI)	P-Value
Baseline Weight (kg)	1.0040 (0.9854, 1.0229)	0.677
Change in Weight from Baseline to 3.5 Years (kg)	1.0088 (0.9700, 1.0491)	0.661
Number of Longstanding Illnesses at 3.5 Years	1.9323 (1.1170, 3.3428)	0.019

Each variable is analyzed separately. Models also adjusted for club as random effect and FFIT RCT group as fixed effect.

Table 8.2.2
Relationship Between Number of Prescriptions During Time Period B and C and the Following:

	Incident Rate Ratio (95% CI)	P-Value
Baseline Weight (kg)	1.0034 (0.9979, 1.0089)	0.225
Change in Weight from Baseline to 3.5 Years (kg)	1.0056 (0.9930, 1.0183)	0.387
Number of Longstanding Illnesses at 3.5 Years	1.7076 (1.5138, 1.9262)	<0.001

Each variable is analyzed separately. Models also adjusted for club as random effect and FFIT RCT group as fixed effect.