



Best evidence for management of peritoneal metastases

Cytoreductive surgery (CRS) with hyperthermic intraoperative peritoneal chemotherapy (HIPEC) versus standard of care (SoC) in people with peritoneal metastases from colorectal, ovarian or gastric origin:

Systematic review and individual participant data meta-analyses (IPD) of effectiveness and cost-effectiveness

Protocol

Conducted by the
EvidencE Review of PEritoneal Tumours (EVERPET) Working Group

February 2019 (Version 1.0)

This project was funded by NIHR HTA programme (HTA - Project: 17/135/02)

EVERPET-IPD research team

Project Management Group

Kurinchi Gurusamy

Division of Surgery and Interventional Science

University College London

Email: k.gurusamy@ucl.ac.uk

Tel: +44 (0)20 7794 0500 Ext: 33943

Claire Vale

MRC Clinical Trials Unit at UCL

Email: claire.vale@ucl.ac.uk

Tel: +44 (0)20 7670 4723

Rajesh Bhanot (patient representative)

Email: will be disclosed on a need to know basis

Research associate (to be appointed)

Research Steering Group

Kurinchi Gurusamy

Claire Vale

Brian Davidson

University College London

Email: b.davidson@ucl.ac.uk

Sarah O'Dwyer

The Christie NHS Foundation Trust

Email: Sarah.O'Dwyer@christie.nhs.uk

Omer Aziz

The Christie NHS Foundation Trust

Email: Omer.Aziz@christie.nhs.uk

Mark Saunders

The Christie NHS Foundation Trust

Email: Mark.Saunders@christie.nhs.uk

Tim Mould

University College London NHS Foundation Trust

Email: t.mould@nhs.net

Muntzer Mughal

University College London NHS Foundation Trust

Email: muntzer.mughal@nhs.net

Elena Pizzo

Centre of Applied Health Research

University College London

Email: e.pizzo@ucl.ac.uk

Two additional patient representatives

Meta-analysis Group

Kurinchi Gurusamy

Claire Vale

Research associate (to be appointed)

Collaborating Trialists

To be completed once additional confirmation is obtained

Contents

EVERPET-IPD research team	2
Project Management Group	2
Research Steering Group.....	2
Meta-analysis Group	3
Collaborating Trialists.....	3
Background and rationale	5
What is the problem being addressed?.....	5
Treatment of peritoneal metastases.....	5
Why is this research important to patients and health and care services?	5
Review of existing evidence	5
Justification for IPD	6
Aims and objectives	6
Primary objectives	6
Secondary objectives	6
General Methods	7
Eligibility criteria.....	7
Search strategy	7
Electronic searches	7
Other resources	8
Data collection.....	8
Selection of studies.....	8
Data collection	8
Data checking and management	9
Assessment of risk of bias in included studies.....	9
Meta-analysis of clinical effectiveness	9
Measures of treatment effect and data synthesis.....	9
Dealing with missing data	10
Assessment and investigation of heterogeneity	10
Sensitivity analysis.....	10
Cost-effectiveness analysis.....	10
Model	10
Measuring cost-effectiveness	11
Project management.....	12
Dissemination and reporting plan	13
Project / research timetable.....	13
Appendix 1: Search Strategies	14
Medline.....	14
Embase	14
Cochrane	15
Science Citation Index.....	16
WHO trials register	16
ClinicalTrials.gov	16
Cost-Effectiveness Analysis (CEA) Registry	16
References	17

Background and rationale

What is the problem being addressed?

Approximately 7 million people worldwide and 160,000 people in UK develop colorectal, ovarian, or gastric cancer each year [1], of whom 8% to 50% develop peritoneal metastases. The peritoneum is one of the commonest sites of metastases from these cancers [2-8], and is often the only site of metastases [7-9]. In general, people with peritoneal metastases have poorer prognosis than those with other metastases (liver or lung) [10], with median reported survival ranging from 6 to 24 months depending on from the primary cancers and treatment received [11-13].

Treatment of peritoneal metastases

The current standard of care of people with peritoneal metastases from these cancers is systemic chemotherapy alone or in combination with either cytoreductive surgery (CRS) or palliative surgery [7, 8, 12-15]. CRS + hyperthermic intraoperative peritoneal chemotherapy (HIPEC) is an alternative treatment for these patients. The main principle of CRS+HIPEC is to remove all visible (macroscopic) peritoneal metastases followed by HIPEC to treat any remaining microscopic peritoneal metastases [16]. HIPEC involves peritoneal circulation of chemotherapy drugs (usually mitomycin C, 5-Fluorouracil, and oxaliplatin, or cisplatin) [17] heated to temperatures of 42° C, at which the chemotherapy drugs are potentiated [18]. Until only a decade ago, less than 5% of patients with peritoneal metastases underwent CRS+HIPEC, however this has progressively increased to about 10% of patients by 2012 [8, 9, 14]. CRS+HIPEC has been commissioned by NHS England for patients with peritoneal metastases from appendiceal tumours and colorectal adenocarcinoma.

Why is this research important to patients and health and care services?

Although CRS+HIPEC has the potential to improve the survival and health-related quality of life (HRQoL) in people with peritoneal metastases [14, 19, 20], there have been concerns raised about its safety. Reports have shown a 30-day mortality after CRS+HIPEC of 1-3% [6], and a major complication rate of 32% [6, 21], albeit that local audit data from high volume centres suggest that mortality and morbidity rates are somewhat less than in these reports (local audit data). The average costs of CRS+HIPEC per patient varies from about 20,000 USD to 80,000 USD [22-28]. Because of these reasons, this research is important to address the significant uncertainty about the benefits of an intervention that carries significant risk of harm to patients and costs to the NHS.

Review of existing evidence

There have been several overviews, systematic reviews, and health technology assessments (HTA) investigating this area. Sixteen systematic reviews of comparative studies have been undertaken, comparing CRS+HIPEC to other treatment modalities in peritoneal metastases from colorectal, ovarian, or gastric cancer [6, 17, 20, 29-41]. Ten of these included at least one RCT, but the conclusions were largely based on non-randomised studies [6, 17, 20, 29, 31-33, 35, 39, 41]. Although most of these systematic reviews concluded that CRS+HIPEC can improve survival in people with peritoneal metastases, all had limitations and deficiencies. Firstly, all are at high risk of bias according to ROBIS (risk of bias in systematic reviews) tool [42] with concern about bias across all domains. Secondly, the systematic reviews included only a single RCT [13] and/or based their evidence predominantly on non-randomised studies, without any adjustment for baseline differences in

disease-related or patient-related prognostic characteristics [6, 17, 20, 29, 31-33, 35, 39, 41]. Finally, meta-analyses could only include a small proportion of the results from the studies because of the way these results had been reported (e.g. proportion survived versus median survival) [17, 20, 29, 35, 37]. Therefore, there is still considerable uncertainty about the benefits of CRS+HIPEC and which patient groups will benefit from it.

There have been two formal HTAs on this issue [26, 43]. The HTA reviewing patients with peritoneal disease from colorectal cancer concluded that there was moderate quality evidence that CRS+HIPEC prolonged survival based on a single RCT, but the costs were high [26]. The HTA on ovarian cancer (which did not include any RCTs) concluded there was no clear benefit of CRS+HIPEC for ovarian peritoneal metastases [43].

Justification for IPD

Through the collection and reanalysis of IPD from all relevant randomised controlled trials, we aim to overcome the limitations of the existing evidence and provide the highest quality evidence synthesis of the benefits and harms of CRS + HIPEC in patients with peritoneal metastases to inform clinical practice and future research. Importantly, the main advantages of using IPD over aggregate data in this setting are the following.

1. Overcome lack of reporting of key survival outcomes: The key survival outcomes have not been reported in a format that can be meta-analysed. This can be overcome with IPD.
2. Harmonise definitions of performance indicators and outcome: Use of IPD can ensure that the definitions of the prognostic and confounding factors, and outcomes are harmonised.
3. Improve the quality of the analysis: IPD is commonly reported to improve the quality of analyses [44, 45].
4. Investigate whether any patient-related or disease-related characteristics impact on the treatment effect at the individual level

Aims and objectives

The overarching aim of this project is to answer the following research question:

Does CRS+HIPEC improve survival and/or quality of life compared to SoC in people with peritoneal metastases (from colorectal, ovarian, or gastric cancers) who can withstand major surgery and is it cost-effective in the NHS setting?

Primary objectives

To compare the relative benefits and harms of CRS+HIPEC versus SoC in people with peritoneal metastases from colorectal, ovarian, or gastric cancers eligible to undergo CRS+HIPEC by a systematic review and IPD meta-analysis.

Secondary objectives

To compare the cost-effectiveness of CRS+HIPEC versus SoC from an NHS and PSS (personal social services) perspective using a model-based cost-utility analysis.

General Methods

Eligibility criteria

Type of studies

All RCTs regardless of the publication status, year of publication, and language of publication will be included.

Setting

Secondary or tertiary care with expertise to perform CRS+HIPEC

Type of participants

Inclusion criteria

People with synchronous or metachronous peritoneal metastases from colorectal cancer, ovarian cancer, or gastric cancer, eligible to undergo CRS+HIPEC regardless of the involvement of other organs and whether the primary cancer was resected completely (i.e. R0 resection). We will also include people with appendiceal adenocarcinomas under colorectal cancer as they behave in a similar way to colorectal adenocarcinomas.

Exclusion criteria

Studies on pseudomyxoma peritonei (PMP) will be excluded.

Intervention

CRS+HIPEC

Control

Standard of care (SoC)

Outcomes

Primary outcomes

1. Overall survival, defined as time from randomisation until death by any cause.
2. HRQoL using any validated measure
3. Serious adverse events or Clavien-Dindo grade III or above [46, 47]

Secondary outcomes

4. Time to disease progression: defined as time from diagnosis to death in people who died of treatment or disease-related causes, time from diagnosis to recurrence in people in whom complete CRS was achieved, and time from diagnosis to disease progression as defined by RECIST (Response Evaluation Criteria in Solid Tumors) criteria of 20% increase in size of the tumour or appearance of new lesions [48], or similar criteria used by authors
5. Non-serious adverse events or Clavien-Dindo grade I or II [46, 47]

Search strategy

Electronic searches

We will search MEDLINE, EMBASE, Cochrane library, and the Science Citation Index for published trials as well as ClinicalTrials.gov, and WHO ICTRP trial registers for ongoing or unreported studies. The search strategies, which combine the Cochrane sensitivity maximising RCT filter [49] with a

combination of subject headings and free text terms relating to the interventions and diseases of interest, are provided in Appendix 1. Searches will be updated periodically until October 2019.

Other resources

We will also search the references of all identified studies for additional studies eligible for inclusion. We will also contact the American Society of Peritoneal Surface Malignancies, the Canadian HIPEC Collaborative Group (CHICG), The Peritoneal Surface Oncology Group International (PSOGI), and the study authors who agree to participate in this project for further studies.

Data collection

Selection of studies

Two review authors will independently screen the titles and abstracts of all records retrieved and make the final selection based on full text (after translation if required, i.e. there will be no language restrictions). We will document the process to enable completion of the PRISMA flow-chart. We will resolve discrepancies through discussion and arbitration.

Data collection

At the study level, we will record the contact details of the study author and the study contact, information required to assess the risk of bias, details of the treatment centres (name and the average number of CRS+HIPEC performed per year).

At the participant level, we will collect the following details:

1. Centre at which treated
2. Patient demographics: age, gender, comorbidities, performance index
3. Cancer details (including severity)
4. Intervention details
5. Control details
6. Follow-up details
7. Outcome data
8. Resource utilisation data
 - a. Operating time
 - b. Quantity of blood and blood products transfused
 - c. Length of hospital stay (including readmissions)
 - d. Length of intensive care unit stay
 - e. Chemotherapy regimen used in HIPEC and in control group if applicable
 - f. Proportion in whom surgery was performed and the nature of surgery in the control group
 - g. Additional surgery and other palliative treatments

These data will be sought for all patients randomised into each trial. Up to date follow-up will be requested in order to report on longer-term outcomes: the existing ethical approval for the studies usually cover collection of data.

The proposed data format and coding conventions for these data will be developed as part of the project to obtain the EVERPET-IPD data dictionary. Transfer guide will be developed as part of the project. Although the aim of the conventions is to facilitate data transfer, they are not essential. Data will be accepted in the format most convenient for the individual trial investigator or data centre, however, **all personal identifiers (e.g. names) are to be removed before sharing**. Data should be transferred by encrypted email or source ftp site. Further details are included in the data transfer guide.

Data checking and management

Once trial investigators have agreed to provide the IPD, they will be asked to sign a data transfer agreement that covers the transfer, use and storage of that data. By signing up to the agreement, investigators also declare that they have complied with all laws and regulations relating to the conduct of their studies and the collection of data as part of those studies.

On receipt, data will be cleaned and checked for accuracy, consistency and validity. This will include checks for missing data, randomisation integrity, follow-up and censoring. We will query any anomalies with the study contact to ensure that the data are represented accurately, and send a summary of the final dataset from each trial to the study contacts for verification.

Once checked and verified, we will store the trial data securely. Access to the data will be restricted to the Project Management Group, who are all trained in data protection and personal data confidentiality and who will act as custodians of the data under the terms of the data transfer agreement, which will be developed as part of this project. In line with that agreement, data will be deposited in the EVERPET-IPD repository.

Assessment of risk of bias in included studies

We will use the Cochrane risk of bias tool to assess the risk of bias in RCTs [49]. If the RoB 2.0 [50] is validated, we will use the RoB 2.0 for assessment of risk of bias.

Meta-analysis of clinical effectiveness

Measures of treatment effect and data synthesis

We will use risk ratio for binary outcomes (proportion of people with serious and non-serious adverse events), mean difference (if same scales are reported in the studies) or standardised mean difference (if different scales are reported in the studies) for continuous outcomes (HRQoL), rate ratios for count outcomes (number of serious and non-serious adverse events), and hazard ratio for time-to-event outcomes (overall all-cause mortality and time to progression) with their respective 95% confidence intervals.

We will perform a two-step IPD, i.e. calculate the adjusted effect estimate from each included study and then perform a random-effects model meta-analysis using DerSimonian and Laird method [51] for binary outcomes and inverse variance method for other types of outcomes. The reason for choosing the two-step IPD over one-step IPD is the way the confounding factors are reported in the studies, for example, comorbidities can be reported as different types of performance indices and the extent of peritoneal disease can be reported in different ways [52, 53]. However, if we agree on

an approximation to convert different performance indices into a single measure and convert the different measures of extent of peritoneal involvement into a single measure, we will perform a single-step meta-analysis to check the robustness of the two-step meta-analysis. We will test our assumptions in approximations (of the different performance indices into a single measure and different measures of extent of peritoneal involvement into a single measure) by sensitivity analyses. We will use multilevel modelling to take the clustering of data in the studies into account for the one-step IPD meta-analysis, as the unit of analysis will be the individual participant.

Dealing with missing data

We will perform an intention-to-treat analysis whenever possible [54]. If data on the classification of the treatment as intervention or control is missing, and cannot be ascertained through discussion with trialists, we will exclude such participants. If outcome data are missing, we will use multiple imputation method if the data are likely to be missing at random or best-case and worst-case scenarios analysis if it is felt that the outcome data are not missing at random.

Assessment and investigation of heterogeneity

We will assess the clinical and methodological heterogeneity by carefully examining the characteristics and design of included trials. Clinical heterogeneity could be due to the type of participants included in the studies (performance index, stage of cancer, extent of peritoneal involvement, other organ involvement), different interventions (complete CRS or not, chemotherapy agents used), different controls (chemotherapy alone or CRS or both), whether complete CRS was achieved (if the control group was CRS), or different follow-up methods (routine imaging versus clinical examination). Different study designs and risk of bias may contribute to methodological heterogeneity. We will calculate and report the between-trial standard deviation and I^2 as measures of heterogeneity.

If we identify substantial clinical, methodological, or statistical heterogeneity, we will explore and address it in subgroup analyses and/or metaregression using participant level covariates on the sources of clinical heterogeneity mentioned above except for routine imaging which will be a trial-level covariate. All sources of methodological heterogeneity will be trial-level covariates.

Sensitivity analysis

We will perform the following sensitivity analyses to assess the impact of:

- data not missing at random
- non-participation in the IPD
- methods (two-step versus single-step) and model (fixed-effect versus random-effects model) used for meta-analysis
- risk of bias.

Cost-effectiveness analysis

Model

We will perform a model-based cost-utility analysis estimating mean costs and quality-adjusted life years (QALYs) per patient. We will compare CRS+HIPEC versus SoC in each of the three cancers by three separate cost-effectiveness analyses. The time horizon will be life-time time horizon. We will

calculate the costs from the NHS and personal social services (PSS) perspective. We will discount the costs and utilities at the rate of 3.5% per annum [55].

We will create a decision tree model (one for each cancer) along the lines of the model that we used to compare two types of surgeries in pancreatic cancer [56]. Briefly, a patient with peritoneal metastases from one of the three cancers (colorectal cancer, ovarian cancer, or gastric cancer) and eligible for CRS+HIPEC can either undergo CRS+HIPEC or SoC. A proportion of patients undergoing CRS+HIPEC will have complete CRS (i.e. all macroscopic tumour is removed). A proportion of patients in whom CRS+HIPEC will develop complications (whether complete CRS was achieved or not), a proportion of whom may die within 90 days. Those who are alive at 90 days may die between 90 days and 1 year; a proportion of people who are alive at 1 year may die between 1 year and 2 years; and so on. The decision tree pathways in the people who had SoC will be identical: some will have complete CRS, some will have complications, some will die within 90 days, and some will die after 90 days.

Most of the information required for populating the decision tree (including resource utilisation data) will be obtained from the systematic review and IPD meta-analysis. For information not available from the systematic review and IPD meta-analysis, we will perform literature searches of NHS Economic Evaluation Database (NHS EED), the Health Economic Evaluations Database (HEED), MEDLINE, and EMBASE (for MEDLINE and EMBASE, we will combine the search strategy from Table 5 with sensitivity maximising 'economics' filter developed as a part of [The Hedges Project of the Health Information Research Unit of McMaster University](#)). We will also review the [Cost-Effectiveness Analysis Registry \(CEA\) at Tufts University](#) for information on quality of life. Currently, there is no HRG (Healthcare Resource Group) code available for CRS+HIPEC and SoC (which will vary according to the nature of the treatment). We will obtain resource utilisation data as part of the systematic review and IPD (please see above) and convert these to costs on the basis of [NHS National Tariff](#), [NHS National Schedule of Reference costs](#), British National Formulary, and/or local estimates as required.

We will assume that the people who die in each period will do so at a constant rate during the period and check whether this assumption is true using the IPD. If this assumption is not true, then we will use more complex models to mirror the survival curves based on the IPD. When no data are available from the IPD or published sources, a range of values will be used in the model. We will tabulate the inputs used in the decision tree model and the source of these inputs in the project report.

Measuring cost-effectiveness

We will measure cost-effectiveness using net monetary benefits (NMBs). For each treatment, we will calculate the NMB as the mean QALYs per patient accruing to that treatment multiplied by decision-makers' maximum willingness to pay for a QALY (also referred to as the cost-effectiveness threshold), minus the mean cost per patient for the treatment. In the UK, the lower and upper limit of the maximum willingness to pay for a QALY are £20,000 and £30,000 respectively [55]. NMBs will be calculated using the base case parameter values to obtain the deterministic results, which do not depend on chance. The option with the highest NMB represents best value for money. The NMB for CRS+HIPEC minus the NMB for SoC is the incremental NMB. If the incremental NMB is positive then

CRS+HIPEC represents better value for money; if it is negative, the SoC represents better value for money.

A probabilistic sensitivity analysis (PSA) will also be undertaken [55]. The PSA involves Monte Carlo simulation and takes variability of all selected inputs into account simultaneously. Distributions will be assigned to parameters to reflect the uncertainty with each parameter value. A random value from the corresponding distribution for each parameter will be selected (by the computer). This will generate an estimate of the mean cost and mean QALYs and the NMB associated with each treatment. This will be repeated 5000 times and the results for each simulation will be noted. The mean costs, QALYs and NMB for each treatment will be calculated from the 5000 simulations; these are probabilistic results because they depend on chance. The NMB will also be calculated for each of the 5000 simulations and the proportion of times each treatment had the highest NMB will be calculated for a range of values for the maximum willingness to pay for a QALY. These will be summarised graphically using cost-effectiveness acceptability curves. We will derive the 95% confidence intervals around the base case values using the 2.5 and 97.5 percentiles calculated from the PSA.

For the deterministic univariate sensitivity analysis, each variable in the cost-effectiveness model will be varied one at a time. The results of the sensitivity analysis will be represented in the tornado diagram which reflects the variation in the NMB within the range of the lowest and highest value used for a parameter with all else equal. If the variation in the NMB includes 0, then there is uncertainty in the cost-effectiveness due to the variation of the parameter.

We will also perform various subgroup analyses guided by the results of the systematic reviews and IPD meta-analyses, but will include subgroup analysis of different types of control (i.e. CRS alone or systemic chemotherapy alone or both) as a minimum. We will also perform a sensitivity analysis using information from 'real-life' prospective data from Christie NHS foundation trust (and from other NHS specialist centres if such information is available).

Project management

The overall project will be managed by the principal applicant, Prof Kurinchi Gurusamy (KG). The day-to-day management, research co-ordination, data collection, and data cleaning will be performed by a research associate (RA) appointed as part of this grant. Methodological support will be provided by KG and Dr Claire Vale (CV). The health economics supervision will be provided by Prof Steve Morris. The clinical co-applicants will help with recruitment of trialists through their professional connections with those trialists. The Research Management Group (RMG) comprising of the KG, CV, a patient representative, and the newly appointed RA will be established to ensure that the project is progressing as planned, and to take any remedial actions to ensure that the project milestones are met. The RMG will meet either via teleconferences or with face to face meetings monthly. The RMG will report on progress every 6 months to the research steering group (RSG) made up of all co-applicants and key collaborators. The RMG and RSG will ensure smooth running of the project and adherence to key milestones.

Dissemination and reporting plan

The authorship of the systematic review manuscript will comprise the Project Management Group, International Advisory Group, representatives from the included trials and patient representatives. Author names will be listed “for the EVERPET-IPD Working Group”. We aim to present the findings at appropriate international meetings and publish the review, irrespective of the findings, in a peer-reviewed journal. A manuscript will be drafted, circulated to the Working Group for comment prior to being submitted for publication.

Project / research timetable

The timetable for key activities of the project are as follows.

Proposed Time Table	
Spring 2019	Finalise and register protocol
Summer / Autumn 2019	Appoint Research Associate Finalise and establish data use agreements
Winter 2019	Set up secure database Initiate collaboration and data collection
Spring – Autumn 2020	Collate, check and verify incoming data for the first cancer type Analyse individual trials
Winter 2020	Complete meta-analysis for the first cancer type Report preliminary results to Collaborative Group
Spring 2021	Publish and present results for the first cancer type Collate, check and verify incoming data for the remaining cancer types
Summer-Winter 2021	Analyse individual trials Complete meta-analysis for the remaining cancer types Report preliminary results to Collaborative Group
Spring 2022	Publish and present results for the remaining cancer types Complete and report results for the cost effectiveness analysis

Appendix 1: Search Strategies

Medline

1. Hyperthermia, Induced/
2. ((hyperthermic or heated) adj3 (intraperitoneal or intra-peritoneal) adj3 (chemotherapy or chemotherapies)).ti,ab.
3. (intraperitoneal adj3 chemohyperthermia).ti,ab.
4. (HIPEC or IPHC or HIIC).ti,ab.
5. 1 or 2 or 3 or 4
6. Cytoreduction Surgical Procedures/
7. ((cytoreductive or cytoreduction or debulking) adj3 (surgery or surgeries or surgical or procedure or procedures)).ti,ab.
8. 6 or 7
9. 5 or 8
10. exp Colorectal Neoplasms/
11. exp Ovarian Neoplasms/
12. Stomach Neoplasms/
13. ((colorectal or bowel or colon or colonic or rectum or rectal or ovary or ovaries or ovarian or gastric or stomach) adj3 (cancer or cancers or carcinoma or carcinomas or tumour or tumours or tumor or tumors or neoplasm or neoplasms)).ti,ab.
14. 10 or 11 or 12 or 13
15. 9 and 14
16. randomized controlled trial.pt.
17. controlled clinical trial.pt.
18. randomized.ab.
19. placebo.ab.
20. drug therapy.fs.
21. randomly.ab.
22. trial.ab.
23. groups.ab.
24. 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23
25. exp animals/ not humans.sh.
26. 24 not 25
27. 15 and 26
28. (cost: or cost benefit analys: or health care costs).mp.
29. 15 and 28
30. 27 or 29

Embase

1. hyperthermic intraperitoneal chemotherapy/
2. ((hyperthermic or heated) adj3 (intraperitoneal or intra-peritoneal) adj3 (chemotherapy or chemotherapies)).ti,ab.
3. (intraperitoneal adj3 chemohyperthermia).ti,ab.
4. (HIPEC or IPHC or HIIC).ti,ab.

5. 1 or 2 or 3 or 4
6. cytoreductive surgery/
7. ((cytoreductive or cytoreduction or debulking) adj3 (surgery or surgeries or surgical or procedure or procedures)).ti,ab.
8. 6 or 7
9. 5 or 8
10. exp colon cancer/
11. exp rectum cancer/
12. exp ovary cancer/
13. exp stomach cancer/
14. ((colorectal or bowel or colon or colonic or rectum or rectal or ovary or ovaries or ovarian or gastric or stomach) adj3 (cancer or cancers or carcinoma or carcinomas or tumour or tumours or tumor or tumors or neoplasm or neoplasms)).ti,ab.
15. 10 or 11 or 12 or 13 or 14
16. 9 and 15
17. exp crossover-procedure/ or exp double-blind procedure/ or exp randomized controlled trial/ or single-blind procedure/
18. ((((((random* or factorial* or crossover* or cross over* or cross-over* or placebo* or double*) adj blind*) or single*) adj blind*) or assign* or allocat* or volunteer*).af.
19. 17 or 18
20. 16 and 19
21. (cost or costs).tw.
22. 16 and 21
23. 20 or 22

Cochrane

- #1 MeSH descriptor: [Hyperthermia, Induced] this term only
- #2 ((hyperthermic or heated) near/3 (intraperitoneal or intra-peritoneal) near/3 (chemotherapy or chemotherapies))
- #3 (intraperitoneal near/3 chemohyperthermia)
- #4 (HIPEC or IPHC or HIIC)
- #5 #1 or #2 or #3 or #4
- #6 MeSH descriptor: [Cytoreduction Surgical Procedures] this term only
- #7 ((cytoreductive or cytoreduction or debulking) near/3 (surgery or surgeries or surgical or procedure or procedures))
- #8 #6 or #7
- #9 #5 or #8
- #10 MeSH descriptor: [Colorectal Neoplasms] explode all trees
- #11 MeSH descriptor: [Ovarian Neoplasms] explode all trees
- #12 MeSH descriptor: [Stomach Neoplasms] this term only
- #13 ((colorectal or bowel or colon or colonic or rectum or rectal or ovary or ovaries or ovarian or gastric or stomach) near/3 (cancer or cancers or carcinoma or carcinomas or tumour or tumours or tumor or tumors or neoplasm or neoplasms))
- #14 #10 or #11 or #12 or #13

#15 #9 and #14

Science Citation Index

1 TS=((hyperthermic or heated) near/3 (intraperitoneal or intra-peritoneal) near/3 (chemotherapy or chemotherapies))

2 TS=(intraperitoneal near/3 chemohyperthermia)

3 TS=(HIPEC or IPHC or HIIC)

4 #3 OR #2 OR #1

5 TS=((cytoreductive or cytoreduction or debulking) near/3 (surgery or surger-ies or surgical or procedure or procedures))

6 #5 or #4

7 TS=((colorectal or bowel or colon or colonic or rectum or rectal or ovary or ovaries or ovarian or gastric or stomach) near/3 (cancer or cancers or carci-noma or carcinomas or tumour or tumours or tumor or tumors or neoplasm or neoplasms))

#8 TS=(random* or placebo* or blind* or meta-analysis or cost or costs)

#9 #8 AND #7 AND #6

WHO trials register

Condition: colorectal OR bowel OR colon OR colonic OR rectum OR rectal OR ovary OR ovaries OR ovarian OR gastric OR stomach

Intervention: HIPEC OR hyperthermic intraperitoneal chemotherapy OR IPHC OR intraperitoneal chemohyperthermia OR HIIC OR heated intraoperative intraperitoneal chemotherapy OR cytoreductive surgery OR CRS

ClinicalTrials.gov

Condition: colorectal OR bowel OR colon OR colonic OR rectum OR rectal OR ovary OR ovaries OR ovarian OR gastric OR stomach

Study Type: Interventional Studies (Clinical Trials)

Intervention/treatment: HIPEC OR hyperthermic intraperitoneal chemotherapy OR IPHC OR intraperitoneal chemohyperthermia OR HIIC OR heated intraoperative intraperitoneal chemotherapy OR cytoreductive surgery OR CRS

Interventional studies, phase 2,3,4

Interventional Studies | colorectal OR bowel OR colon OR colonic OR rectum OR rectal OR ovary OR ovaries OR ovarian OR gastric OR stomach | HIPEC OR hyperthermic intraperitoneal chemotherapy OR IPHC OR intraperitoneal chemohyperthermia OR HIIC OR heated intraoperative intraperitoneal chemotherapy OR cytoreductive surgery OR CRS | Phase 2, 3, 4

Cost-Effectiveness Analysis (CEA) Registry

The following terms were searched:

Hyperthermic

Cytoreduction

Cytoreductive

References

1. International Agency for Research on Cancer (World Health Organization). Estimated number of incident cases, both sexes, all cancers excluding non-melanoma skin cancer, worldwide in 2012. https://gco.iarc.fr/today/online-analysis-table?mode=cancer&mode_population=continents&population=900&sex=0&cancer=29&type=0&statistic=0&prevalence=0&color_palette=default (accessed on 3rd December 2017) 2017.
2. Lemmens VE, Klaver YL, Verwaal VJ et al. Predictors and survival of synchronous peritoneal carcinomatosis of colorectal origin: A population-based study. *International Journal of Cancer* 2011; 128: 2717-2725.
3. Riihimäki M, Hemminki A, Sundquist J, Hemminki K. Patterns of metastasis in colon and rectal cancer. *Scientific Reports* 2016; 6: 29765.
4. Segelman J, Granath F, Holm T et al. Incidence, prevalence and risk factors for peritoneal carcinomatosis from colorectal cancer. *British Journal of Surgery* 2012; 99: 699-705.
5. van Gestel YRBM, Thomassen I, Lemmens VEPP et al. Metachronous peritoneal carcinomatosis after curative treatment of colorectal cancer. *European Journal of Surgical Oncology (EJSO)* 2014; 40: 963-969.
6. Baratti D, Kusamura S, Pietrantonio F et al. Progress in treatments for colorectal cancer peritoneal metastases during the years 2010-2015. A systematic review. *Critical Reviews in Oncology/Hematology* 2016; 100: 209-222.
7. Thomassen I, van Gestel YR, van Ramshorst B et al. Peritoneal carcinomatosis of gastric origin: A population-based study on incidence, survival and risk factors. *International Journal of Cancer* 2014; 134: 622-628.
8. Quere P, Facy O, Manfredi S et al. Epidemiology, Management, and Survival of Peritoneal Carcinomatosis from Colorectal Cancer: A Population-Based Study. *Dis Colon Rectum* 2015; 58: 743-752.
9. Klaver YL, Lemmens VE, Creemers GJ et al. Population-based survival of patients with peritoneal carcinomatosis from colorectal origin in the era of increasing use of palliative chemotherapy. *Ann Oncol* 2011; 22: 2250-2256.
10. Hwang M, Jayakrishnan TT, Green DE et al. Systematic review of outcomes of patients undergoing resection for colorectal liver metastases in the setting of extra hepatic disease. *Eur J Cancer* 2014; 50: 1747-1757.
11. Spiliotis J, Halkia E, Lianos E et al. Cytoreductive surgery and HIPEC in recurrent epithelial ovarian cancer: a prospective randomized phase III study. *Annals of Surgical Oncology* 2015; 22: 1570-1575.
12. Yang XJ, Huang CQ, Suo T et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy improves survival of patients with peritoneal carcinomatosis from gastric cancer: Final results of a phase III randomized clinical trial. *Annals of Surgical Oncology* 2011; 18: 1575-1581.
13. Verwaal VJ, Van Ruth S, De Bree E et al. Randomized trial of cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy and palliative surgery in patients with peritoneal carcinomatosis of colorectal cancer. *Journal of Clinical Oncology* 2003; 21: 3737-3743.
14. Razenberg LG, van Gestel YR, Creemers GJ et al. Trends in cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for the treatment of synchronous peritoneal carcinomatosis of colorectal origin in the Netherlands. *Eur J Surg Oncol* 2015; 41: 466-471.
15. Verwaal VJ, Bruin S, Boot H et al. 8-year follow-up of randomized trial: cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy in patients with peritoneal carcinomatosis of colorectal cancer. *Ann Surg Oncol* 2008; 15: 2426-2432.
16. Elias D, Goere D, Dumont F et al. Role of hyperthermic intraoperative peritoneal chemotherapy in the management of peritoneal metastases. *Eur J Cancer* 2014; 50: 332-340.
17. Desiderio J, Chao J, Melstrom L et al. The 30-year experience-A meta-analysis of randomised and high-quality non-randomised studies of hyperthermic intraperitoneal chemotherapy in the treatment of gastric cancer. *European Journal of Cancer* 2017; 79: 1-14.

18. Schaaf L, van der Kuip H, Zopf W et al. A Temperature of 40 degrees C Appears to be a Critical Threshold for Potentiating Cytotoxic Chemotherapy In Vitro and in Peritoneal Carcinomatosis Patients Undergoing HIPEC. *Annals of Surgical Oncology* 2015; 22: S758-S765.
19. Seretis C, Youssef H. Quality of life after cytoreductive surgery and intraoperative hyperthermic intraperitoneal chemotherapy for peritoneal surface malignancies: a systematic review. *Eur J Surg Oncol* 2014; 40: 1605-1613.
20. Huang CQ, Min Y, Wang SY et al. Cytoreductive surgery plus hyperthermic intraperitoneal chemotherapy improves survival for peritoneal carcinomatosis from colorectal cancer: A systematic review and meta-analysis of current evidence. *Oncotarget* 2017; 8: 55657-55683.
21. Cavaliere D, Cirocchi R, Coccolini F et al. 1st Evidence-based Italian consensus conference on cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinosis from ovarian cancer. *Tumori* 2017; 103: 525-536.
22. Baratti D, Scivales A, Balestra MR et al. Cost analysis of the combined procedure of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy (HIPEC). *Ejso* 2010; 36: 463-469.
23. Bonastre J, Chevalier J, Elias D et al. Cost-effectiveness of intraperitoneal chemohyperthermia in the treatment of peritoneal carcinomatosis from colorectal cancer. *Value in Health* 2008; 11: 347-353.
24. Chua TC, Martin S, Saxena A et al. Evaluation of the cost-effectiveness of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy (peritonectomy) at the st george hospital peritoneal surface malignancy program. *Annals of Surgery* 2010; 251: 323-329.
25. Lee Z, Chia C, Teo M. Is cytoreductive surgery and hyperthermic intraperitoneal chemotherapy cost-effective for metastatic colorectal cancer? *Annals of Surgical Oncology* 2016; 23: S77.
26. Ludwigs K, Breimer ME, Brorson F et al. Cytoreductive surgery and intraperitoneal chemotherapy (HIPEC or EPIC) in patients with colorectal adenocarcinoma and peritoneal carcinomatosis. (Structured abstract). *Health Technology Assessment Database* 2013.
27. Naffouje SA, O'Donoghue C, Salti GI. Evaluation of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy in a community setting: A cost-utility analysis of a hospital's initial experience and reflections on the health care system. *Journal of Surgical Oncology* 2016; 113: 544-547.
28. Tentes AAK, Pallas N, Korakianitis O et al. The cost of cytoreductive surgery and perioperative intraperitoneal chemotherapy in the treatment of peritoneal malignancy in one Greek institute. *Journal of B.U.ON.* 2012; 17: 776-780.
29. Cao C, Yan TD, Black D, Morris DL. A systematic review and meta-analysis of cytoreductive surgery with perioperative intraperitoneal chemotherapy for peritoneal carcinomatosis of colorectal origin. *Annals of Surgical Oncology* 2009; 16: 2152-2165.
30. Ceelen WP, Hesse U, de Hemptinne B, Pattyn P. Hyperthermic intraperitoneal chemoperfusion in the treatment of locally advanced intra-abdominal cancer. *British Journal of Surgery* 2000; 87: 1006-1015.
31. Chia CS, Seshadri RA, Kepenekian V et al. Survival outcomes after Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis from gastric cancer: A systematic review. *Pleura and Peritoneum* 2016; 1: 67-77.
32. Chua TC, Esquivel J, Pelz JO, Morris DL. Summary of current therapeutic options for peritoneal metastases from colorectal cancer. *Journal of Surgical Oncology* 2013; 107: 566-573.
33. Di Vita M, Cappellani A, Piccolo G et al. The role of HIPEC in the treatment of peritoneal carcinomatosis from gastric cancer: Between lights and shadows. *Anti-Cancer Drugs* 2015; 26: 123-138.
34. Eveno C, Pocard M. Randomized controlled trials evaluating Cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC) in prevention and therapy of peritoneal metastasis: A Systematic review. *Pleura and Peritoneum* 2016; 1: 169-182.

35. He TF, Chen ZR, Xing CG. Cytoreductive surgery combined with intraperitoneal chemotherapy in the treatment of colorectal peritoneal metastasis: a meta-analysis. *International Journal of Clinical and Experimental Medicine* 2016; 9: 20562-20570.
36. Hotouras A, Desai D, Bhan C et al. Heated IntraPERitoneal Chemotherapy (HIPEC) for patients with recurrent ovarian cancer: A systematic literature review. *International Journal of Gynecological Cancer* 2016; 26: 661-670.
37. Huo YR, Richards A, Liauw W, Morris DL. Hyperthermic intraperitoneal chemotherapy (HIPEC) and cytoreductive surgery (CRS) in ovarian cancer: A systematic review and meta-analysis. *European Journal of Surgical Oncology* 2015; 41: 1578-1589.
38. Lopez-Lopez V, Cascales-Campos PA, Schneider MA et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy (HIPEC) in elderly patients. A systematic literature review. *Surgical Oncology-Oxford* 2016; 25: 378-384.
39. Mirnezami R, Moran BJ, Harvey K et al. Cytoreductive surgery and intraperitoneal chemotherapy for colorectal peritoneal metastases. *World Journal of Gastroenterology* 2014; 20: 14018-14032.
40. Kireeva GS, Gafton GI, Guseynov KD et al. HIPEC in patients with primary advanced ovarian cancer: Is there a role? A systematic review of short- and long-term outcomes. *Surg Oncol* 2018; 27: 251-258.
41. Morano WF, Khalili M, Chi DS et al. Clinical studies in CRS and HIPEC: Trials, tribulations, and future directions-A systematic review. *J Surg Oncol* 2018; 117: 245-259.
42. Whiting P, Savovic J, Higgins JP et al. ROBIS: A new tool to assess risk of bias in systematic reviews was developed. *J Clin Epidemiol* 2016; 69: 225-234.
43. Ubago-Pérez R, Matas-Hoces A, Beltrán-Calvo C, Romero-Tabares A. Hyperthermic intraperitoneal chemotherapy. Efficacy and safety in the treatment of ovarian cancer peritoneal carcino matosis (Structured abstract). *Health Technology Assessment Database* 2013.
44. Riley RD, Lambert PC, Abo-Zaid G. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ* 2010; 340.
45. Tierney JF, Vale C, Riley R et al. Individual Participant Data (IPD) Meta-analyses of Randomised Controlled Trials: Guidance on Their Use. *PLoS Med* 2015; 12: e1001855.
46. Clavien PA, Barkun J, de Oliveira ML et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg* 2009; 250: 187-196.
47. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004; 240: 205-213.
48. Therasse P, Arbuck SG, Eisenhauer EA et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000; 92: 205-216.
49. Higgins J, Green S, (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration. Available from www.handbook.cochrane.org 2011.
50. Higgins J, Sterne J, Savović J et al. A revised tool for assessing risk of bias in randomized trials. <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool> (accessed 14 May 2019) 2016.
51. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-188.
52. Ng JL, Ong WS, Chia CS et al. Prognostic Relevance of the Peritoneal Surface Disease Severity Score Compared to the Peritoneal Cancer Index for Colorectal Peritoneal Carcinomatosis. *Int J Surg Oncol* 2016; 2016: 2495131.
53. Esquivel J, Lowy AM, Markman M et al. The American Society of Peritoneal Surface Malignancies (ASPSM) Multiinstitution Evaluation of the Peritoneal Surface Disease Severity Score (PSDSS) in 1,013 Patients with Colorectal Cancer with Peritoneal Carcinomatosis. *Annals of Surgical Oncology* 2014; 21: 4195-4201.

54. Newell DJ. Intention-to-treat analysis: implications for quantitative and qualitative research. *Int J Epidemiol* 1992; 21: 837-841.
55. National Institute for Health and Care Excellence. Guide to the methods of technology appraisal 2013. <https://www.nice.org.uk/process/pmg9/chapter/foreword> (accessed on 4th June 2018) 2013.
56. Gurusamy KS, Riviere D, van Laarhoven CJH et al. Cost-effectiveness of laparoscopic versus open distal pancreatectomy for pancreatic cancer. *PLoS One* 2017; 12: e0189631.