Table S1. Studies excluded following full-text screening

	Reference	Reason for exclusion
1	Aaronson JA, van Bennekom CA, Hofman WF, van Bezeij T, van den Aardweg JG, Groet E, et al. Obstructive Sleep Apnea is Related to Impaired Cognitive and Functional Status after Stroke. <i>Sleep</i> 2015; 38 :1431-7. https://doi.org/10.5665/sleep.4984	Phenomenon of interest
2	Alastair A, Fiona OM. Issues in the definition and implementation of "best practice" for staff delivery of interventions for challenging behaviour. <i>Journal of Intellectual and Developmental Disability</i> 2001; 26 :243-56.	Outcome
3	Alderman N, Knight C, Birkett-Swan L. Inappropriate sexual behavior and aggression observed within a neurobehavioral rehabilitation service: Sasba and OAS-MNR outcomes over a three-month period. <i>Journal of Cyber Therapy and Rehabilitation</i> 2009; 2 :205-20.	Phenomenon of interest
4	Alderman N, Knight C, Stewart I, Gayton A. Measuring behavioural outcome in neurodisability. <i>British Journal of Neuroscience Nursing</i> 2011; 7 :691-5.	Population
5	Alderman N, Major G, Brooks J. What can structured professional judgement tools contribute to management of neurobehavioural disability? Predictive validity of the Short-Term Assessment of Risk and Treatability (START) in acquired brain injury. <i>Neuropsychol Rehabil</i> 2018; 28 :448-65.	Validity only (i.e. does not measure reliability)
6	Alderman N, Pink AE, Williams C, Ramos SDS, Oddy M, Knight C, et al. Optimizing measurement for neurobehavioural rehabilitation services: A multisite comparison study and response to UKROC. Neuropsychol Rehabil 2019; 10.1080/09602011.2019.1582432:1-30. https://doi.org/10.1080/09602011.2019.1582432	Phenomenon of interest
7	Alderman N, Pink AE, Williams C, Ramos SDS, Oddy M, Knight C, et al. Optimizing measurement for neurobehavioural rehabilitation services: A multisite comparison study and response to UKROC. Neuropsychol Rehabil 2019; 10.1080/09602011.2019.1582432:1-30. https://doi.org/10.1080/09602011.2019.1582432	Phenomenon of interest
8	Alderman N, Williams C, Knight C, Wood RL. Measuring Change in Symptoms of Neurobehavioural Disability: Responsiveness of the St Andrew's-Swansea Neurobehavioural Outcome Scale. <i>Arch Clin Neuropsychol</i> 2017; 32 :951-62. https://doi.org/10.1093/arclin/acx026	Phenomenon of interest
9	Alderman N, Williams C, Wood RL. When normal scores don't equate to independence: Recalibrating ratings of neurobehavioural disability from the 'St Andrew's - Swansea Neurobehavioural Outcome Scale' to reflect context-dependent support. <i>Brain Inj</i> 2018; 32 :218-29. https://doi.org/10.1080/02699052.2017.1406989	Validity only (i.e. does not measure reliability)
10	Alderman N, Wood RL, Williams C. The development of the St Andrew's-Swansea Neurobehavioural Outcome Scale: validity and reliability of a new measure of neurobehavioural disability and social handicap. Brain Inj 2011;25:83-100. https://doi.org/10.3109/02699052.2010.532849	Study duplicated in included systematic review

11	Alderson AL, Novack TA. Reliable serial measurement of cognitive processes in rehabilitation: the Cognitive Log. <i>Arch Phys Med Rehabil</i> 2003; 84 :668-72. https://doi.org/10.1016/s0003-9993(02)04842-6	Outcome
12	Allely CS. Prevalence and assessment of traumatic brain injury in prison inmates: A systematic PRISMA review. <i>Brain Inj</i> 2016; 30 :1161-80. https://doi.org/10.1080/02699052.2016.1191674	Study type
13	Alway Y, Gould KR, Johnston L, McKenzie D, Ponsford J. A prospective examination of Axis I psychiatric disorders in the first 5 years following moderate to severe traumatic brain injury. <i>Psychol Med</i> 2016; 46 :1331-41. https://doi.org/10.1017/S0033291715002986	Phenomenon of interest
14	Andaloro RR. The effects of diabetes, hypertension, and hypercholesterolemia on the severity of traumatic brain injury [PhD]. Indiana: Indiana University of Pennsylvania; 2012.	Phenomenon of interest
15	Andelic N, Sigurdardottir S, Schanke AK, Sandvik L, Sveen U, Roe C. Disability, physical health and mental health 1 year after traumatic brain injury. <i>Disabil Rehabil</i> 2010; 32 :1122-31. https://doi.org/10.3109/09638280903410722	Phenomenon of interest
16	Appelros P. Characteristics of Mini-Mental State Examination 1 year after stroke. <i>Acta Neurol Scand</i> 2005; 112 :88-92. https://doi.org/10.1111/j.1600-0404.2005.00441.x	Population
17	Armengol CG. Acute oxygen deprivation: neuropsychological profiles and implications for rehabilitation. Brain Inj 2000;14:237-50. https://doi.org/10.1080/026990500120718	Phenomenon of interest
18	Awad CP. Establishing the validity of the Neurobehavioral Functioning Inventory. Missouri: University of Missouri-Columbia; 2003.	Study duplicated in included systematic review
19	Bailie JM, King LC, Kinney D, Nitch SR. The relationship between self-reported neuropsychological risk factors and RBANS test performance among forensically committed psychiatric inpatients. <i>Appl Neuropsychol Adult</i> 2012; 19 :279-86. https://doi.org/10.1080/09084282.2012.670146	Phenomenon of interest
20	Baird Alison E. A three-item scale for the early prediction of stroke recovery. <i>Lancet</i> ; 357 :2095-9.	Outcome
21	Balasch IBM, Balasch IPS, Noe Sebastian E, Duenas Moscardo L, Ferri Campos J, Lopez-Bueno L. Study of the Recovery Patterns of Elderly Subacute Stroke Patients in an Interdisciplinary Neurorehabilitation Unit. <i>J Stroke Cerebrovasc Dis</i> 2015; 24 :2213-8. https://doi.org/10.1016/j.jstrokecerebrovasdis.2015.05.014	Phenomenon of interest
22	Ballard C, Stephens S, Kenny R, Kalaria R, Tovee M, O'Brien J. Profile of neuropsychological deficits in older stroke survivors without dementia. <i>Dement Geriatr Cogn Disord</i> 2003; 16 :52-6. https://doi.org/10.1159/000069994	Outcome
23	Bartolo M, Zucchella C, Tortola P, Spicciato F, Sandrini G, Pierelli F. Clinical scales for measuring stroke rehabilitation promote functional recovery by supporting teamwork. <i>Eur J Phys Rehabil Med</i> 2016; 52 :195-202.	Outcome
24	Beck KD. Personality and the prediction of outcome following rehabilitation in persons with acquired brain	Phenomenon of

	injuries: the Millon Behavioral Medicine Diagnostic (MBMD) [PhD]. Texas: University of North Texas; 2008.	interest
25	Belanger HG, Vanderploeg RD, Soble JR, Richardson M, Groer S. Validity of the Veterans Health	Phenomenon of
	Administration's traumatic brain injury screen. Arch Phys Med Rehabil 2012; 93 :1234-9.	interest
	https://doi.org/10.1016/j.apmr.2012.03.003	
26	Bennett HE, Thomas SA, Austen R, Morris AM, Lincoln NB. Validation of screening measures for assessing	Phenomenon of
	mood in stroke patients. <i>Br J Clin Psychol</i> 2006; 45 :367-76. https://doi.org/10.1348/014466505x58277	interest
27	Berthier ML, Kulisevsky JJ, Gironell A, Lopez OL. Obsessivecompulsive disorder and traumatic brain injury:	Study type
	behavioral, cognitive, and neuroimaging findings. Neuropsychiatry Neuropsychol Behav Neurol 2001;14:23-	
	31.	
28	Bertisch HC, Long C, Langenbahn DM, Rath JF, Diller L, Ashman T. Anxiety as a primary predictor of	Phenomenon of
	functional impairment after acquired brain injury: a brief report. Rehabil Psychol 2013;58:429-35.	interest
	https://doi.org/10.1037/a0034554	
29	Bezeau SC, Bogod NM, Mateer CA. Sexually intrusive behaviour following brain injury: approaches to	Study type
	assessment and rehabilitation. <i>Brain Inj</i> 2004; 18 :299-313. https://doi.org/10.1080/02699050310001617398	
30	Blake H, McKinney M, Treece K, Lee E, Lincoln NB. An evaluation of screening measures for cognitive	Phenomenon of
	impairment after stroke. Age Ageing 2002;31:451-6. https://doi.org/10.1093/ageing/31.6.451	interest
31	Blakey SM, Wagner HR, Naylor J, Brancu M, Lane I, Sallee M, et al. Chronic Pain, TBI, and PTSD in Military	Study type
	Veterans: A Link to Suicidal Ideation and Violent Impulses? <i>J Pain</i> 2018; 19 :797-806.	
	https://doi.org/10.1016/j.jpain.2018.02.012	
32	Boakye NT, Scott R, Parsons A, Betteridge S, Smith MA, Cluckie G. All change: a stroke inpatient service's	Phenomenon of
	experience of a new clinical neuropsychology delivery model. BMJ Open Quality 2019;8:e000184.	interest
33	Boan BK. The relationship between rehabilitation services, cognitive status and functional ability post brain	Phenomenon of
	injury: Adler University; 2012.	interest
34	Bogner J, Corrigan JD. Reliability and predictive validity of the Ohio State University TBI identification	Phenomenon of
	method with prisoners. <i>J Head Trauma Rehabil</i> 2009; 24 :279-91.	interest
	https://doi.org/10.1097/HTR.0b013e3181a66356	
35	Bogner JA, Corrigan JD, Mysiw WJ, Clinchot D, Fugate L. A comparison of substance abuse and violence in	Outcome
	the prediction of long-term rehabilitation outcomes after traumatic brain injury. Arch Phys Med Rehabil	
	2001; 82 :571-7. https://doi.org/10.1053/apmr.2001.22340	
36	Bogner JA, Whiteneck GG, MacDonald J, Juengst SB, Brown AW, Philippus AM, et al. Test-Retest Reliability of	Phenomenon of
	Traumatic Brain Injury Outcome Measures: A Traumatic Brain Injury Model Systems Study. J Head Trauma	interest
	Rehabil 2017; 32 :E1-E16. https://doi.org/10.1097/HTR.000000000000291	
37	Bond J, Gregson B, Smith M, Lecouturier J, Rousseau N, Rodgers H. Predicting place of discharge from	Population
	hospital for patients with a stroke or hip fracture on admission. J Health Serv Res Policy 2000;5:133-9.	

	https://doi.org/10.1177/135581960000500303	
38	Bondari S, Bondari D, Pirscoveanu M, Morosanu DV, Musetescu AE, Tudorica V, et al. Study on cognitive	Country
	decline in patients diagnosed with brain tumors. Rom J Morphol Embryol 2017;58:1185-92.	
39	Borgaro SR, Kwasnica C, Cutter N, Alcott S. The use of the BNI screen for higher cerebral functions in	Outcome
	assessing disorientation after traumatic brain injury. J Head Trauma Rehabil 2003;18:284-91.	
	https://doi.org/10.1097/00001199-200305000-00006	
40	Borgaro SR, Prigatano GP. Modification of the Patient Competency Rating Scale for use on an acute	Phenomenon of
	neurorehabilitation unit: the PCRS-NR. <i>Brain Inj</i> 2003; 17 :847-53.	interest
	https://doi.org/10.1080/0269905031000089350	
41	Borgaro SR, Prigatano GP, Alcott S, Kwasnica C, Cutter N. The Patient Distress Scale questionnaire: factor	Outcome
	structure and internal consistency. Brain Inj 2003;17:545-51.	
	https://doi.org/10.1080/0269905031000070206	
42	Borgaro SR, Prigatano GP, Kwasnica C, Rexer JL. Cognitive and affective sequelae in complicated and	Phenomenon of
	uncomplicated mild traumatic brain injury. <i>Brain Inj</i> 2003; 17 :189-98.	interest
	https://doi.org/10.1080/0269905021000013183	
43	Bowen C. Family therapy and neuro-rehabilitation: forging a linkincluding commentary by Charles N,	Study type
	Butera-Prinzi F and Perlesz A. International Journal of Therapy & Rehabilitation 2007; 14 :344-9.	
44	Brands I, Bol Y, Stapert S, Kohler S, van H. Is the effect of coping styles disease specific? Relationships with	Phenomenon of
	emotional distress and quality of life in acquired brain injury and multiple sclerosis. <i>Clin Rehabil</i>	interest
4.5	2018; 32 :116-26.	0.1
45	Brenner LA, Carlson NE, Harrison-Felix C, Ashman T, Hammond FM, Hirschberg RE. Self-inflicted traumatic	Outcome
	brain injury: Characteristics and outcomes. <i>Brain Inj</i> 2009; 23 :991-8. https://doi.org/10.3109/02699050903379362	
46	Brenner LA, Homaifar BY, Olson-Madden JH, Nagamoto HT, Huggins J, Schneider AL, et al. Prevalence and	Outcome
40	screening of traumatic brain injury among veterans seeking mental health services. J Head Trauma Rehabil	Outcome
	2013; 28 :21-30. https://doi.org/10.1097/HTR.0b013e31827df0b5	
47	Brickell TA, Lange RT, French LM. Health-related quality of life within the first 5 years following military-	Phenomenon of
7/	related concurrent mild traumatic brain injury and polytrauma. <i>Mil Med</i> 2014; 179 :827-38.	interest
	https://doi.org/10.7205/MILMED-D-13-00506	
48	Brooks BL, Holdnack JA, Iverson GL. Advanced clinical interpretation of the WAIS-IV and WMS-IV: prevalence	Phenomenon of
	of low scores varies by level of intelligence and years of education. <i>Assessment</i> 2011; 18 :156-67.	interest
	https://doi.org/10.1177/1073191110385316	
49	Bryant Richard A, O'Donnell Meaghan L, Creamer M, McFarlane Alexander C, Clark C, Silove D. The	Phenomenon of
	psychiatric sequelae of traumatic injury. <i>Am J Psychiatry</i> 2010; 167 :312-20.	interest

50	Campbell N, Rice D, Friedman L, Speechley M, Teasell RW. Screening and facilitating further assessment for cognitive impairment after stroke: application of a shortened Montreal Cognitive Assessment (miniMoCA). <i>Disabil Rehabil</i> 2016; 38 :601-4. https://doi.org/10.3109/09638288.2015.1047968	Phenomenon of interest
51	Caplain S, Truelle J-L, Hinglais E, Baarir N, Vignaud F, Rozec G, et al. After a mild traumatic injury, can a persistent post-concussion syndrome be predicted? A prospective clinical study on 55 cases. <i>Acta Neuropsychologica</i> 2010; 8 :123-41.	Phenomenon of interest
52	Carlozzi NE, Kirsch NL, Kisala PA, Tulsky DS. An examination of the Wechsler Adult Intelligence Scales, Fourth Edition (WAIS-IV) in individuals with complicated mild, moderate and Severe traumatic brain injury (TBI). <i>Clin Neuropsychol</i> 2015; 29 :21-37. https://doi.org/10.1080/13854046.2015.1005677	Phenomenon of interest
53	Carroll E, Coetzer R. Identity, grief and self-awareness after traumatic brain injury. <i>Neuropsychol Rehabil</i> 2011; 21 :289-305. https://doi.org/10.1080/09602011.2011.555972	Phenomenon of interest
54	Carroll Linda J, Cassidy JD, Garritty C, Giles-Smith L, Peloso Paul M. Systematic search and review procedures: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. <i>J Rehabil Med</i> 2004; 36 :11-4.	Study type
55	Castano Monsalve B, Laxe S, Bernabeu Guitart M, Vilarrasa AB, Quemada JI. Behavioral scales used in severe and moderate traumatic brain injury. <i>NeuroRehabilitation</i> 2014; 35 :67-76. https://doi.org/10.3233/NRE-141103	Study type
56	Cattran C, Oddy M, Ramos SDS, Goodson A, Wood R. The development of a measure of social cognition following acquired brain injury. <i>Neuropsychol Rehabil</i> 2018; 28 :633-48. https://doi.org/10.1080/09602011.2016.1202121	Phenomenon of interest
57	Cattran C, Oddy M, Wood R. The development of a measure of emotional regulation following acquired brain injury. <i>J Clin Exp Neuropsychol</i> 2011; 33 :672-9. https://doi.org/10.1080/13803395.2010.550603	Phenomenon of interest
58	Cattran CJ, Oddy M, Wood RL, Moir JF. Post-injury personality in the prediction of outcome following severe acquired brain injury. <i>Brain Inj</i> 2011; 25 :1035-46. https://doi.org/10.3109/02699052.2011.607787	Outcomes
59	Chapman JC, Andersen AM, Roselli LA, Meyers NM, Pincus JH. Screening for mild traumatic brain injury in the presence of psychiatric comorbidities. <i>Arch Phys Med Rehabil</i> 2010; 91 :1082-6. https://doi.org/10.1016/j.apmr.2010.03.018	Phenomenon of interest
60	Chen YK, Wong KS, Mok V, Ungvari GS, Tang WK. Health-related quality of life in patients with poststroke emotional incontinence. <i>Arch Phys Med Rehabil</i> 2011; 92 :1659-62. https://doi.org/10.1016/j.apmr.2011.04.016	Phenomenon of interest
61	Cheng C, Chi NC, Williams E, Thompson HJ. Examining age-related differences in functional domain impairment following traumatic brain injury. <i>Int J Older People Nurs</i> 2018; 13 :e12208. https://doi.org/10.1111/opn.12208	Phenomenon of interest
62	Clark-Wilson J, Giles GM, Seymour S, Tasker R, Baxter DM, Holloway M. Factors influencing community case	Phenomenon of

	management and care hours for clients with traumatic brain injury living in the UK. <i>Brain Inj</i> 2016; 30 :872-82. https://doi.org/10.3109/02699052.2016.1146799	interest
63	ClinicalTrials.gov. Executive Dysfunction and Suicide in Psychiatric Outpatients and Inpatients. NCT01043432; 2015. URL: https://clinicaltrials.gov/ct2/show/NCT01043432 (accessed 19 Feb, 2020).	Unable to retrieve full- text
64	Coban E, Mutluay B, Sen A, Keskek A, Atakl D, Soysal A. Characteristics, diagnosis and outcome of patients referred to a specialized neurology emergency clinic: prospective observational study. <i>Ann Saudi Med</i> 2016; 36 :51-6. https://doi.org/10.5144/0256-4947.2016.51	Outcome
65	Colantonio A, Stamenova V, Abramowitz C, Clarke D, Christensen B. Brain injury in a forensic psychiatry population. <i>Brain Inj</i> 2007; 21 :1353-60. https://doi.org/10.1080/02699050701785054	Study type
66	Comerford VE, Geffen GM, May C, Medland SE, Geffen LB. A rapid screen of the severity of mild traumatic brain injury. <i>J Clin Exp Neuropsychol</i> 2002; 24 :409-19. https://doi.org/10.1076/jcen.24.4.409.1044	Outcome
67	Cooper-Evans S, Alderman N, Knight C, Oddy M. Self-esteem as a predictor of psychological distress after severe acquired brain injury: an exploratory study. <i>Neuropsychol Rehabil</i> 2008; 18 :607-26. https://doi.org/10.1080/09602010801948516	Phenomenon of interest
68	Corvo K, Halpern J, Ferraro FR. Frontal lobe deficits and alcohol abuse: possible interactions in predicting domestic violence. <i>J Aggress Maltreat Trauma</i> 2006; 13 :49-63.	Study type
69	Cullen NK, Crescini C, Bayley MT. Rehabilitation outcomes after anoxic brain injury: a case-controlled comparison with traumatic brain injury. <i>PM R</i> 2009; 1 :1069-76. https://doi.org/10.1016/j.pmrj.2009.09.013	Phenomenon of interest
70	Cusimano MD, Holmes SA, Sawicki C, Topolovec-Vranic J. Assessing aggression following traumatic brain injury: a systematic review of validated aggression scales. <i>J Head Trauma Rehabil</i> 2014; 29 :172-84. https://doi.org/10.1097/HTR.0b013e31827c7d15	Study type
71	Dailey NS, Smith R, Bajaj S, Alkozei A, Gottschlich MK, Raikes AC, et al. Elevated Aggression and Reduced White Matter Integrity in Mild Traumatic Brain Injury: A DTI Study. Front Behav Neurosci 2018;12:118. https://doi.org/10.3389/fnbeh.2018.00118	Study type
72	Daniels JE, Wirth JB, Herrera DG, Simpson EB, Auchincloss EL, Occhiogrosso MB, et al. Head banging on an inpatient psychiatric unit: a vicious circle. <i>Harv Rev Psychiatry</i> 2007; 15 :70-9. https://doi.org/10.1080/10673220701307588	Study type
73	de Guise E, le Blanc J, Feyz M, Meyer K, Duplantie J, Thomas H, et al. Long-term outcome after severe traumatic brain injury: the McGill interdisciplinary prospective study. J Head Trauma Rehabil 2008;23:294-303. https://doi.org/10.1097/01.HTR.0000336842.53338.f4	Outcome
74	Deb S. Almost half of people suffering traumatic brain injury may later be diagnosed with axis I disorders. Evidence Based Mental Health 2003; 6 :59	Study type
75	Demeyere N, Riddoch MJ, Slavkova ED, Bickerton WL, Humphreys GW. The Oxford Cognitive Screen (OCS): validation of a stroke-specific short cognitive screening tool. <i>Psychol Assess</i> 2015; 27 :883-94.	Phenomenon of interest

	https://doi.org/10.1037/pas0000082	
76	Dennis JP, Ghahramanlou-Holloway M, Cox DW, Brown GK. A guide for the assessment and treatment of	Study type
	suicidal patients with traumatic brain injuries. J Head Trauma Rehabil 2011; 26 :244-56.	
	https://doi.org/10.1097/HTR.0b013e3182225528	
77	Diaz AP, Schwarzbold ML, Thais ME, Hohl A, Bertotti MM, Schmoeller R, et al. Psychiatric disorders and	Country
	health-related quality of life after severe traumatic brain injury: a prospective study. J Neurotrauma	
	2012; 29 :1029-37. https://doi.org/10.1089/neu.2011.2089	
78	Dickens G, Alderman N, Bowers L. Potential severity of aggressive behaviour after acquired brain injury:	Phenomenon of
	implications for recording. <i>J Psychiatr Ment Health Nurs</i> 2011; 18 :586-94. https://doi.org/10.1111/j.1365-	interest
	2850.2011.01707.x	
79	Dickens G, Picchioni M, Long C. Aggression in specialist secure and forensic inpatient mental health care:	Population
	incidence across care pathways. <i>The Journal of Forensic Practice</i> 2013; 15 :206-17.	
90	https://doi.org/10.1108/jfp-09-2012-0017	Demulation
80	Dinn WM, Gansler DA, Moczynski N, Fulwiler C. Brain Dysfunction and Community Violence in Patients With Major Mental Illness. <i>Crim Justice Behav</i> 2008; 36 :117-36. https://doi.org/10.1177/0093854808327507	Population
81	Doninger NA, Ehde DM, Bode RK, Knight K, Bombardier CH, Heinemann AW. Measurement properties of the	Outcome
01	Neurobehavioral Cognitive Status Examination (Cognistat) in traumatic brain injury rehabilitation. <i>Rehabil</i>	Outcome
	Psychol 2006; 51 :281-8. https://doi.org/10.1037/0090-5550.51.4.281	
82	Dowler RN, Bush BA, Novack TA, Jackson WT. Cognitive orientation in rehabilitation and neuropsychological	Outcome/
<u> </u>	outcome after traumatic brain injury. <i>Brain Inj</i> 2000; 14 :117-23. https://doi.org/10.1080/026990500120781	Phenomenon of
	, , , , , , , , , , , , , , , , , , ,	interest
83	Dreer LE, Tang X, Nakase-Richardson R, Pugh MJ, Cox MK, Bailey EK, et al. Suicide and traumatic brain injury:	Study type
	a review by clinical researchers from the National Institute for Disability and Independent Living	
	Rehabilitation Research (NIDILRR) and Veterans Health Administration Traumatic Brain Injury Model	
	Systems. Curr Opin Psychol 2018; 22 :73-8. https://doi.org/10.1016/j.copsyc.2017.08.030	
84	Edge D, Walker T, Meacock R, Wilson H, McNair L, Shaw J, et al. Secure pathways for women in the UK:	Population
	lessons from the women's enhanced medium secure services (WEMSS) pilots. J Forens Psychiatry Psychol	
	2016; 28 :206-25. https://doi.org/10.1080/14789949.2016.1244279	
85	Fazel S, Lichtenstein P, Grann M, Langstrom N. Risk of violent crime in individuals with epilepsy and	Population
	traumatic brain injury: a 35-year Swedish population study. <i>PLoS Med</i> 2011; 8 :e1001150.	
	https://doi.org/10.1371/journal.pmed.1001150	
86	Fergus G. Characterising neuropsychological rehabilitation service users for service design. <i>Social Care and</i>	Phenomenon of
	Neurodisability 2014; 5 :16-28.	interest
87	Ferguson SD, Coccaro EF. History of mild to moderate traumatic brain injury and aggression in physically	Population

	health martining to with and with and with a second life discussion of the second seco	
	healthy participants with and without personality disorder. <i>J Pers Disord</i> 2009; 23 :230-9.	
	https://doi.org/10.1521/pedi.2009.23.3.230	
88	Finn JA, Lamberty GJ, Tang X, Saylors ME, Stevens LF, Kretzmer T. Postrehabilitation Mental Health	Phenomenon of
	Treatment Utilization in Veterans With Traumatic Brain Injury: A VA TBI Model Systems Study. J Head	interest
	Trauma Rehabil 2018; 33 :E1-E9. https://doi.org/10.1097/HTR.00000000000357	
89	Fitzgerald A, Aditya H, Prior A, McNeill E, Pentland B. Anoxic brain injury: Clinical patterns and functional	Phenomenon of
	outcomes. A study of 93 cases. <i>Brain Inj</i> 2010; 24 :1311-23. https://doi.org/10.3109/02699052.2010.506864	interest
90	Foebel AD, Hirdes JP, Heckman GA, Kergoat MJ, Patten S, Marrie RA, et al. Diagnostic data for neurological	Phenomenon of
	conditions in interRAI assessments in home care, nursing home and mental health care settings: a validity	interest
	study. BMC Health Serv Res 2013;13:457. https://doi.org/10.1186/1472-6963-13-457	
91	Frank B, Schlote A, Hasenbein U, Wallesch CW. Prognosis and prognostic factors in ADL-dependent stroke	Phenomenon of
	patients during their first in-patient rehabilitationa prospective multicentre study. Disabil Rehabil	interest
	2006; 28 :1311-8. https://doi.org/10.1080/09638280600633597	
92	Gagnon J, Rochat L, Messier F, Chiocchio F, Sordes C, Beaulieu J, et al. Development and validation of a task	Non-English language
	to detect the risk of showing socially inappropriate behavior following a craniocerebral trauma: the task of	study
	social decision. Canadian Journal of Behavioural Science-Revue Canadienne Des Sciences Du Comportement	
	2017; 49 :100-11. https://doi.org/10.1037/cbs0000067	
93	Gagnon J, Simpson GK, Kelly G, Godbout D, Ouellette M, Drolet J. A French adaptation of the Overt	Non-English language
	Behaviour Scale (OBS) measuring challenging behaviours following acquired brain injury: The Echelle des	study
	comportements observables (ECO). Brain Inj 2016; 30 :1019-25.	
	https://doi.org/10.3109/02699052.2016.1148197	
94	Gao L, Li SC, Xia L, Pan S, Velakoulis D, Walterfang M. Validation of the Chinese version of the NUCOG	Country
	cognitive screening tool in patients with epilepsy, dementia and other neurological disorders. J Clin Neurosci	
	2014; 21 :980-7. https://doi.org/10.1016/j.jocn.2013.09.020	
95	Gass CS, Luis CA. MMPI-2 short form: psychometric characteristics in a neuropsychological setting.	Population
	Assessment 2001;8:213-9. https://doi.org/10.1177/107319110100800209	
96	Ghika-Schmid F, Bogousslavsky J. The acute behavioral syndrome of anterior thalamic infarction: a	Phenomenon of
	prospective study of 12 cases. Ann Neurol 2000;48:220-7.	interest
97	Ghose SS. The effects of post-stroke depression on inpatient and outpatient medical utilization: A	Phenomenon of
	retrospective database study. Indiana: Indiana University; 2002.	interest
98	Glover N, Gorgens K, Lehto M, Meyer L, Detmer J, Gaford J. Sensitivity and Specificity of the Ohio State	Outcome
	University Traumatic Brain Injury Identification Method to Neuropsychological Impairment. Crim Justice	
	Behav 2018; 45 :885-901. https://doi.org/10.1177/0093854818765043	
99	Gould KR, Ponsford JL, Johnston L, Schonberger M. Relationship between psychiatric disorders and 1-year	Phenomenon of

	psychosocial outcome following traumatic brain injury. <i>J Head Trauma Rehabil</i> 2011; 26 :79-89. https://doi.org/10.1097/HTR.0b013e3182036799	interest
100	Gould KR, Ponsford JL, Johnston L, Schonberger M. The nature, frequency and course of psychiatric disorders in the first year after traumatic brain injury: a prospective study. <i>Psychol Med</i> 2011; 41 :2099-109. https://doi.org/10.1017/S003329171100033X	Phenomenon of interest
101	Gracey F, Malley D, P. Wagner A, Clare I. Characterising neuropsychological rehabilitation service users for service design. <i>Social Care and Neurodisability</i> 2014; 5 :16-28. https://doi.org/10.1108/scn-09-2013-0034	Phenomenon of interest
102	Gralton E, Bernard SH. Inpatient assessment of young people with developmental disabilities who offend. Adv Ment Health Intellect Disabil 2013;7:108-16. https://doi.org/10.1108/20441281311310207	Study type
103	Grauwmeijer E, Heijenbrok-Kal MH, Peppel LD, Hartjes CJ, Haitsma IK, de Koning I, et al. Cognition, Health-Related Quality of Life, and Depression Ten Years after Moderate to Severe Traumatic Brain Injury: A Prospective Cohort Study. J Neurotrauma 2018; 35 :1543-51. https://doi.org/10.1089/neu.2017.5404	Phenomenon of interest
104	Greenall PV, Jellicoe-Jones L. Themes and risk of sexual violence among the mentally ill: implications for understanding and treatment. <i>Sexual and Relationship Therapy</i> 2007; 22 :323-37. https://doi.org/10.1080/14681990701391269	Population
105	Gualtieri CT, Johnson LG. A computerized test battery sensitive to mild and severe brain injury. <i>Medscape J Med</i> 2008; 10 :90.	Outcome
106	Hanks RA, Jackson AM, Crisanti LK. Predictive validity of a brief outpatient neuropsychological battery in individuals 1-25 years post traumatic brain injury. <i>Clin Neuropsychol</i> 2016; 30 :1074-86. https://doi.org/10.1080/13854046.2016.1194479	Phenomenon of interest
107	Harbinson M, Zarshenas S, Cullen NK. Long-Term Functional and Psychosocial Outcomes After Hypoxic-Ischemic Brain Injury: A Case-Controlled Comparison to Traumatic Brain Injury. <i>PM R</i> 2017; 9 :1200-7. https://doi.org/10.1016/j.pmrj.2017.04.015	Phenomenon of interest
108	Hashmi F, Krady A, Qayum F, Grossberg G. Sexually disinhibited behavior in the cognitively impaired elderly. <i>Clin Geriatr</i> 2000; 8 :61-8.	Study type
109	Haushalter J. Cognitive Distortions and Antisocial Behavior among Adults with Traumatic Brain Injury [UG Thesis]. Ohio: Ohio State University; 2015.	Phenomenon of interest
110	Hawley CA, Maden A. Mentally disordered offenders with a history of previous head injury: are they more difficult to discharge? <i>Brain Inj</i> 2003; 17 :743-58.	Phenomenon of interest
111	Heald A, Parr C, Gibson C, O'Driscoll K, Fowler H. A cross-sectional study to investigate long-term cognitive function in people with treated pituitary Cushing's disease. <i>Exp Clin Endocrinol Diabetes</i> 2006; 114 :490-7. https://doi.org/10.1055/s-2006-924332	Phenomenon of interest
112	Healey AK, Kneebone, II, Carroll M, Anderson SJ. A preliminary investigation of the reliability and validity of the Brief Assessment Schedule Depression Cards and the Beck Depression Inventory-Fast Screen to screen	Phenomenon of interest

	for depression in older stroke survivors. <i>Int J Geriatr Psychiatry</i> 2008; 23 :531-6. https://doi.org/10.1002/gps.1933	
113	Hellweg S, Schuster-Amft C. German version, inter-and intrarater reliability and internal consistency of the "Agitated Behavior Scale" (ABS-G) in patients with moderate to severe traumatic brain injury. <i>Health and quality of life outcomes</i> 2016; 14 :106.	Validity only (i.e. does not measure reliability)
114	Herman MA, Tremont-Lukats I, Meyers CA, Trask DD, Froseth C, Renschler MF, et al. Neurocognitive and functional assessment of patients with brain metastases: a pilot study. <i>Am J Clin Oncol</i> 2003; 26 :273-9. https://doi.org/10.1097/01.COC.0000020585.85901.7C	Outcome
115	Hernandez-Cardenache R. <i>The relationship between the Glasgow Coma Scale and the Functional Independence Measure in a sample of traumatic brain injury patients in a neurorehabilitation setting</i> . Malibu: Pepperdine University; 2007.	Outcome
116	Hicks AJ, Gould KR, Hopwood M, Kenardy J, Krivonos I, Ponsford JL. Behaviours of concern following moderate to severe traumatic brain injury in individuals living in the community. <i>Brain Inj</i> 2017; 31 :1312-9. https://doi.org/10.1080/02699052.2017.1317361	Outcome
117	Holtzer R, Burright RG, Lynn SJ, Donovick PJ. Behavioural differences between psychiatric patients with confirmed versus non-confirmed traumatic brain injuries. <i>Brain Inj</i> 2000; 14 :959-73. https://doi.org/10.1080/02699050050191904	Phenomenon of interest
118	Huh PW, Yoo DS, Cho KS, Park CK, Kang SG, Park YS, et al. Diagnostic method for differentiating external hydrocephalus from simple subdural hygroma. <i>J Neurosurg</i> 2006; 105 :65-70. https://doi.org/10.3171/jns.2006.105.1.65	Outcome
119	Hunt AW, Turner GR, Polatajko H, Bottari C, Dawson DR. Executive function, self-regulation and attribution in acquired brain injury: A scoping review. <i>Neuropsychol Rehabil</i> 2013; 23 :914-32. https://doi.org/10.1080/09602011.2013.835739	Phenomenon of interest
120	Hux K, Schneider T, Bennett K. Screening for traumatic brain injury. Brain Inj 2009;23:8-14. https://doi.org/10.1080/02699050802590353	Phenomenon of interest
121	Intiso D, Lombardi T, Grimaldi G, Iarossi A, Tolfa M, Russo M, et al. Long-term outcome and health status in decompressive craniectomized patients with intractable intracranial pressure after severe brain injury. Brain Inj 2011; 25 :379-86. https://doi.org/10.3109/02699052.2011.558046	Phenomenon of interest
122	Jackson HF, Tunstall V, Hague G, Daniels L, Crompton S, Taplin K. The Behavioural Assessment of Self-Structuring (BASS): psychometric properties in a post-acute brain injury rehabilitation programme. NeuroRehabilitation 2014;34:695-708. https://doi.org/10.3233/NRE-141087	Outcome
123	Jennekens N, de Casterle BD, Dobbels F. A systematic review of care needs of people with traumatic brain injury (TBI) on a cognitive, emotional and behavioural level. <i>J Clin Nurs</i> 2010; 19 :1198-206. https://doi.org/10.1111/j.1365-2702.2009.03114.x	Phenomenon of interest

124	Jimenez M. An analysis of neuropsychological functioning and psychopathic traits in correctional patients	Population
125	with histories of suicidal behavior. Palo Alto: Palo Alto University; 2014. Johnson CI. Effect of self-reported head injury on cognitive ability within forensic population. Fresno: Alliant International University; 2016.	Phenomenon of interest
126	Kalmar K, Novack TA, Nakase-Richardson R, Sherer M, Frol AB, Gordon WA, et al. Feasibility of a brief neuropsychologic test battery during acute inpatient rehabilitation after traumatic brain injury. Arch Phys Med Rehabil 2008;89:942-9. https://doi.org/10.1016/j.apmr.2008.01.008	Outcome
127	Kean J, Malec JF, Altman IM, Swick S. Rasch measurement analysis of the Mayo-Portland Adaptability Inventory (MPAI-4) in a community-based rehabilitation sample. <i>J Neurotrauma</i> 2011; 28 :745-53. https://doi.org/10.1089/neu.2010.1573	Phenomenon of interest
128	Kelly G, Simpson GK, Brown S, Kremer P, Gillett L. The Overt Behaviour Scale-Self-Report (OBS-SR) for acquired brain injury: exploratory analysis of reliability and validity. <i>Neuropsychol Rehabil</i> 2019; 29 :704-22. https://doi.org/10.1080/09602011.2017.1322523	Study duplicated in included systematic review
129	Kelly G, Todd J, Simpson G, Kremer P, Martin C. The Overt Behaviour Scale (OBS): a tool for measuring challenging behaviours following ABI in community settings. <i>Brain Inj</i> 2006; 20 :307-19. https://doi.org/10.1080/02699050500488074	Study duplicated in included systematic review
130	Kennedy Mary R, Yorkston Kathryn M. The effects of frontal injury on "on-line" self-monitoring during verbal learning by adults with diffuse brain injury. <i>Neuropsychol Rehabil</i> 2004; 14 :449-65.	Outcome
131	Kennedy RE, Livingston L, Riddick A, Marwitz JH, Kreutzer JS, Zasler ND. Evaluation of the Neurobehavioral Functioning Inventory as a depression screening tool after traumatic brain injury. <i>J Head Trauma Rehabil</i> 2005; 20 :512-26. https://doi.org/10.1097/00001199-200511000-00004	Phenomenon of interest
132	Kilmer RP, Demakis GJ, Hammond FM, Grattan KE, Cook JR, Kornev AA. Use of the neuropsychiatric inventory in traumatic brain injury: A pilot investigation. <i>Rehabil Psychol</i> 2006; 51 :232-8. https://doi.org/10.1037/0090-5550.51.3.232	Study duplicated in included systematic review
133	King JT, Jr., Horowitz MB, Kassam AB, Yonas H, Roberts MS. The short form-12 and the measurement of health status in patients with cerebral aneurysms: performance, validity, and reliability. <i>J Neurosurg</i> 2005; 102 :489-94. https://doi.org/10.3171/jns.2005.102.3.0489	Phenomenon of interest
134	Knight C, Alderman N, Johnson C, Green S, Birkett-Swan L, Yorstan G. The St Andrew's Sexual Behaviour Assessment (SASBA): development of a standardised recording instrument for the measurement and assessment of challenging sexual behaviour in people with progressive and acquired neurological impairment. <i>Neuropsychol Rehabil</i> 2008; 18 :129-59. https://doi.org/10.1080/09602010701822381	Population
135	Knopf A. TBI 'sequelae' require special care by behavioral health providers. <i>Behavioral Healthcare</i> 2012; 32 :42-6.	Study type
136	Kolakowsky-Hayner SA, Gourley EV, 3rd, Kreutzer JS, Marwitz JH, Meade MA, Cifu DX. Post-injury substance	Phenomenon of

	abuse among persons with brain injury and persons with spinal cord injury. <i>Brain Inj</i> 2002; 16 :583-92. https://doi.org/10.1080/02699050110119475	interest
137	Koponen S, Taiminen T, Portin R, Himanen L, Isoniemi H, Heinonen H, et al. Axis I and II psychiatric disorders after traumatic brain injury: a 30-year follow-up study. <i>Am J Psychiatry</i> 2002; 159 :1315-21.	Phenomenon of interest
138	https://doi.org/10.1176/appi.ajp.159.8.1315 Kumari P, Mohsin H, Koola M. Dyke-Davidoff-Masson syndrome presenting with bipolar I mania with	Study type
	psychosis. In: Indian J Psychiatry; 2018: 149-51. https://doi.org/10.4103/psychiatry.IndianJPsychiatry_229_17	
139	Landre N, Poppe CJ, Kiolbasa T. Prevalence of neuropathological indicators among acute psychiatric inpatients. <i>J Nerv Ment Dis</i> 2001; 189 :263-5. https://doi.org/10.1097/00005053-200104000-00009	Study type
140	Lane KS, St Pierre ME, Lauterbach MD, Koliatsos VE. Patient Profiles of Criminal Behavior in the Context of Traumatic Brain Injury. <i>J Forensic Sci</i> 2017; 62 :545-8. https://doi.org/10.1111/1556-4029.13289	Study type
141	Langeluddecke PM, Lucas SK. Wechsler Adult Intelligence Scale-Third Edition findings in relation to severity of brain injury in litigants. <i>Clin Neuropsychol</i> 2003; 17 :273-84. https://doi.org/10.1076/clin.17.2.273.16499	Phenomenon of interest
142	Lannin NA, Cusick A, McLachlan R, Allaous J. Observed recovery sequence in neurobehavioral function after severe traumatic brain injury. <i>Am J Occup Ther</i> 2013; 67 :543-9. https://doi.org/10.5014/ajot.2013.008094	Phenomenon of interest
143	Lapadatu I, Morris R. The relationship between stroke survivors' perceived identity and mood, self-esteem and quality of life. <i>Neuropsychol Rehabil</i> 2019; 29 :199-213. https://doi.org/10.1080/09602011.2016.1272468	Phenomenon of interest
144	Larrabee Glenn J, Rohling Martin L. Neuropsychological Differential Diagnosis of Mild Traumatic Brain Injury. Behav Sci Law 2013; 31 :686-701.	Study type
145	Lauterbach Margo D, Notarangelo Paula L, Nichols Stephen J, Lane Kristy S, Koliatsos Vassilis E. Diagnostic and treatment challenges in traumatic brain injury patients with severe neuropsychiatric symptoms: Insights into psychiatric practice. <i>Neuropsychiatr Dis Treat</i> 2015; 11 .	Study type
146	Laxe S, Terre R, Leon D, Bernabeu M. How does dysautonomia influence the outcome of traumatic brain injured patients admitted in a neurorehabilitation unit? <i>Brain Inj</i> 2013; 27 :1383-7.	Outcome
147	Laxe S, Zasler N, Tschiesner U, Lopez-Blazquez R, Tormos JM, Bernabeu M. ICF use to identify common problems on a TBI neurorehabilitation unit in Spain. <i>NeuroRehabilitation</i> 2011; 29 :99-110. https://doi.org/10.3233/NRE-2011-0683	Outcome/study type
148	Leigh AJ, O'Hanlon K, Sheldrick R, Surr C, Hare DJ. Care mapping in clinical neuroscience settings: Cognitive impairment and dependency. <i>Neuropsychol Rehabil</i> 2015; 25 :574-92. https://doi.org/10.1080/09602011.2014.951366	Outcome
149	Liesbeth De C, et al. The sensitivity of somatic symptoms in post-stroke depression: a discriminant analytic approach. <i>Int J Geriatr Psychiatry</i> 2005; 20 :358-62.	Phenomenon of interest

150	Lincoln NB, Nicholl CR, Flannaghan T, Leonard M, Van der Gucht E. The validity of questionnaire measures for assessing depression after stroke. <i>Clin Rehabil</i> 2003; 17 :840-6. https://doi.org/10.1191/0269215503cr687oa	Phenomenon of interest
151	Ll Wood R, Alderman N, Williams C. Assessment of neurobehavioural disability: a review of existing measures and recommendations for a comprehensive assessment tool. <i>Brain Inj</i> 2008; 22 :905-18. https://doi.org/10.1080/02699050802491271	Study type
152	Lloyd-James L. Facing reality: discharge challenges after neuro-rehabilitation. <i>Paediatr Nurs</i> 2006; 18 :28. https://doi.org/10.7748/paed.18.6.28.s25	Study type
153	Longworth C, Deakins J, Rose D, Gracey F. The nature of self-esteem and its relationship to anxiety and depression in adult acquired brain injury. <i>Neuropsychol Rehabil</i> 2018; 28 :1078-94. https://doi.org/10.1080/09602011.2016.1226185	Phenomenon of interest
154	Lowings G, Trout S, Braham L. Seizing the opportunities to improve neuropsychological services in a High Secure hospital. <i>Adv Ment Health Intellect Disabil</i> 2013; 7 :356-64.	Population
155	Mahoney D, Gutman SA, Gillen G. A Scoping Review of Self-Awareness Instruments for Acquired Brain Injury. <i>Open J Occup Ther</i> 2019; 7 :1-15. https://doi.org/10.15453/2168-6408.1529	Phenomenon of interest
156	Mainio A, Hakko H, Niemela A, Koivukangas J, Rasanen P. Depression in relation to anxiety, obsessionality and phobia among neurosurgical patients with a primary brain tumor: a 1-year follow-up study. <i>Clin Neurol Neurosurg</i> 2011; 113 :649-53. https://doi.org/10.1016/j.clineuro.2011.05.006	Phenomenon of interest
157	Malec JF, Stump TE, Monahan PO, Kean J, Neumann D, Hammond FM. Rasch Analysis, Dimensionality, and Scoring of the Neuropsychiatric Inventory Irritability and Aggression Subscales in Individuals With Traumatic Brain Injury. <i>Arch Phys Med Rehabil</i> 2018; 99 :281-8 e2. https://doi.org/10.1016/j.apmr.2017.07.020	Study duplicated in included systematic review
158	Marcano-Cedeno A, Chausa P, Garcia A, Caceres C, Tormos JM, Gomez EJ. Artificial metaplasticity prediction model for cognitive rehabilitation outcome in acquired brain injury patients. <i>Artif Intell Med</i> 2013; 58 :91-9. https://doi.org/10.1016/j.artmed.2013.03.005	Phenomenon of interest
159	Marsh NV, Kersel DA. Frequency of behavioural problems at one year following traumatic brain injury: correspondence between patient and caregiver reports. <i>Neuropsychol Rehabil</i> 2006; 16 :684-94. https://doi.org/10.1080/09602010500220290	Outcome
160	Mateo MA, Glod CA, Hennen J, Price BH, Merrill N. Mild traumatic brain injury in psychiatric inpatients. <i>J Neurosci Nurs</i> 2005; 37 :28-33. https://doi.org/10.1097/01376517-200502000-00005	Study type
161	McAleese A, Wilson CF, McEvoy M, Caldwell S. Comparison of SMART and WHIM as measurement tools in routine assessment of PDOC patients. <i>Neuropsychol Rehabil</i> 2018; 28 :1266-74. https://doi.org/10.1080/09602011.2016.1264977	Outcome
162	McBride WF, rd, Crighton AH, Wygant DB, Granacher RP, Jr. It's not all in your head (or at least your brain): association of traumatic brain lesion presence and location with performance on measures of response bias	Outcome

	in forensic evaluation. <i>Behav Sci Law</i> 2013; 31 :779-88.	
163	McCoy TH, Jr., Yu S, Hart KL, Castro VM, Brown HE, Rosenquist JN, et al. High Throughput Phenotyping for	Population
103	Dimensional Psychopathology in Electronic Health Records. <i>Biol Psychiatry</i> 2018; 83 :997-1004.	ropulation
	https://doi.org/10.1016/j.biopsych.2018.01.011	
164	McCrea M, Iverson GL, Echemendia RJ, Makdissi M, Raftery M. Day of injury assessment of sport-related	Phenomenon of
104	concussion. <i>Br J Sports Med</i> 2013; 47 :272-84. https://doi.org/10.1136/bjsports-2013-092145	interest
165	McGilloway E, Mitchell J, Dharm-Datta S, Roberts A, Tilley H, Etherington J. The Mayo Portland Adaptability	Phenomenon of
103	Inventory-4 outcome measure is superior to UK FIM+FAM in a British military population. <i>Brain Inj</i>	interest
	, , , , , , , , , , , , , , , , , , , ,	interest
166	2016; 30 :1208-12. https://doi.org/10.1080/02699052.2016.1188215	Phenomenon of
100	McGinley A, McMillan T. The prevalence, characteristics, and impact of head injury in female prisoners: a	
1.67	systematic PRISMA review. <i>Brain Inj</i> 2019; 33 :1581-91. https://doi.org/10.1080/02699052.2019.1658223	interest
167	McKeon A. Physiological predictors of behavioral dysregulation in adults with traumatic brain injury: A novel	Phenomenon of
4.60	ecological momentary assessment method [PhD]. Pittsburgh: University of Pittsburgh; 2016.	interest
168	McKeon A, Terhorst L, Ding D, Cooper R, McCue M. Naturalistic physiological monitoring as an objective	Phenomenon of
	approach for detecting behavioral dysregulation after traumatic brain injury: A pilot study. <i>J Vocat Rehabil</i>	interest
100	2018; 49 :379-88. https://doi.org/10.3233/Jvr-180981	
169	McKinnon IG, Grubin D. Health screening of people in police custodyevaluation of current police screening	Phenomenon of
	procedures in London, UK. Eur J Public Health 2013; 23 :399-405. https://doi.org/10.1093/eurpub/cks027	interest
170	McMillan TM, Laurie M. Young adults with acquired brain injury in nursing homes in Glasgow. <i>Clin Rehabil</i>	Phenomenon of
	2004; 18 :132-8. https://doi.org/10.1191/0269215504cr712oa	interest
171	Meijer R, van Limbeek J, Kriek B, Ihnenfeldt D, Vermeulen M, de Haan R. Prognostic social factors in the	Phenomenon of
	subacute phase after a stroke for the discharge destination from the hospital stroke-unit. A systematic	interest
	review of the literature. <i>Disabil Rehabil</i> 2004; 26 :191-7. https://doi.org/10.1080/09638280310001636437	
172	Meijer R, van Limbeek J, Peusens G, Rulkens M, Dankoor K, Vermeulen M, et al. The Stroke Unit Discharge	Outcome
	Guideline, a prognostic framework for the discharge outcome from the hospital stroke unit. A prospective	
	cohort study. Clin Rehabil 2005; 19 :770-8. https://doi.org/10.1191/0269215505cr875oa	
173	Monaco F, Mazzini L, Marchetti C, Torta R, Cicolin A, Mantegazza P, et al. The structured assessment of	Phenomenon of
	depression in brain-damaged individuals: translation and validation study of the Italian version. Neurol Sci	interest
	2005; 26 :182-4. https://doi.org/10.1007/s10072-005-0459-9	
174	Myers A. Wechsler adult intelligence scale findings in mildly to severely traumatic brain-injured patients.	Phenomenon of
	Davie: Nova Southeastern University; 2011.	interest
175	Nelson NW, Hoelzle JB, Sweet JJ, Arbisi PA, Demakis GJ. Updated meta-analysis of the MMPI-2 symptom	Outcome
	validity scale (FBS): verified utility in forensic practice. <i>Clin Neuropsychol</i> 2010; 24 :701-24.	
	https://doi.org/10.1080/13854040903482863	

176	Neumann D, Babbage DR, Zupan B, Willer B. A randomized controlled trial of emotion recognition training	Phenomenon of
	after traumatic brain injury. <i>J Head Trauma Rehabil</i> 2015; 30 :E12-23.	interest
	https://doi.org/10.1097/HTR.00000000000054	
177	Newman AC, Garmoe W, Beatty P, Ziccardi M. Self-awareness of traumatically brain injured patients in the	Outcome
	acute inpatient rehabilitation setting. Brain Inj 2000;14:333-44. https://doi.org/10.1080/026990500120628	
178	Niemeier JP, Marwitz JH, Lesher K, Walker WC, Bushnik T. Gender differences in executive functions	Outcome
	following traumatic brain injury. Neuropsychol Rehabil 2007;17:293-313.	
	https://doi.org/10.1080/09602010600814729	
179	Nokleby K, Boland E, Bergersen H, Schanke AK, Farner L, Wagle J, et al. Screening for cognitive deficits after	Phenomenon of
	stroke: a comparison of three screening tools. Clin Rehabil 2008;22:1095-104.	interest
	https://doi.org/10.1177/0269215508094711	
180	Norris G, Tate RL. The Behavioural Assessment of the Dysexecutive Syndrome (BADS): Ecological, concurrent	Phenomenon of
	and construct validity. <i>Neuropsychol Rehabil</i> 2000; 10 :33-45. https://doi.org/Doi 10.1080/096020100389282	interest
181	Olai L, Omne-Ponten M, Borgquist L, Svardsudd K. Prognosis assessment in stroke patients at discharge from	Outcome
	hospital. Age Ageing 2007; 36 :184-9. https://doi.org/10.1093/ageing/afl146	
182	Olson-Madden JH, Homaifar BY, Hostetter TA, Matarazzo BB, Huggins J, Forster JE, et al. Validating the	Study type
	traumatic brain injury-4 screening measure for veterans seeking mental health treatment with psychiatric	
	inpatient and outpatient service utilization data. Arch Phys Med Rehabil 2014;95:925-9.	
	https://doi.org/10.1016/j.apmr.2014.01.008	
183	Osborne-Crowley K, McDonald S, Francis H. Development of an observational measure of social disinhibition	Phenomenon of
	after traumatic brain injury. J Clin Exp Neuropsychol 2016; 38 :341-53.	interest
	https://doi.org/10.1080/13803395.2015.1115824	
184	O'Shea LE, Thaker DK, Picchioni MM, Mason FL, Knight C, Dickens GL. Predictive validity of the HCR-20 for	Population
	violent and non-violent sexual behaviour in a secure mental health service. Crim Behav Ment Health	
	2016; 26 :366-79. https://doi.org/10.1002/cbm.1967	
185	Osterberg K, Karlson B, Orbaek P. Personality, mental distress, and risk perception in subjects with multiple	Population
	chemical sensitivity and toxic encephalopathy. <i>Scand J Psychol</i> 2002; 43 :169-75.	
	https://doi.org/10.1111/1467-9450.00283	
186	O'Sullivan M, Glorney E, Sterr A, Oddy M, Ramos SD. Traumatic brain injury and violent behavior in females:	Study type
	A systematic review. Aggress Violent Behav 2015;25:54-64. https://doi.org/10.1016/j.avb.2015.07.006	
187	Ownsworth T, Gooding K, Beadle E. Self-focused processing after severe traumatic brain injury: Relationship	Phenomenon of
	to neurocognitive functioning and mood symptoms. Br J Clin Psychol 2019; 58 :35-50.	interest
	https://doi.org/10.1111/bjc.12185	
188	Palijan TZ, Radeljak S, Kovac M, Kovacevic D. Relationship between comorbidity and violence risk	Study type

	assessment in forensic psychiatry - the implication of neuroimaging studies. <i>Psychiatria Danubina</i> 2010; 22 :253-6.	
189	Panaite V, Brown R, Henry M, Garcia A, Powell-Cope G, Vanderploeg RD, et al. Post-deployment Mental Health Screening: A Systematic Review of Current Evidence and Future Directions. Adm Policy Ment Health 2018;45:850-75. https://doi.org/10.1007/s10488-018-0869-7	Phenomenon of interest
190	Parente R, Demott E, Johnson C, Jennings P, Silver R. Measuring and manipulating subjective organization after traumatic brain injury. <i>NeuroRehabilitation</i> 2011; 29 :117-24. https://doi.org/10.3233/NRE-2011-0685	Outcome
191	Parry-Jones BL, Vaughan FL, Miles Cox W. Traumatic brain injury and substance misuse: a systematic review of prevalence and outcomes research (1994-2004). <i>Neuropsychol Rehabil</i> 2006; 16 :537-60. https://doi.org/10.1080/09602010500231875	Validity only (i.e. does not measure reliability)
192	Patrick L, Leclerc C, Perugini M. Is rehabilitation neuropsychology an evidence-based practice? Insights from a continuous quality improvement perspective. <i>Top Geriatr Rehabil</i> 2003; 19 :160-8.	Outcome
193	Paulus EM, Fabian TC, Savage SA, Zarzaur BL, Botta V, Dutton W, et al. Blunt cerebrovascular injury screening with 64-channel multidetector computed tomography: more slices finally cut it. <i>J Trauma Acute Care Surg</i> 2014; 76 :279-83; discussion 84-5. https://doi.org/10.1097/TA.000000000000101	Outcome
194	Pendlebury ST, Mariz J, Bull L, Mehta Z, Rothwell PM. Impact of different operational definitions on mild cognitive impairment rate and MMSE and MoCA performance in transient ischaemic attack and stroke. Cerebrovasc Dis 2013;36:355-62. https://doi.org/10.1159/000355496	Outcome
195	Perino C, Rago R, Cicolini A, Torta R, Monaco F. Mood and behavioural disorders following traumatic brain injury: clinical evaluation and pharmacological management. <i>Brain Inj</i> 2001; 15 :139-48. https://doi.org/10.1080/026990501458371	Phenomenon of interest
196	Peter H, Jolyon M, Carolyn T. The Weigl Colour-Form Sorting Test: a quick and easily administered bedside screen for dementia and executive dysfunction. <i>Int J Geriatr Psychiatry</i> 2007; 22 :909-15.	Phenomenon of interest
197	Petruseviciene D, Krisciunas A. Evaluation of activity and effectiveness of occupational therapy in stroke patients at the early stage of rehabilitation. <i>Medicina-Lithuania</i> 2008; 44 :216-24. https://doi.org/DOI 10.3390/medicina44030028	Phenomenon of interest
198	Pewter SM, Williams WH, Haslam C, Kay JM. Neuropsychological and psychiatric profiles in acute encephalitis in adults. <i>Neuropsychol Rehabil</i> 2007; 17 :478-505. https://doi.org/10.1080/09602010701202238	Study type
199	Pickard JD, Seeley HM, Kirker S, Maimaris C, McGlashan K, Roels E, et al. Mapping rehabilitation resources for head injury. J R Soc Med 2004; 97 :384-9. https://doi.org/10.1258/jrsm.97.8.384	Study type
200	Pignat JM, Mauron E, Johr J, Gilart de Keranflec'h C, Van De Ville D, Preti MG, et al. Outcome Prediction of Consciousness Disorders in the Acute Stage Based on a Complementary Motor Behavioural Tool. <i>PLoS One</i> 2016; 11 :e0156882. https://doi.org/10.1371/journal.pone.0156882	Outcome

201	Pinsker Donna M, Stone V, Pachana N, Greenspan S. Social Vulnerability Scale for older adults: Validation study. <i>Clinical Psychologist</i> 2006; 10 :109-19.	Outcome
202	, , ,	Dhanan an af
202	Port Ingrid GLvd. Determinants of depression in chronic stroke: a prospective cohort study. <i>Disabil Rehabil</i>	Phenomenon of
	2007; 29 :353-8.	interest
203	Prince C, Bruhns ME. Evaluation and Treatment of Mild Traumatic Brain Injury: The Role of	Study type
	Neuropsychology. Brain Sci 2017;7:17. https://doi.org/10.3390/brainsci7080105	
204	Procaccio F, Stocchetti N, Citerio G, Berardino M, Beretta L, Della Corte F, et al. Guidelines for the treatment	Unable to retrieve full-
	of adults with severe head trauma (part II). Criteria for medical treatment. J Neurosurg Sci 2000;44:11-8.	text
205	Prouteau A, Stefan A, Wiart L, Mazaux JM. The evaluation of behavioural changes in brain-injured patients:	Study type
	SOFMER recommendations for clinical practice. Ann Phys Rehabil Med 2016;59:23-30.	
	https://doi.org/10.1016/j.rehab.2015.12.002	
206	Pryor J. What do nurses do in response to their predictions of aggression? Journal of Neuroscience Nursing	Study type
	2006; 38 :177-82.	7,77
207	Qiu HC, Liu HZ, Li X, Zeng X, Zhao JZ. Insulin resistance as estimated by homeostasis model assessment	Country
207	predicts incident post-stroke depression in Chinese subjects from ischemic stroke. J Affect Disord	Country
	2018; 231 :1-7. https://doi.org/10.1016/j.jad.2018.01.023	
208	Rahman B, Alderman N, Oliver C. Use of the structured descriptive assessment to identify possible functions	Study type
200	of challenging behaviour exhibited by adults with brain injury. <i>Neuropsychol Rehabil</i> 2013; 23 :501-27.	Study type
	https://doi.org/10.1080/09602011.2013.787938	
200		Charles to an
209	Rahman B, Oliver C, Alderman N. Descriptive analysis of challenging behaviours shown by adults with	Study type
	acquired brain injury. Neuropsychol Rehabil 2010; 20 :212-38. https://doi.org/10.1080/09602010903021097	
210	Ramos SDS, Oddy M, Haye L, Goodson A. Preliminary investigation of the reliability and validity of the BIRT	Phenomenon of
	Independent Living Scale. <i>Disabil Rehabil</i> 2018; 40 :2817-23.	interest
	https://doi.org/10.1080/09638288.2017.1362594	
211	Rankin KP, Santos-Modesitt W, Kramer JH, Pavlic D, Beckman V, Miller BL. Spontaneous social behaviors	Population
	discriminate behavioral dementias from psychiatric disorders and other dementias. J Clin Psychiatry	
	2008; 69 :60-73. https://doi.org/10.4088/jcp.v69n0109	
212	Rao R. "Sadly confused': The detection of depression and dementia on medical wards. <i>Psychiatric Bulletin</i>	Population
	2001; 25 :177-9.	
213	Rao R. Outcomes from liaison psychiatry referrals for older people with alcohol use disorders in the UK.	Phenomenon of
	Mental Health and Substance Use 2013; 6 :362-8.	interest
214	Renee B, et al. The needs of people with alcohol-related brain injury (ARBI): a review of the international	Phenomenon of
	literature. Drugs Alcohol Today 2013;13:205-14.	interest
215	Richardson C, McKay A, Ponsford JL. Factors influencing self-awareness following traumatic brain injury. J	Phenomenon of
	instances of monay, it is sold the ractors in identified being wareness following traditions of an injury.	c.ioinciion oi

	Head Trauma Rehabil 2015; 30 :E43-54. https://doi.org/10.1097/HTR.0000000000000048	interest
216	Ritter AC, Wagner AK, Szaflarski JP, Brooks MM, Zafonte RD, Pugh MJ, et al. Prognostic models for predicting	Outcome
	posttraumatic seizures during acute hospitalization, and at 1 and 2 years following traumatic brain injury.	
	Epilepsia 2016; 57 :1503-14. https://doi.org/10.1111/epi.13470	
217	Ron L, Suzanne C. A comparison of psychopathy, attention deficit hyperactivity disorder, and brain	Population
	dysfunction among sex offenders. <i>Journal of Forensic Psychology Practice</i> 2010; 10 :177-200.	
218	Royle J, Whitehill Y. Development of a clinical screening tool to differentiate levels of challenging patient	Study duplicated in
	behaviourincluding commentary by Hillier S and Farrell GA. International Journal of Therapy &	included systematic
	Rehabilitation 2010; 17 :405-15.	review
219	Ruff R. Best practice guidelines for forensic neuropsychological examinations of patients with traumatic	Study type
	brain injury. <i>J Head Trauma Rehabil</i> 2009; 24 :131-40. https://doi.org/10.1097/01.HTR.0000348755.42649.e9	
220	Saini M, Tan CS, Hilal S, Dong Y, Ting E, Ikram MK, et al. Computer tomography for prediction of cognitive	Phenomenon of
	outcomes after ischemic cerebrovascular events. J Stroke Cerebrovasc Dis 2014;23:1921-7.	interest
	https://doi.org/10.1016/j.jstrokecerebrovasdis.2014.02.007	
221	Salmond CH, Sahakian BJ. Cognitive outcome in traumatic brain injury survivors. Curr Opin Crit Care	Study type
	2005; 11 :111-6. https://doi.org/10.1097/01.ccx.0000155358.31983.37	
222	Salter K, Jutai JW, Teasell R, Foley NC, Bitensky J. Issues for selection of outcome measures in stroke	Study type
	rehabilitation: ICF Body Functions. <i>Disabil Rehabil</i> 2005; 27 :191-207.	
	https://doi.org/10.1080/09638280400008537	
223	Satomi J, Ghaibeh AA, Moriguchi H, Nagahiro S. Predictability of the future development of aggressive	Outcome
	behavior of cranial dural arteriovenous fistulas based on decision tree analysis. <i>J Neurosurg</i> 2015; 123 :86-90.	
	https://doi.org/10.3171/2014.10.JNS141429	
224	Schiltz K, Witzel JG, Bausch-Holterhoff J, Bogerts B. High prevalence of brain pathology in violent prisoners: a	Population
	qualitative CT and MRI scan study. Eur Arch Psychiatry Clin Neurosci 2013; 263 :607-16.	
	https://doi.org/10.1007/s00406-013-0403-6	
225	Schumacher R, Walder B, Delhumeau C, Muri RM. Predictors of inpatient (neuro)rehabilitation after acute	Phenomenon of
	care of severe traumatic brain injury: An epidemiological study. <i>Brain Inj</i> 2016; 30 :1186-93.	interest
	https://doi.org/10.1080/02699052.2016.1183821	
226	Seel RT, Macciocchi S, Kreutzer JS. Clinical considerations for the diagnosis of major depression after	Study type
	moderate to severe TBI. J Head Trauma Rehabil 2010; 25 :99-112.	
	https://doi.org/10.1097/HTR.0b013e3181ce3966	
227	Senior G, Douglas L. Misconceptions and misuse of the MMPI-2 in assessing personal injury claimants.	Study type
	NeuroRehabilitation 2001; 16 :203-13.	
228	Seniow J, Mroziak B, Czlonkowska A, Jedryka-Goral A. Self-rated emotional functioning of patients with	Population

	neurological or asymptomatic form of Wilson's disease. <i>Clin Neuropsychol</i> 2003; 17 :367-73. https://doi.org/10.1076/clin.17.3.367.18085	
229	Sherer M, Yablon SA, Nick TG. Psychotic symptoms as manifestations of the posttraumatic confusional state: prevalence, risk factors, and association with outcome. <i>J Head Trauma Rehabil</i> 2014; 29 :E11-8. https://doi.org/10.1097/HTR.0b013e318287f894	Unable to retrieve full- text
230	Shirley AT, Nadina BL. Factors relating to depression after stroke. <i>Br J Clin Psychol</i> 2006; 45 :49-61.	Phenomenon of interest
231	Siegert RJ, Jackson DM, Turner-Stokes L. The Needs and Provision Complexity Scale: a first psychometric analysis using multicentre data. <i>Clin Rehabil</i> 2014; 28 :687-95.	Population
232	Silk-Eglit GM, Miele AS, Stenclik JH, Lynch JK, McCaffrey RJ. Evaluation of the Generalizability of the Number of Abnormal Scores and the Overall Test Battery Mean as Measures of Performance Validity to a Different Test Battery. <i>Appl Neuropsychol Adult</i> 2015; 22 :399-406. https://doi.org/10.1080/23279095.2014.949719	Outcome
233	Silva S, Sousa D, Rezende V, Teixeira A, Peixoto B. The psychometric properties of the Portuguese version of the Stroke Quality of Life Scale. <i>Adv Clin Exp Med</i> 2012; 21 :781-90.	Phenomenon of interest
234	Simpson GK, Sabaz M, Daher M, Gordon R, Strettles B. Challenging behaviours, co-morbidities, service utilisation and service access among community-dwelling adults with severe traumatic brain injury: A multicentre study. <i>Brain Impairment</i> 2014;15:28-42.	Outcome
235	Singh R, Mason S, Lecky F, Dawson J. Prevalence of depression after TBI in a prospective cohort: The SHEFBIT study. <i>Brain Inj</i> 2018; 32 :84-90. https://doi.org/10.1080/02699052.2017.1376756	Phenomenon of interest
236	Singh R, Mason S, Lecky F, Dawson J. Comparison of early and late depression after TBI; (the SHEFBIT study). Brain Inj 2019;33:584-91. https://doi.org/10.1080/02699052.2019.1566837	Phenomenon of interest
237	Singh R, Venkateshwara G, Nair KP, Khan M, Saad R. Agitation after traumatic brain injury and predictors of outcome. <i>Brain Inj</i> 2014; 28 :336-40. https://doi.org/10.3109/02699052.2013.873142	Phenomenon of interest
238	Smeets SM, Vink M, Ponds RW, Winkens I, van Heugten CM. Changes in impaired self-awareness after acquired brain injury in patients following intensive neuropsychological rehabilitation. <i>Neuropsychol Rehabil</i> 2017; 27 :116-32. https://doi.org/10.1080/09602011.2015.1077144	Phenomenon of interest
239	Soendergaard PL, Siert L, Poulsen I, Wood RL, Norup A. Measuring Neurobehavioral Disabilities Among Severe Brain Injury Survivors: Reports of Survivors and Proxies in the Chronic Phase. <i>Front Neurol</i> 2019; 10 :51. https://doi.org/10.3389/fneur.2019.00051	Phenomenon of interest
240	Stefan A, Mathe JF, group S. What are the disruptive symptoms of behavioral disorders after traumatic brain injury? A systematic review leading to recommendations for good practices. <i>Ann Phys Rehabil Med</i> 2016; 59 :5-17. https://doi.org/10.1016/j.rehab.2015.11.002	Study type
241	Stein MB, Jain S, Giacino JT, Levin H, Dikmen S, Nelson LD, et al. Risk of Posttraumatic Stress Disorder and Major Depression in Civilian Patients After Mild Traumatic Brain Injury: A TRACK-TBI Study. <i>JAMA Psychiatry</i>	Phenomenon of interest

	2019; 76 :249-58. https://doi.org/10.1001/jamapsychiatry.2018.4288	
242	Stewart CA, Enders FT, Schneider N, Felmlee-Devine D, Kamath PS, Smith GE. Development of a three-factor	Phenomenon of
	neuropsychological approach for detecting minimal hepatic encephalopathy. <i>Liver Int</i> 2010; 30 :841-9.	interest
	https://doi.org/10.1111/j.1478-3231.2010.02246.x	
243	Stulemeijer M, Vos PE, Bleijenberg G, van der Werf SP. Cognitive complaints after mild traumatic brain	Phenomenon of
	injury: things are not always what they seem. J Psychosom Res 2007;63:637-45.	interest
	https://doi.org/10.1016/j.jpsychores.2007.06.023	
244	Suter-Riederer S, Mahrer Imhof R, Kesselring J, Imhof L. "Care on the ground floor" for agitated people after	Non-English language
	traumatic brain injury. Neurologie und Rehabilitation 2008; 14 :70-8.	study
245	Svensson S, Sonn U, Sunnerhagen KS. Reliability and validity of the Northwick Park Dependency Score	Outcome
	(NPDS) Swedish version 6.0. Clin Rehabil 2005;19:419-25. https://doi.org/10.1191/0269215505cr808oa	
246	Swaine B, Cullen N, Messier F, Bayley M, Lavoie A, Marshall S, et al. Post-acute care referral and inpatient	Outcome
	rehabilitation admission criteria for persons with brain injury across two Canadian provinces. Disabil Rehabil	
	2018; 40 :697-704. https://doi.org/10.1080/09638288.2016.1262911	
247	Tang CZ, Zhang YL, Wang WS, Li WG, Shi JP. Elevated Serum Levels of Neopterin at Admission Predicts	Country
	Depression After Acute Ischemic Stroke: a 6-Month Follow-Up Study. <i>Mol Neurobiol</i> 2016; 53 :3194-204.	
	https://doi.org/10.1007/s12035-015-9220-4	
248	Tang WK, Ungvari GS, Chiu HF, Sze KH, Yu AC, Leung TL. Screening post-stroke depression in Chinese older	Phenomenon of
	adults using the hospital anxiety and depression scale. Aging Ment Health 2004;8:397-9.	interest
	https://doi.org/10.1080/13607860410001725027	
249	Taylor Brent C, Campbell Emily H, Nugent S, Bidelspach Douglas E, Kehle-Forbes Shannon M, Scholten J, et	Study type
	al. Three year trends in veterans health administration utilization and costs after traumatic brain injury	
	screening among veterans with mild traumatic brain injury. <i>J Neurotrauma</i> 2017; 34 :2567-74.	
250	Tellier A, Marshall SC, Wilson KG, Smith A, Perugini M, Stiell IG. The heterogeneity of mild traumatic brain	Phenomenon of
	injury: Where do we stand? Brain Inj 2009; 23 :879-87.	interest
251	Terrill AL, Schwartz JK, Belagaje S. Understanding Mental Health Needs After Mild Stroke. <i>Arch Phys Med</i>	Study type
	Rehabil 2019; 100 :1003-8. https://doi.org/10.1016/j.apmr.2018.12.017	
252	Thomas A. Thinking it Through. <i>OT Practice</i> 2017:14-7.	Study type
253	Todd KL. A neuropsychological study of traumatic brain injury among a Canadian sample of male federal	Phenomenon of
	offenders. Regina: University of Regina; 2011.	interest
254	Tornas S, Stubberud J, Solbakk AK, Evans J, Schanke AK, Lovstad M. Moderators, mediators and nonspecific	Phenomenon of
	predictors of outcome after cognitive rehabilitation of executive functions in a randomised controlled trial.	interest
	Neuropsychol Rehabil 2019; 29 :844-65. https://doi.org/10.1080/09602011.2017.1338587	
255	Toyokura M, Yamashita K, Hayashi T, Nishimura Y, Sawatari M, Kikui R, et al. A newly developed assessment	Phenomenon of

scale for attentional disturbance based on behavioral problems: Behavioral Assessment of Attentional	interest
Disturbance (BAAD). Tokai J Exp Clin Med 2006; 31 :29-33.	
experience. Asian J Neurosurg 2010; 5 :24-31.	Study type
Tucha O, Smely C, Preier M, Lange KW. Cognitive deficits before treatment among patients with brain tumors. <i>Neurosurgery</i> 2000; 47 :324-33; discussion 33-4. https://doi.org/10.1097/00006123-200008000-00011	Phenomenon of interest
Tuninger E, Levander S. Neuropsychological impairment in patients treated with depot neuroleptics: a longitudinal study. <i>Acta Psychiatr Scand Suppl</i> 2001; 104 :75-80. https://doi.org/10.1034/j.1600-0447.2001.00109.x	Population
Turner-Stokes L. The Needs and Provision Complexity Scale: a multicentre prospective cohort analysis of met and unmet needs and their cost implications for patients with complex neurological disability. <i>BMJ Open</i> 2013; 3 .	Population
Turner-Stokes L, Kalmus M, Hirani D, Clegg F. The Depression Intensity Scale Circles (DISCs): a first evaluation of a simple assessment tool for depression in the context of brain injury. <i>J Neurol Neurosurg Psychiatry</i> 2005; 76 :1273-8. https://doi.org/10.1136/jnnp.2004.050096	Phenomenon of interest
Turner-Stokes L, Siegert RJ. A comprehensive psychometric evaluation of the UK FIM + FAM. <i>Disabil Rehabil</i> 2013; 35 :1885-95. https://doi.org/10.3109/09638288.2013.766271	Phenomenon of interest
Turner-Stokes L, Thu A, Williams H, Casey R, Rose H, Siegert RJ. The Neurological Impairment Scale: reliability and validity as a predictor of functional outcome in neurorehabilitation. <i>Disabil Rehabil</i> 2014; 36 :23-31. https://doi.org/10.3109/09638288.2013.775360	Outcome
Turner-Stokes L, Williams H, Rose H, Harris S, Jackson D. Deriving a Barthel Index from the Northwick Park Dependency Scale and the Functional Independence Measure: are they equivalent? <i>Clin Rehabil</i> 2010; 24 :1121-6.	Population
Vanderploeg RD, Groer S, Belanger HG. Initial developmental process of a VA semistructured clinical interview for TBI identification. <i>J Rehabil Res Dev</i> 2012; 49 :545-56. https://doi.org/10.1682/jrrd.2011.04.0069	Study type
Vitacco MJ, Van Rybroek GJ, Rogstad JE, Yahr LE, Tomony JD, Saewert E. Predicting short-term institutional aggression in forensic patients: a multi-trait method for understanding subtypes of aggression. <i>Law Hum Behav</i> 2009; 33 :308-19. https://doi.org/10.1007/s10979-008-9155-7	Population
Volbrecht ME, Meyers JE, Kaster-Bundgaard J. Neuropsychological outcome of head injury using a short battery. <i>Arch Clin Neuropsychol</i> 2000; 15 :251-65. https://doi.org/Doi 10.1016/S0887-6177(99)00016-5	Phenomenon of interest
Von Hilsheimer G. A psychobiological study of delinquents: Humanistic Psychology Institute; 2003.	Population
	Disturbance (BAAD). Tokai J Exp Clin Med 2006;31:29-33. Truelle JL, Wild KV, Onillon M, Montreuil M. Social reintegration of traumatic brain-injured: the French experience. Asian J Neurosurg 2010;5:24-31. Tucha O, Smely C, Preier M, Lange KW. Cognitive deficits before treatment among patients with brain tumors. Neurosurgery 2000;47:324-33; discussion 33-4. https://doi.org/10.1097/00006123-200008000-00011 Tuninger E, Levander S. Neuropsychological impairment in patients treated with depot neuroleptics: a longitudinal study. Acta Psychiatr Scand Suppl 2001;104:75-80. https://doi.org/10.1034/j.1600-0447.2001.00109.x Turner-Stokes L. The Needs and Provision Complexity Scale: a multicentre prospective cohort analysis of met and unmet needs and their cost implications for patients with complex neurological disability. BMJ Open 2013;3. Turner-Stokes L, Kalmus M, Hirani D, Clegg F. The Depression Intensity Scale Circles (DISCs): a first evaluation of a simple assessment tool for depression in the context of brain injury. J Neurol Neurosurg Psychiatry 2005;76:1273-8. https://doi.org/10.1136/jnnp.2004.050096 Turner-Stokes L, Siegert RJ. A comprehensive psychometric evaluation of the UK FIM + FAM. Disabil Rehabil 2013;35:1885-95. https://doi.org/10.3109/09638288.2013.766271 Turner-Stokes L, Thu A, Williams H, Casey R, Rose H, Siegert RJ. The Neurological Impairment Scale: reliability and validity as a predictor of functional outcome in neurorehabilitation. Disabil Rehabil 2014;36:23-31. https://doi.org/10.3109/09638288.2013.775360 Turner-Stokes L, Williams H, Rose H, Harris S, Jackson D. Deriving a Barthel Index from the Northwick Park Dependency Scale and the Functional Independence Measure: are they equivalent? Clin Rehabil 2010;24:1121-6. Vanderploeg RD, Groer S, Belanger HG. Initial developmental process of a VA semistructured clinical interview for TBI identification. J Rehabil Res Dev 2012;49:545-56. https://doi.org/10.1682/jrrd.2011.04.0069 Vitacco MJ, Van Rybroek GJ, Rogstad JE, Yahr LE, Tomony JD, Saewe

	in older people: Alcohol Research UK; 2013.	
269	Walker WC, Stromberg KA, Marwitz JH, Sima AP, Agyemang AA, Graham KM, et al. Predicting Long-Term	Phenomenon of
	Global Outcome after Traumatic Brain Injury: Development of a Practical Prognostic Tool Using the	interest
	Traumatic Brain Injury Model Systems National Database. <i>J Neurotrauma</i> 2018; 35 :1587-95.	
	https://doi.org/10.1089/neu.2017.5359	
270	Walters GD, Duncan SA. Performance-verbal discrepancies and facets of psychopathy: assessing the	Population
	relationship between WAIS–R/III summary IQs/index scores and PCL–R facet scores. J Crim Psychol	
	2018; 8 :234-46. https://doi.org/10.1108/jcp-12-2017-0045	
271	Wasserman L, Shaw T, Vu M, Ko C, Bollegala D, Bhalerao S. An overview of traumatic brain injury and	Study type
	suicide. <i>Brain Inj</i> 2008; 22 :811-9. https://doi.org/10.1080/02699050802372166	
272	Willemse-van Son AH, Ribbers GM, Verhagen AP, Stam HJ. Prognostic factors of long-term functioning and	Phenomenon of
	productivity after traumatic brain injury: a systematic review of prospective cohort studies. Clin Rehabil	interest
	2007; 21 :1024-37. https://doi.org/10.1177/0269215507077603	
273	Williams WH, Evans JJ, Needham P, Wilson BA. Neurological, cognitive and attributional predictors of	Phenomenon of
	posttraumatic stress symptoms after traumatic brain injury. <i>J Trauma Stress</i> 2002; 15 :397-400.	interest
	https://doi.org/10.1023/A:1020185325026	
274	Winocur G, Palmer H, Stuss DT, Alexander MP, Craik FI, Levine B, et al. Cognitive rehabilitation in clinical	Study type
	neuropsychology. <i>Brain Cogn</i> 2000; 42 :120-3. https://doi.org/10.1006/brcg.1999.1179	
275	Wong GK, Lam SW, Chan SS, Lai M, Tse PP, Mok V, et al. Neuropsychiatric disturbance after aneurysmal	Phenomenon of
	subarachnoid hemorrhage. <i>J Clin Neurosci</i> 2014; 21 :1695-8. https://doi.org/10.1016/j.jocn.2014.02.018	interest
276	Wood A. Rey-Osterrieth Figure Copy size and cognitive functioning in veterans. Massachusetts: William	Phenomenon of
	James College; 2019.	interest
277	Wygant DB, Ben-Porath YS, Arbisi PA, Berry DT, Freeman DB, Heilbronner RL. Examination of the MMPI-2	Outcome
	restructured form (MMPI-2-RF) validity scales in civil forensic settings: findings from simulation and known	
	group samples. Arch Clin Neuropsychol 2009;24:671-80. https://doi.org/10.1093/arclin/acp073	
278	Xu KY, Rossi KC, Kim AM, Jette N, Yoo JY, Hung K, et al. Risk of readmission for suicide attempt after epilepsy	Phenomenon of
	hospitalization. <i>Epilepsy Behav</i> 2018; 83 :124-30. https://doi.org/10.1016/j.yebeh.2018.03.037	interest
279	Yamaki T, Suzuki K, Sudo Y, Niitsu T, Okai M, Oka N, et al. Association between uncooperativeness and the	Phenomenon of
	glucose metabolism of patients with chronic behavioral disorders after severe traumatic brain injury: a	interest
	cross-sectional retrospective study. <i>Biopsychosoc Med</i> 2018; 12 :6. https://doi.org/10.1186/s13030-018-	
	0125-0	
280	Yantz CL, Gavett BE, Lynch JK, McCaffrey RJ. Potential for interpretation disparities of Halstead-Reitan	Phenomenon of
	neuropsychological battery performances in a litigating sample. Arch Clin Neuropsychol 2006;21:809-17.	interest
	https://doi.org/10.1016/j.acn.2006.09.001	

28	Yu KH, Cho SJ, Oh MS, Jung S, Lee JH, Shin JH, et al. Cognitive impairment evaluated with Vascular Cognitive	Population
	Impairment Harmonization Standards in a multicenter prospective stroke cohort in Korea. Stroke	
	2013; 44 :786-8. https://doi.org/10.1161/STROKEAHA.112.668343	