

**Evaluating graduated progress towards, and impacts of, the implementation
of indoor smokefree prison facilities in Scotland**

The Tobacco in Prisons (TIPs) study. Version 2.5

**Kate Hunt, Helen Sweeting, Ashley Brown, Sean Semple, Douglas Eadie on behalf of the TIPs
research team**

09/04/19

© University of Glasgow and University of Stirling

Scientific title: Evaluating graduated progress towards, and impacts of, the implementation of indoor smokefree prison facilities in Scotland: the Tobacco in Prisons study (TIPs)

Lay title: The Tobacco in Prisons study (TIPs)



Protocol date: 9/04/19

Protocol version: 2.5

Principal Investigator: Professor Kate Hunt*, Institute for Social Marketing, University of Stirling.

Co-Investigators: Dr Helen Sweeting¹, Dr Sean Semple², Professor Linda Bauld³, Douglas Eadie³, Professor Jill Pell¹, Professor Alastair Leyland¹, Dr Peter Craig¹, Dr Kathleen Boyd¹, Dr Evangelia Demou¹, Dr Phillip Conaglen⁴, Dr Emily Tweed¹

Researchers: Ashley Brown^{1*}; Dr Richard Purves³

* University of Glasgow (until Feb 2018), University of Stirling (from Mar 2018);

¹University of Glasgow; ²University of Aberdeen (until Dec 2017; University of Stirling from Jan 2018); ³University of Stirling; ⁴NHS Lothian

Funding reference: NIHR-PHR-15/55/44

Table of Contents

1. Background	3
1.1 Existing research	3
Description of the problem	3
Tobacco and smoking within prison culture	5
Evidence and evidence gaps in introducing smokefree prison policies	5
Type chapter title (level 3)	6
Type chapter title (level 3)	6
1.2 Risks and benefits	6
1.3 Rationale for current study	7
2. Research Objectives	7
3. Research Design	8
3.1 WP1 – Scoping the international landscape	8
3.2 WP2 – Evaluating exposure and outcomes	8
3.3 WP3 & 4 – Surveys and qualitative research with prison staff and prisoners	11
3.4 WP5 – Cessation services :experiences, provision and outcomes	13
4. Study Population	14
5. Socio-economic position and inequalities	14
6. Planned Interventions	15
7. Proposed outcome measures	15
8. Assessment and follow-up	15
9. Proposed sample size	15
10. Analysis	16
11. Ethical arrangements	17
12. Research Governance	18
13. Project timetable and milestones	19
14. Expertise	20
15. Partner Collaboration	20
16. Funding	20
17. References	21

Project title. Evaluating graduated progress towards, and impacts of, the implementation of indoor smokefree prison facilities in Scotland: the Tobacco in Prisons (TIPs) study

Background to revised protocol, March 2018

On 17 July 2017, the Scottish Prison Service (SPS) announced that Scotland's prisons would become smoke free in November 2018, informed by the measurement of exposures to secondhand smoke (SHS) conducted within Scotland's prisons as part of Phase 1 of the Tobacco in Prisons (TIPs) study (Semple, Sweeting et al. 2017). This necessitated revisions to Version 1.4 of the Protocol to take account of the known implementation date, 30 November 2018. In addition, our experience of conducting Phase 1 data collection led to other changes to the protocol. In particular, the collection and analysis of three measures of SHS exposure data demonstrated that quantification of SHS concentrations in prisons using Dylos DC1700 particle counters was highly feasible and entailed relatively little disruption to the prison regime. Importantly the phase 1 data showed that there were high levels of correlation between Dylos measurements and airborne nicotine concentrations. By contrast, collection of saliva samples to assess nicotine/SHS intake among non-smoking prison staff was very onerous for both research and prison staff and made additional demands on the regime at crucial times of the working day (as staff came on and off shift). Given this and the impossibility of linking saliva measurements of cotinine (for individual staff members) pre and post implementation and our expectation based on Phase 1 fieldwork that the proportion of staff volunteering to provide a saliva sample will be low following implementation of a complete ban on smoking within Scotland's prisons, we considered that the potential scientific value of Phase 3 saliva measurements was not high, and that resources could be more effectively redeployed.

Principal changes between our original Protocol (Version 1.4), and the current version (Version 2.5) thus include: i) an extended time frame to 30 May 2020 to take account of the known timing of implementation of smokefree prisons in Scotland; ii) removal of saliva sampling in non-smoking staff in Phase 3, following our experience in Phase 1 fieldwork and the high correlations between measures of SHS (Semple, Sweeting et al. 2017); iii) reconfiguration of qualitative data collection for WP3-5, to take account of expected changes in the lead up to implementation in November 2018, to examine use and experience of smoking cessation services post-implementation and staff views on what influences successful/less successful implementation (via focus groups with staff in Phase 3 post-implementation); iv) an increase in the number of 'case-study' prisons in response to feedback from Prison Governors, SPS and our SPS Research Advisory Group; and v) an extension of WP1 fieldwork into Phase 2 to allow for additional ethical application procedures in various countries and the inclusion of more interviews with personnel in prisons in England and Wales given views expressed by SPS staff in Phase 1 focus groups that they were most interested in the experiences of colleagues in the UK (the four Welsh prisons and initially four English pilot prisons implemented smokefree policies in 2016, with subsequent phased roll-out across the service in England).

1. Background

Smoking bans in public places decrease exposure to SHS (Semple, Macalman et al. 2007) with direct health benefit (Pell, Haw et al. 2008), but national smoking bans are very heterogeneous, especially in custodial settings (Hartwig, Stover et al. 2008). In the UK, prisons (as both staff's workplace and prisoners' 'home') have had partial exemption from smokefree legislation, and have been one of the few institutions in which smoking remained normative prior to the introduction of smoking bans. Whilst rates have fallen markedly in the general population, except in the most deprived (Hiscock, Bauld et al. 2012, Scottish Government 2013), across Europe 64-88% of prisoners are smokers (Hartwig, Stover et al. 2008). Internationally, smoking in prison staff is also thought to be high but evidence is sparse, with only one study (Carpenter, Hughes et al. 2001) identified in a recent review.

Prisoners' high levels of smoking can lead to high levels of SHS within prisons (Semple, Ibrahim et al. 2015), despite the fact that smoking is already subject to some restrictions in these settings. (In Scotland, prior to new policy to be introduced on 30.11.18, prisoners are only permitted to smoke in their cells or during outdoor recreation.) As a result, prisoners' smoking poses a threat to the health of both smoking and non-smoking prisoners and staff, particularly those opening or entering prisoners' cells. Exposure to SHS in prisons is attracting increasing concern both internationally and within the UK. Several countries have implemented total smoke-free policies (i.e. all indoor and outdoor areas) across their prison estate. In the UK,

comprehensive smoke-free policies were adopted by Broadmoor Secure Hospital in 2007 and by the State Hospital in Scotland in 2011. In October 2015, following five years of preparation, it was announced that a comprehensive smoke-free policy would be implemented in all Welsh prisons from January 2016 and at four early adopter sites in England from March 2016.

The Scottish Government's current Tobacco Strategy states that "[C]reating a smoke-free prison service should be seen as a key step on our journey to creating a smoke-free Scotland" (Scottish Government 2013), p26). The Scottish Government, NHS Boards and the Scottish Prison Service (SPS) are considering options on how best to proceed towards a smoke-free prison service, as the issues around implementing/delaying the introduction of such a policy are increasingly debated in the public domain. Following a recommendation from the Ministerial Group on Offender Reintegration, a NHS specification was launched in June 2015 to harmonise smoking cessation services across Scottish prisons (NHS Health Scotland 2015), and a joint action plan concerning smoking in prisons was submitted to ministers in early 2016. Although the exact timing of any policy change in Scotland was yet to be decided prior to Phase 1 of the TIPs study, those concerned about SHS exposures were thought likely to press for the shortest feasible timescale, whilst acknowledging the need for sufficient time and resources to make any change in policy effective, supported and enforceable. Best evidence to date, following New Zealand's experience (Gautam, Glover et al. 2011, Collinson, Wilson et al. 2012) suggests that a minimum period of around one year is needed, but some countries may require longer to mobilise resources for adequate smoking cessation and support.

1.1. Existing research

Description of the problem: high smoking rates in prisoners and exposure of prisoners and prison workers to SHS

The high prevalence of smoking in prisons has been described as "one of the most pernicious public health problems affecting prisons .. all too often .. ignored [in] community based tobacco control policies" (Binswanger, Carson et al. 2014) (p291). A European Commission (EC) report cites the prevalence of smoking in prisoners as "64 to 88%" (Hartwig, Stover et al. 2008), and identified only one study (Carpenter, Hughes et al. 2001) of smoking prevalence among prison staff. Public Health England suggest smoking rates in prisoners are approximately four times higher than in the general population (Public Health England 2015). The national biannual survey of prisoners in Scotland conducted in 2013 reported that 74% smoke¹, three times the national average, with little evidence of reductions in smoking as seen in the general population. These "staggeringly high" (Binswanger, Carson et al. 2014) rates partially reflect the fact that prisoners are disproportionately drawn from deprived and socially excluded communities which have high smoking rates (Hiscock, Bauld et al. 2012). However, imprisonment can also lead to uptake of, or increased, smoking (Donahue 2009, GlobalSmokefree Partnership 2009, MacDonald, Angus et al. 2010, Baybutt, Ritter et al. 2014, Australian Institute of Health and Welfare 2015, Public Health England 2015), but it can present the opportunity for interventions to improve health too.

The persistence of high levels of smoking poses a threat to prisoner health and a challenge to aspirations to achieve smoke-free societies and decrease inequalities in health (Scottish Government 2013). Prisoners' high levels of smoking can lead to high levels of SHS within prisons², despite existing restrictions on smoking in prisons. (In Scotland, for example, Prison Rule 36 was amended in 2006 so that prisoners are only permitted to smoke in their cells or during outdoor recreation.) WHO considers that "there is no safe level of exposure to second-hand tobacco smoke", citing evidence that SHS increases the risk of CHD, lung cancer, breast cancer, and respiratory symptoms and illnesses in adults, and, for those exposed during pregnancy, of low birth weight and preterm delivery (World Health Organisation 2007). Both prisoners and staff (particularly those entering or opening up prisoners' cells) are potentially at risk. Binswanger et al note that "People in prisons live and work in confined indoor spaces that can be crowded and poorly ventilated .. [with] limited movement options and restricted opportunities for outside air to avoid smoke exposure even if they do not smoke" (Public Health England 2015) (p2). Prisons' dual status as 'workplace' for staff and 'home' for prisoners was noted when

¹ The equivalent figure from the 2015 Prisoner Survey is 72%

²

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/469654/SHS_in_Prison_Final_Report_minus_Appendix.pdf

indoor smoke-free legislation was introduced in the UK and elsewhere, and national smoking bans have been described as “very heterogeneous, especially when it comes to the custodial settings” (Hartwig, Stover et al. 2008) (p9).

Exposure to SHS in prisons is attracting increasing concern. A report published in October 2015³ based on data from six prisons in England provided major new evidence. It showed that peaks in SHS in cells where smoking takes place can be considerable; measurements over 24h periods suggested average levels in smokers’ cells that are frequently in the range of 100-200 $\mu\text{g}/\text{m}^3$ of fine Particulate Matter less than 2.5 microns in size ($\text{PM}_{2.5}$ ⁴) (consistent with the upper 10% of levels measured in smokers’ homes). Prison staff reported that about half their time involves exposure to SHS (see footnote 3). Objective measurement of personal exposures indicated that staff spent about one-sixth of their shift in areas where SHS-derived $\text{PM}_{2.5}$ concentrations exceed $25\mu\text{g}/\text{m}^3$ (the 24h guidance level from the WHO for indoor $\text{PM}_{2.5}$ levels) and biological markers (salivary cotinine) validated SHS exposure (see footnote 3). The report concludes that “There is no safe level of exposure to SHS ..[and] it is now widely considered prudent that workers’ exposure to SHS .. should be limited as far as reasonably practicable .. [T]he National Offender and Management Service (NOMS) should consider implementing measures to further reduce or eliminate SHS within the prison estate” (p38). Others have highlighted an “urgent need to ensure the same protection [from SHS] for offenders, [prison] staff and visitors as afforded to the general population” (Public Health England 2015) (p8).

Thus momentum is building throughout the UK, as elsewhere (e.g. Australia (Australian Institute of Health and Welfare 2015)), to increase tobacco control in prisons, both to improve the health of staff and prisoners and as a means of addressing inequalities in health (Butler, Richmond et al. 2007, Scottish Government 2013). The Scottish Government’s Tobacco Strategy states that “consideration should be given to ensuring all prisoners and staff are protected from the harms caused by second-hand smoke .. [C]reating a smoke-free prison service should be seen as a key step on our journey to creating a smoke-free Scotland” (Scottish Government 2013) (p26). The strategy set an action to “work in partnership with the Scottish Prison Service and local NHS Boards to have plans in place by 2015 that set out how indoor smoke-free prison facilities will be delivered” (p26). The English Government’s *Healthy Lives, Healthy People, A Tobacco Control Plan for England* similarly includes an action to “continue progress to reduce secondhand smoke in prisons” and Public Health England argued in 2015 that, “reducing smoking should be given the highest priority across the CJS [Criminal Justice System] and comprehensive nicotine dependence treatment (cessation and/or harm reduction) should be delivered to all smokers in the CJS” (Public Health England 2015) (p4).

However, it is widely recognised, given high smoking amongst prisoners (Hartwig, Stover et al. 2008, Australian Institute of Health and Welfare 2015), that introducing smoke-free legislation in this setting presents particular and considerable challenges, including concerns about prisoner unrest (McInerney 2015, Shaw 2015). In media coverage of the announcement that a smoke-free policy would be implemented in all four prisons in Wales from January 2016 and piloted in four prisons in England in March 2016 as part of a “phased roll out that will eventually see all jails in England and Wales go smoke-free”, Prisons Minister Andrew Selous is quoted as saying this is a “necessary” but “difficult thing to do” given the “unique environment”. The “operational safety and security of our prisons”, he continues, will “always be our top priority .. We have no plans to move to smoke-free prisons overnight and will only do so in a phased way that takes into account operational resilience and readiness of each prison”. Similar concerns are shared by the Scottish Government and Prison Service.

Tobacco and smoking within prison culture

The challenges underlying the introduction of smoke-free prisons are underpinned by: the social and cultural meanings of tobacco products and smoking in day-to-day prison life; and the complexities of the management of nicotine addiction and the regulation and delivery of nicotine replacement, as with other medications, within the prison environment (Lankenau 2001, de Viggiani 2007, Richmond, Butler et al. 2009, Richmond,

3

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/469654/SHS_in_Prisons_Final_Report_minus_Appendix.pdf

⁴ $\text{PM}_{2.5}$ is an internationally recognised marker of SHS concentrations that has been previously used to quantify SHS concentrations in workplace, domestic and car settings [Hyland, A., M. Travers, C. Dresler, C. Higbee and K. Cummings (2008). "A 32 country comparison of tobacco smoke derived particle levels in indoor public places." *Tobacco Control* 17(3): 159-165.]

Butler et al. 2012). Butler described tobacco smoking as “an integral part of prison life and an established part of the prison culture”. He notes that it serves a range of functions in prison “as a surrogate currency, a means of social control, as a symbol of freedom in a group with few rights and privileges, a stress reliever and as a social lubricant” (Butler, Richmond et al. 2007) (p291). In addition, prisoners frequently cite boredom and coping with stress to explain a stronger felt need to smoke while in prison (Enggist, Moller et al. 2014). Cigarettes can thus represent a means of dealing with the challenge of ‘killing time’ as well as representing “both property and privilege” (Taylor, Ogden et al. 2012). Tobacco-based products offer prisoners “cultural capital to buy and exchange items; favours and protection” (Taylor, Ogden et al. 2012), as a *de facto* (Beck 1995) or alternative (Richmond, Butler et al. 2009) currency. Cigarettes can also give “inmates the opportunity to construct their own identity through material possessions available to them... Perhaps the decision to smoke (or not to smoke) is indicative of one of the last functions that the inmate has control over” (Taylor, Ogden et al. 2012). Ireland notes how, in the context of racketeering, extortion, bullying and ‘baroning’, tobacco can be used to “alleviate immediate danger” and “as a measure of self-protection, insulating against physical attacks of acts of violence”; without nicotine, he suggests, some other “substance or artefact is likely to act as the currency of power and influence”.

Hence, for the implementation of any smoke-free prison policy to be successful and enforceable, it is crucial to understand the social and cultural value of tobacco and smoking that prevails in prisons prior to, and following, any change in legislation, and whether other harmful substances or practices replace the use of tobacco. To date, however, qualitative research on the meaning of smoking in prisons, and particularly on how this changes in the context of increased restrictions, is sparse.

Evidence and evidence gaps in introducing smoke-free prison policies

Although restrictions on smoking in many parts of UK prisons have been in place for some time, there are precedents for ‘total’ smoking bans. New Zealand introduced the first country-wide ban on smoking within prisons and outside areas in 2011 (Gautam, Glover et al. 2011, Collinson, Wilson et al. 2012). It was recognised there that a ban would need to be supported by “comprehensive cessation support for all inmates” to maximise health gain and the likelihood that abstinence is maintained post-release, and to mitigate negative unintended consequences of imposing a ban (Gautam, Glover et al. 2011). Preparatory work included: a detailed communications strategy; offers of psychological and pharmacological support for prisoners and staff; and the establishment of voluntary smoke-free units prior to the ban. Following the ban, negative incidents included: increases in contraband tobacco and a doubling of the black-market price; smoking of alternative substances (e.g. nicotine patches); and increased reported violence between prisoners in one prison. However, no major incidents were reported, and SHS (PM_{2.5}) reduced by 63% and arson-related incidents decreased. By contrast, research in Switzerland, the first to measure exposure to SHS in Europe before and after introduction of a partial smoking ban in prisons, did not show a reduction in particulate matter (PM₁₀⁵), although a reduction in nicotine in the ambient air was detected (from 7.0µg/m³ two months before tobacco restrictions were enacted in 2009 to 2.1µg/m³ 11 months after the ban was introduced). The authors concluded that communication and movement between cells and shared rooms leads to “inevitable” air contamination and that the “suboptimal decline in particulate matter” and the residual nicotine indicated that the ban may not have been “fully respected or enforced” (Ritter, Huynh et al. 2012). This again emphasises the need to understand how increased restrictions can be introduced in the most acceptable, enforceable and therefore effective ways. A further gap in the evidence, given the recent rapid increase in the use of electronic cigarettes (e-cigarettes) in the general population, is whether (and if so how) e-cigarettes may either facilitate the introduction of smoke-free policies in prisons or pose particular risks in this environment.

Evidence from jurisdictions which have adopted total smoke-free policies suggests that the factors required for success include: “comprehensive levels of planning; a long lead-in period; clear communication and consultation with both staff and prisoners; clear instruction and guidance from management; holistic cessation support for staff and prisoners; the provision of alternative activities; and comprehensive staff training and support” (Offender Health Research Network 2014), p28). However, there is little published research in this area.

⁵ As noted earlier, it is more usual to measure PM_{2.5}

This context provides a unique opportunity to conduct research which will: inform policy-makers' detailed planning and communication strategies nationally and internationally; address the scarcity of robust evidence on the *impacts* of the implementation of smoke-free policies in prisons; and allow evaluation of the *process of implementing policies* during the preparatory period.

We have worked in close collaboration with the SPS and their Tobacco Strategy Group to develop a comprehensive research programme to fully understand: (1) the place and prevalence of smoking in Scottish prisons prior to the introduction of any new policies on smoking in prison; and (2) the impact of the implementation of any policy change.

This research comprises six workpackages (see Figure 1a): WP1 - scoping the international landscape; WP2 - (changes in) health, smoking and exposures to SHS; WP3 - staff attitudes and experiences of smoking-related issues in the prison context; WP4 - prisoners' attitudes and experiences of smoking-related issues in the prison context; WP5 - provision and impact of smoking cessation services across Scottish prisons; WP6 - stakeholder partnership working to integrate evidence from WPs1-5 to inform decision-making and communication strategies in relation to Scotland's implementation of indoor smoke-free prison facilities.

1.2 Risks and benefits

Reductions in SHS are highly likely to lead to health benefits for staff and prisoners. A World Health Organisation study (World Health Organisation 2007) estimated that SHS exposure led to 603,000 deaths in 2004. The smoking ban introduced in Scotland in 2006, and similar legislation in England in 2007, led to significant reductions in SHS in public places and positive impacts on health outcomes and public health. In Scotland, a 17% decrease in admissions for heart attack was observed following the smoking ban (Pell, Haw et al. 2008). An American study demonstrated that smoke-free rules in homes increase the likelihood of low income smokers quitting (Vijayaraghavan, Messer et al. 2013). There is evidence too of a reduction in occupational SHS exposures in those previously at high risk. A study of bar workers, for example, reported an 89% reduction in SHS exposure (as measured by salivary cotinine levels) in non-smokers after the ban (Semple, Maccalman et al. 2007) and a reduction in respiratory and sensory symptoms in smoking and non-smoking staff (Ayres, Semple et al. 2009). We are not aware of any research to date that has conducted a similar evaluation of the implementation of prison smoke-free policies on the health of workers in the prison environment (Sweeting and Hunt 2015).

In the US, half of state prisons had some sort of smoking ban by 2001, increasing to 48 states in 2011 (when 44 states banned tobacco indoors and 39 states banned tobacco or smoking indoors or outdoors). In 2014, research on the impact of these US bans on prisoner health was reported for the first time, using self-reported smoking data from a cross-sectional nationally representative survey of 14,499 individuals in state prisons in 2004, mortality data from all US state prisons for 2001-11, and data on tobacco control policies in 50 states for each of these years. This demonstrated that smoking contributed to substantial excess mortality in prisoners, and that smoking bans reduced this excess. States with the most restrictive bans had the lowest mortality rates, and where bans had been in place for longer (9+ years compared with no ban) there were significant reductions in all smoking-related deaths (adjusted incidence rate ratio 0.89, 95% confidence interval 0.85 to 0.94), cancer deaths (0.81, 0.74 to 0.82) and deaths from pulmonary disease (0.66, 0.54 to 0.80) but no reduction in mortality from causes other than tobacco for 2001-11 (Binswanger, Carson et al. 2014). However, there is little evidence on *process* - what facilitates or militates against the timely introduction of smoke-free prison policies (Sweeting and Hunt 2015).

The introduction of smoke-free prisons therefore has great potential benefits to protect and improve the health of prison staff and prisoners. If prisoners can be supported to become life-long non-smokers, these benefits may extend into the community after their release, potentially extending their impact on health inequalities. However, there could be risks to staff and prisoner safety and well-being if prison smoking 'bans' do not attract sufficient support from staff and prisoners to be enforceable, or if increased tobacco control measures threaten security at the institutional or individual level. Awofeso argues that "sustainable, humane and cost effective" approaches, including providing good smoking cessation programmes in prisons, are essential to the goal of improving the health of prisoners and staff long-term rather than a more limited goal to achieve smoke-free prisons (Awofeso 1999). Such programmes, he suggests, will also undermine the growth of a black market in tobacco and the perception of tobacco control policies as primarily punitive.

1.3 Rationale for current study

Momentum is building in many countries to increase tobacco control in prison (Offender Health Research Network 2014); (Scottish Government 2013, Public Health England 2015). This is high on media, and policy agendas (Baybutt, Ritter et al. 2014, McInerney 2015, Shaw 2015), driven by concerns to protect staff and prisoners from SHS, support health improvement in prisoners and tackle a key driver of inequalities. Prison is one of the only environments in which people are exposed to SHS in an indoor workplace or large communal living space, a cause of concern amongst staff and their unions. Whilst the UK governments are committed to immediate progress toward smokefree prisons, the timing of policy implementation was uncertain when this study was initially funded⁶. This research forms a key part of Scotland's strategy to make and evaluate "graduated progress towards smokefree prisons"⁷. The research proposed will provide data to inform this process, and will collect baseline, process and outcome data for robust evaluation of the impact of increased tobacco control in prisons/introduction of smokefree prisons. A European Commission report notes "Hardly any smoking prevention programme has been evaluated and integrated in a prison setting .. Studies addressing the effectiveness .. are few, methodologically weak, and do not provide a comprehensive overview about effects and side effects of a smoking ban in prisons." (Hartwig, Stover et al. 2008) (p6). In addition to the pressing concerns about exposure to SHS, issues around tobacco control and smoking have unique features in prisons (Richmond, Butler et al. 2006, de Viggiani 2007), both in socio-cultural meanings and the management of nicotine addiction and regulation and delivery of nicotine replacement (Eadie, MacAskill et al. 2012). For increasing restriction, or smokefree prisons, to be successful, it is important to understand the cultural value of tobacco in prisons prior to, during and following changes in legislation.

2. Research Objectives

Overall aims:

- a) To evaluate the process of implementing enhanced tobacco control in Scottish prisons to: i) strengthen the evidence-base on what is likely to facilitate successful implementation of smokefree prison policies for other jurisdictions; and ii) feed into planning and communication strategies.
- b) To evaluate: (i) changes in smoking status and exposure to second-hand smoke; and (ii) changes in related health indicators among prisoners and staff; and iii) organisational/cultural impacts in Scottish prisons, following the implementation of an indoor smokefree prison service.

Objectives:

- i. To understand barriers and facilitators to implementation of smoke-free policies, via a scoping of evidence and experiences internationally within other jurisdictions (WP1).
- ii. To evaluate changes in smoking and exposures to SHS following the implementation of a smoke-free policy in Scotland's prison service, and associated health-related indicators and costs, and other intended and unintended consequences (e.g. predictors of compliance and enforceability, incidences of unrest) (WP2).
- iii. To understand staff attitudes to and experiences of smoking-related issues in the prison context, including access to/restriction on tobacco and tobacco-related products (including e-cigarettes) in the prison environment; if/how these vary between prisons; and how these change leading up to and following the implementation of smoke-free policies (WP3).
- iv. To understand prisoners' attitudes and experiences of smoking-related issues in the prison context, including access to/restriction on tobacco/tobacco-related products (including e-cigarettes) in the prison environment; if/how these vary between prisons; and how these change leading up to and following the implementation of smoke-free policies (WP4).
- v. To evaluate provision and impact of smoking cessation services across Scottish prisons, and experiences of providers, users and potential users of these services in the lead up to implementation of indoor⁸ smoke-free prisons; and to evaluate efforts to harmonise smoking cessation services from 2016 (WP5).

⁶ As noted in the introduction, the timing of the implementation of smokefree policies across all of Scotland's prisons, 30 November 2018, was announced on 17.7.17 informed by publication of data from Phase 1 of the TIPS survey.

⁷ See section 4.7, Scotland's Journey towards Smokefree Prisons, published 13th September 2016, available at <http://www.sps.gov.uk/Corporate/Publications/Publication-4405.aspx>

⁸ Since 17.7.17, it is now known that Scottish prisons will go smokefree indoors and outdoors.

- vi. To synthesise this evidence to inform decision-making and communication strategies in relation to Scotland's implementation of indoor⁹ smoke-free prison facilities through stakeholder partnership working, and address international evidence gaps (WP6)

3. Research design

The study will take place in three phases (see Figure 1a).

- **Phase 1** will establish baseline values (for smoking, SHS and relevant health indicators) and cultural/social norms against which to assess the impact on health, economic, cultural and organisational outcomes of the implementation of smokefree prison policies. This will also be informed by scoping the international landscape.
- **Phase 2** entails a process evaluation of initiatives within Scottish prisons and events outwith the Scottish Prison Service in the period leading up to the implementation of indoor smokefree prison facilities in Scotland.
- **Phase 3** entails an evaluation of the impact of the implementation of smokefree policies on health, economic, cultural and organisational outcomes, and predictors of success or difficulties.

In Version 1.4 of this Protocol, we noted that as the exact timing of implementation of a smoke-free prison policy in Scotland was then unknown, the study may be stopped/ 'hibernated' after Phase 2 and that a key role of the independent Study Steering Committee (SSC) would be to review progress towards the implementation of indoor smokefree prison policies in the second half of 2017, and to advise then whether this should occur. Version 1.4 also stated that if "any announcement indicates that implementation will be earlier than mid-2018, the SSC will advise on adjustments required to the research to take account of Phases 1 and/or 2 being concatenated". As noted in the introduction to this version (2.5) of the Protocol, smokefree prison policies will be introduced across Scotland on 30.11.18. The amendments to this Protocol (Version 1.4 to Version 2.3) were discussed with, and approved by, the SSC at their meeting in December 2017.

Phases 1 and 3 and some Phase 2 work will take place in all Scottish prisons (n=15). Data obtained in Phase 1 will inform the selection, in consultation with the SPS, of about 6¹⁰ case-study prisons for more detailed qualitative work, to be conducted in Phase 2.

3.1 WP1 – Scoping the international landscape (Phase 1 and 2)

In order to understand barriers and facilitators to implementation of smokefree policies within other jurisdictions we will conduct telephone interviews (n=c15-25¹¹) with key personnel (including health workers) in prisons which have introduced bans to elicit their views on key challenges, successes, pitfalls and advice. We aim to include prison personnel from: New Zealand; Isle of Man; Canada; California, US; and Victoria, Australia. We may also include a small number of interviews with policy-makers and/or academics from these jurisdictions. We will also interview key personnel in pilot and other prisons introducing smokefree policies in England and Wales in 2016 (n=c.20). Interviews will be semi-structured, based on a topic guide. Participants will be encouraged to express themselves freely and to raise any additional issues which they believe are important. We anticipate that interviews will last around 30 minutes and potential participants will be made aware of this before deciding to take part.

Interviews will be audio recorded with participants' permission. Brief notes will also be taken during the interviews and may be used for reference during analysis; if any interviews take place within the prison setting rather than by telephone (e.g. in England and/or Wales), brief field notes summarising any pertinent observations or comments will be taken and may also be used during analysis. All data will be transcribed either by a member of the MRC/CSO Social & Public Health Sciences Unit, University of Glasgow (SPHSU) administrative support staff or an external transcribing service which has signed a confidentiality agreement with SPHSU/University of Stirling and been vetted for security.

⁹ See footnote above.

¹⁰ As noted in the introduction to this amended Protocol, we have increased the number of case study prisons from 4 to 6 in response to requests and feedback from SPS and our SPS Research Advisory Group.

¹¹ As noted in the introduction to this amended Protocol, in response to views expressed by staff in Phase 1 focus groups, we have shifted the balance of interviews to include more with personnel in England and Wales, and fewer with personnel in other jurisdictions.

All data will be treated confidentially and participants' anonymity will be maintained by de-identifying the data, unless an interviewee agrees to or wishes to be named. All identifiable data will be kept securely in a locked cabinet or in a password-protected computer at the University of Glasgow or the University of Stirling. Only the researchers will have access to any identifiable data.

3.2 WP2- Evaluating exposures and outcomes (Phases 1 and 3)

In this section (3.2), we describe how we will conduct the measurements of exposures to SHS. In the next section (3.3) we describe how we will obtain data on health and smoking outcomes via questionnaires administered to staff and prisoners. We will also explore with SPS the feasibility of obtaining suitably anonymised routinely-collected health data, e.g on sickness absence, among **prison staff**, in the form of a monthly series of data for each prison; we will seek to obtain only anonymised, aggregate data (i.e.no individual level or identifiable data). In respect of **prisoners**, we will also explore whether we can obtain suitably anonymised routinely-collected health care and medication data and/or other prison data and/or use anonymised linkage to deaths and Scottish Morbidity Record (SMR) hospitalisation data that would enable us to model long term changes pre- and post-implementation on smoking related disease outcomes. We will also review routine data available from ISD Scotland (WP5 lead (LB) sits on their advisory board) on quit dates set and cessation outcomes in the 12 months prior to the interviews.

Air Quality Monitoring

Instrument: Recently developed, low-cost, simple to use laser particle counters (Dylos DC1700) will measure and log minute-by-minute concentrations of second-hand smoke (Semple, Ibrahim et al. 2015). The Dylos DC1700 counts particle numbers using a low-powered laser; measures particle numbers in two size ranges (>0.5 microns and >2.5 microns) every second; and logs the average each minute. The output can be converted to provide a Particulate Matter (PM_{2.5}) concentration, a commonly used marker for SHS. Instruments have an internal memory enabling data to be collected for 6 days and 6 hours (which, within prisons, provides detail on changes in SHS levels over the course of the week to identify the impact of visiting days, weekends, and/or provision of tobacco). They require mains electricity to run for periods beyond the 4-5 hour internal battery life.

A Health & Safety (H&S) Co-ordinator and/or his/her designated representative for each prison will be trained in the use of the Dylos device at a half-day training session. Each H&S Co-ordinator will install the Dylos device in the main hall/atrium area of a residential hall within their prison to gather data on PM_{2.5} concentrations during a 6 day period. Data collection forms will be completed to indicate that the machine has been checked each day. The 6-day main hall/atrium data will be downloaded at the end of the week by connecting the device to a laptop during a visit by a member of the research team. The Dylos will then be returned to the staff member so that further 'task-based' data collection can be carried out by the H&S Co-ordinator as soon as possible afterwards to gather a series of short-term (30-min) measurements at a variety of strategic locations where staff and prisoners spend time (e.g. other residential halls/floors, canteen, games room, library) to check on exposures and compliance with existing policies. Since the devices are small enough to carry on a strap, data will also be gathered on particular tasks carried out by staff, including unlocking and entering cells where smoking is permitted. A data collection form will record the time/location for these short/mobile sampling tasks. On completion of this 'task-based' sampling, the Dylos device and the data collection sheet will be returned by the H&S Co-ordinator to one of the research staff, or by prior arrangement by courier to the University of Aberdeen/University of Stirling¹².

In addition to the Dylos, a small 'nicotine badge' will be used in Phase 1 to gather information on airborne nicotine levels over the 6 day measurement period. This 'badge' will provide an overall average of the airborne nicotine concentration at the location used. It will assist in determining the relationship between airborne nicotine and airborne PM_{2.5} concentrations and will thus help quantify how much PM_{2.5} is from tobacco sources compared to non-tobacco sources. The 'badges' (Hargreaves, Lewis et al. 2003) will be supplied by John Hopkins University and will be removed by the H&S Co-ordinator from the supplied container, positioned alongside the Dylos instrument and left exposed to the air throughout the 6-day

¹² Dr Sean Semple, the lead for SHS exposure measurements was at University of Aberdeen during Phase 1 of the TIPs study and moved to University of Stirling prior to the measurements of SHS in Phase 3.

sampling period in the main hall or atrium. At the end of the 6-day sampling prior to the researcher's visit to the prison, the 'badge' will be placed in a sealed container and collected from the prison for return to the University of Aberdeen or the University of Glasgow, together with the air quality monitoring data collection form. Training in the handling and positioning of the nicotine badge will be included in the half-day training for each H&S Coordinator or his/her nominated designate(s).

Datafiles downloaded from the Dylos device will be forwarded to the research team at the University of Aberdeen/University of Stirling¹³ as soon as possible after being downloaded. Should any issues arise, full technical support will be provided by the University of Aberdeen/University of Stirling¹⁴ by telephone and email during data collection. The H&S Coordinator or his/her nominated designate(s) will also have access to a FAQ guide and an abbreviated manual from their training session.

In order to provide comparison with outdoor PM_{2.5} concentrations (from non-tobacco, generally vehicle and industrial combustion emissions), publically available data from the Scottish Government/Local Authority Air Quality Network (<http://www.scottishairquality.co.uk/>) will be acquired from the nearest fixed monitoring site to each prison for the main 6-day monitoring period. The median distance between prisons and these outdoor PM_{2.5} monitors is 16km with only one prison located >50km from a monitoring station. The average outdoor PM_{2.5} concentration during the measurement period will then be subtracted from the indoor Dylos results to provide an estimate of SHS-derived PM_{2.5}. Results will be available with and without this adjustment.

Salivary cotinine measurement (Phase 1)

Non-smoking staff (who do not live with a smoker nor travel to work in a vehicle where smoking occurs) will be invited to participate in a study of salivary cotinine concentration changes over a typical work-shift in each of the Scottish prisons in Phase 1 measurements. Recruitment procedures may vary across prisons and will be discussed with Governors/senior staff. Our preferred method of invitation will be via an e-mail from the research team, to be forwarded by individual Governors/nominated prison staff representative. This will give brief information about the procedure, detail who would be eligible (see below), and ask those interested in participating to contact us. In addition, it may also be possible for researchers to approach and recruit staff during fieldwork visits to the prisons. We envisage high awareness of the study because of its topicality, our publicity materials and communications from the SPS, and the ongoing research activity around air quality monitoring and online staff survey (see below). Interested staff will be provided with further information. Staff volunteering to take part will be asked to complete a consent form and confirm that they understand the purpose and process of sampling and satisfy the inclusion/exclusion criteria. We aim to gather paired samples of saliva from the start and end of shifts from approximately 10% of staff from each prison (c. 400 staff in total). Inclusion criteria are that they are: (1) prison staff (preferably operational staff or staff who are likely to have some exposure to SHS during their working day); and (2) available and willing to provide a saliva sample at the start and end of their working day on the days that our fieldwork team is in attendance at their prison. Exclusion criteria are that they: (1) smoke or use any form of nicotine containing product (electronic cigarettes, patches, gums); (2) live in a home where someone smokes; or (3) travel to work in a vehicle where someone smokes.

The saliva collection protocol is based on that used for the Bar Workers' Health and Exposure to Environmental Tobacco Smoke (BHETSE) study (Semple, Maccalman et al. 2007). Participants will be asked to provide a saliva sample in a salivette tube (plastic test-tube like receptacles with a cotton wool roll inside). The salivettes have three parts: an inner tube with stopper; an outer tube (marked with a bar code and ID number); and a cotton wool roll. A member of the research team will guide the prison staff member through collecting a sample of saliva at the start and end of their working shift. The researcher will complete a data collection sheet; and will ensure the participant provides a completed log of activities they have undertaken over the course of the sampling day (using a standard proforma). The collected saliva samples will initially be temporarily stored at the University of Glasgow before being sent by the research team to the University of Aberdeen for storage and batching before analysis by ABS Laboratories Ltd. Saliva samples will be analysed for the presence of cotinine, a marker of how much nicotine a person has breathed in. By comparing the levels at the start of the shift with those at the end of the shift it is possible to estimate the amount of cigarette smoke inhaled using

¹³ See footnote 9.

¹⁴ See footnote 9.

previously published methods (Repace, Al-Delaimy et al. 2006). Any before-shift samples that are not paired with an after-shift sample will be destroyed.

Staff will also be asked about their perceived exposure to SHS in the online staff questionnaire (see below).

Prison staff SHS exposure estimates

By combining the air monitoring data gathered over 6 days from the main atrium area of one residential hall and the shorter task-based measurements from each prison with details of the typical work-shift patterns of operational staff estimates of full-shift SHS-PM_{2.5} exposure will be generated. These will be produced for a range of specific job titles and for each prison. Where possible and where sufficient samples are available, these estimates of airborne exposure will be validated by comparison with the data gathered from the salivary cotinine measurements.

3.3 WP3 – staff survey/staff focus groups in all prisons; staff telephone interviews in c.6 case-study prisons; WP4 – prisoner survey in all prisons and interviews in c.6 case-study prisons (Phases 1, 2 and 3)

Staff and prisoner questionnaires – Phases 1, 2 and 3

Questionnaires will be administered to staff and prisoners during each study phase, covering similar, but not identical topics. Both include items on: basic demographics; physical, respiratory and mental health; smoking history and behaviour, including quit attempts and use of electronic cigarettes; second-hand smoke exposure; opinions on smoking and smoking restrictions, both generally and in prisons. A Research Advisory Group (RAG), constituted and convened by the SPS, will ensure consultation with relevant parties (e.g. trade union representatives, SPS Research Access and Ethics Committee).

Staff questionnaires will be administered online, to all prison staff via an email link, forwarded by a nominated representative in each prison (see WP2 SHS measures above). The online survey will be generated and housed by IT staff at SPSU, ensuring highest standards of security. Given that both we as researchers and SPS management believe it is important that the research is, and is seen to be, independent of SPS management, the questionnaire link will be forwarded directly by us as researchers to the nominated staff member, for him/her to forward to all staff in their prison. We believe the alternative, an email forwarded by SPS Headquarters staff, would not be satisfactory in this respect.

We consulted widely on whether: staff should be asked to enter their unique SPS employee ID, in order to ensure that we could send reminders to those who had not yet completed a questionnaire and remove any duplicate responses from any member of staff who completes the survey more than once, with assurances that staff IDs would be immediately removed; or be completely anonymous. The most widely expressed view was that staff would not wish to divulge their staff ID or other potentially identifying information; hence all staff questionnaires are entirely anonymous and untraceable back to individual staff members by any means. Union representatives of prison staff have indicated that we should anticipate a high response rate from both smoking and non-smoking staff, at least in Phase 1, given the concern expressed by some prison staff about exposure to second-hand smoke during the course of their work.

Prisoner questionnaires will be administered as paper copies. Our target is to achieve a 20% randomly selected sample from all prisons; we will be advised on how best to achieve this and how best to invite prisoners to take part by each Governor-in-Charge, or his/her appointed representative. Some prisons have indicated that, on equality and pragmatic grounds, they may ask for questionnaires to be available for a greater proportion of their prisoner population. On the basis of the SPS biannual prisoner survey, we might anticipate a response of 50-60% of those approached, but response levels may be higher if prisoners view this as an opportunity to express their views on a potentially highly emotive topic. Prisoners will be provided with information about the questionnaire beforehand. In ideal circumstances, they will be: given the opportunity to discuss this and ask questions with research staff before deciding whether to participate; able to complete their questionnaires without disturbance from other prisoners, given the role of smoking and tobacco-products in prison life, including in relation to bullying and intimidation; and given the opportunity to have impartial (researcher) assistance if they have literacy or other learning difficulties. When SPS conduct their biannual prisoner survey, prisoners with literacy/learning difficulties are offered help with questionnaire completion by other prisoners; ideally we wish to avoid this in this context, and we will offer to make trained field-workers available to assist prisoners who require help (ideally checking whether prisoners have a

preference for being supported by a research staff or SPS staff member). However, operational considerations will determine the preferred means of administration of the prisoner survey in each prison. We anticipate fieldwork administering the prisoner questionnaires will be completed over a short time period (c1-3 days, depending on operational considerations and the size of the prison) in each prison, going into each prison with enough field-workers to ensure sufficient help is available given local guidance about the means of administration of the survey. Since prisons vary in both their size and individual prisoner characteristics (e.g. literacy levels), we will liaise closely with Governors/their appointed representative over the best way to achieve our aims in respect of prisoner questionnaire administration in their prison (e.g. numbers of field-workers required).

Staff focus groups – Phase 1 and Phase 3

Staff in each of the Scottish prisons will also be invited to attend focus groups (maximum participants per group=c.8, including both smokers and non-smokers) conducted around the time of the saliva collection/survey administration in Phase 1 and post-implementation of smokefree policies in Phase 3. Recruitment procedures may vary across prisons and will be discussed with Governors/senior staff. Our suggested method will be an e-mail/letter from us as researchers, to be forwarded by individual Governors/appointed representative (i.e. the same as the anticipated method for the online staff questionnaire link). This would give brief information about the focus groups and ask those interested in participating to contact us by email. Interested staff will be provided with further information. Participants in each focus group will complete a consent form before participating. SPS have suggested that focus groups should be conducted during work-time on prison premises where possible and at a time which suits participants, taking account of operational constraints in each prison and fieldwork constraints. In Phase 1 the focus groups will cover experiences of smoking and exposures to second-hand smoke, particularly within prisons; smoking norms and perceived prevalence within the prison; the 'culture' of smoking within prisons; restrictions on smoking and opinions on these; and views of prisoners who smoke. In Phase 3, they will cover experiences of the transition to smokefree prisons and any ongoing issues with maintaining smokefree policies (e.g. with new admissions, prisoners on remand, etc.)

Staff and prisoner face-to-face / telephone interviews – Phases 2 and 3

Individual interviews will be conducted with staff (face-to-face or by telephone) and prisoners (face-to-face) in case-study prisons, selected, in consultation with SPS, for more detailed qualitative work (e.g. based on more/less positive attitudes towards smoking restrictions). The number of 'case-study' prisons has been increased from four to six in response to feedback from Prison Governors, SPS and our SPS Research Advisory Group. As with the questionnaires, staff and prisoner interviews will cover similar topics, but are not identical. We anticipate similar content in phases 2 and 3, allowing us to compare responses over time. Both staff and prisoner topic guides will aim to cover: time in prison (current and other prisons – as staff or prisoner); experience of smoking and use of e-cigarettes; changes over time (particularly, for prisoners, associated with coming into prison); experiences of the working environment as a smoker/non-smoker; smoking norms and perceived prevalence within the prison; tobacco availability, trading and as 'currency'; locations of prisoner smoking; restrictions on smoking and opinions on these; views of prisoners who smoke. All interviews will be semi-structured and participants will be encouraged to express themselves freely and to raise any additional issues which they believe are important. We anticipate that interviews will last around 45 minutes and potential participants will be made aware of this before deciding to take part.

In each case-study prison in Phase 2, face-to-face / telephone interviews will be conducted with 6-8 **staff** (~3-4 smoking, 3-4 non-smoking). In order to recruit them, as with the staff focus groups, staff will be invited to participate and will complete a consent form before participating (for those participating via telephone, agreement to each item of the consent form will be recorded prior to the interview). Again, as with the staff focus groups, recruitment procedures may vary across prisons and will be discussed with Governors/senior staff, but we hope to send an initial email invite with further information for those who show interest by providing contact details.

Over Phases 2 and 3 we also anticipate conducting c.4 interviews per case study prison (n=24) with the Smokefree Point of Contact (SPOC)/implementation lead or equivalent.

In each case-study prison, individual face-to-face interviews will be conducted in Phase 2 with 6 **prisoners** (approximate ratio of 3 current and 3 ex/never smokers). We will discuss precise details on exactly how this

can be done within the operational/physical constraints of each prison with Governors/senior staff from each prison. Potential participants will be provided with an invitation to participate and will complete a consent form (or have their consent audio-taped in the case of telephone interviews) before participating, should they indicate that they wish to take part. Individual interviews will be conducted in interview rooms in the residential halls, or other suitable facilities as advised by those in charge of local prison operating facilities. Detailed practical arrangements (e.g. interview location) will be discussed with individual prison Governors/senior staff; safety and operational concerns will be paramount. In Phase 3 we anticipate conducting c.6 interviews with prisoners per case study prison.

Focus groups and interviews will be audio recorded with participants' permission. Brief field-notes will also be taken and may be used for reference during analysis and reporting of findings. All data will be transcribed either by a member of SPHSU/Institute for Social Marketing (ISM) administrative support staff or an external transcribing service which has signed a confidentiality/data processing agreement with SPHSU or University of Stirling and been vetted for security.

All data will be treated confidentially and participants' anonymity will be maintained by de-identifying the data. All identifiable data will be kept securely in a locked room/cabinet or in a password-protected computer within the SPHSU, University of Glasgow or University of Stirling¹⁵. Only the researchers will have access to any identifiable data.

3.4 WP5 – Cessation services: experiences, provision and outcomes (Phases 1, 2 and 3)

Individual interviews with cessation service leads in all prisons and associated health boards - Phase 1

Members of staff with lead responsibility for cessation support in each prison and in those health boards where prisons are located will be invited to take part in an interview either by telephone or face-to-face. These staff will be identified via the health boards, SPS, individual prison Governors/senior staff and through established NHS smoking cessation networks. Identified staff will be provided with further information (via email/post to work addresses) and asked for opt-in consent (scanned and returned via email/post or recorded prior to the interview). Interviews will be semi-structured, last up to one hour and be undertaken during work time. Topics covered will include: time (working) in prison (current and other prisons – as staff or prisoner); experience of smoking within prisons; awareness and understanding of smoking restrictions, compliance with these provisions; norms and attitudes to quitting and remaining abstinent while in prison; expectancies post-release (prisoners only); attitudes and experience of smoking cessation support within and between prisons; attitudes towards smokefree policy within prisons; views on factors likely to influence effective implementation of smokefree policy; and views on the value of e-cigarettes as a cessation and abstinence aid. Participants will be encouraged to express themselves freely and to raise any additional issues which they believe are important.

All identifiable data from WP5 will be kept securely in a locked room/cabinet or in a password-protected computer within the University of Stirling. Copies of all research data will also be archived by the University of Glasgow. Only the researchers will have access to any identifiable data.

Interviews with cessation service providers in prisons in c.6 case-study prisons – Phase 2

In each case study prison, up to six members of staff responsible for delivering and supporting cessation support in the selected case-study prisons (including a mix of professionals such as staff dedicated to providing cessation support and staff responsible for general prisoner health who are supporting cessation delivery/provision of e-cigarettes as part of their remit) will be invited to take part in an interview. It is anticipated that most of these interviews will be conducted face-to-face and will include service leads, trainers and prison healthcare staff. Telephone interviews may be offered in some instances where there are significant organisational barriers to arranging direct access. These participants will be identified from the Phase 1 interviews with prison and health board leads. In some cases it may be appropriate to interview participants together, for example if they assume similar or joint roles.

¹⁵ Prof Kate Hunt and Ashley Brown (appointed researcher on TIPs) moved from University of Glasgow to University of Stirling during Phase 2 of the TIPs Study.

Following the decision by the SPS to implement a comprehensive smokefree policy in all Scottish prisons on 30th November 2018, it is proposed merging the two stages of field work originally envisaged for Phase 2 into a single stage to extend over the remaining eight month period leading up to the implementation deadline in November 2018. The decision to collapse the two stages of field work into one gives the study the scope to expand the number of case study prisons from 4 to 6 in response to Prison Governors, SPS and our SPS Research Advisory Group, and to represent a wider range of establishments and prisoner groups. It is intended that the Phase 2 fieldwork will assess existing smoking cessation services and development needs, as well as examine preparations and progress towards implementation, including barriers and facilitators. As with Phase 1, all interviews and discussion groups will be semi-structured, will last up to one hour and will be undertaken in work-time and at a time suitable to participants. The issues covered will be similar to those described at Phase 1 with a focus on implementation and service development. Staff who have been identified to take part will be provided with further information and asked to complete a consent form before participating. For those participating via telephone, and where the respondent chooses not to provide a scanned version of the consent form, oral consent will be recorded prior to the interview.

Interviews or discussion groups with prisoners (smokers and former smokers) in c. 6 case-study prisons – Phase 2

In each case study prison, interviews or small discussion groups will be conducted with prisoners who smoke or who are ex-smokers and who have used, are currently using or are planning to use structured smoking cessation support. We would anticipate speaking with up to 9 prisoners in each prison, ideally in two small discussion groups, although individual or paired interviews would be conducted if necessary, depending on operational need. Precise details on recruitment for and conduct of the prisoner focus groups/interviews will be discussed with Governors/senior staff from each prison. Participants who have been identified to take part will be provided with further information and asked to complete a consent form before participating.

Prisoner interviews will cover similar topics to the staff interviews from a prisoner's perspective and will examine the perceived value and delivery of cessation support and e-cigarette provision in the lead up to going smokefree, as well as expectations and intentions re. smoking and vaping post-release, and any ongoing cessation support needs as appropriate.

Individual interviews with cessation service leads in all prisons and associated health boards - Phase 3

The original proposal was designed to examine the provision of cessation support within the prison service in Scotland. Given the SPS is now committed to implementing a comprehensive smokefree policy across the full prison estate within the lifetime of the study, WP5 has been extended to include an additional phase of research (Phase 3) to examine the role cessation support plays in facilitating the transition to smokefree prisons and the need for cessation support in maintaining a smokefree environment post-implementation. In broad terms this will replicate the Phase 1 methodology, by carrying out a series of interviews (n~30, telephone or face-to-face) with prison healthcare leads responsible for providing cessation support and/or community quit services responsible for delivering cessation support within their local prison(s) depending on organisational approach. Field work will be undertaken in April- August 2019 once the new policy has had an opportunity to bed in and will cover all 15 prisons and leads from those health boards where prisons are located.

4. Study population

WP1: representatives from prisons internationally with experience of implementing partial/total smoking bans within prisons

WP2-4: adult (age 18 or over) prisoners and prison staff within all 15 prisons in Scotland. This includes a female-only prison, two private prisons, one open prison, and a Young Offenders Institution (16-21 years) that operates a smokefree policy for its under 18 population. For the qualitative work, participants will be purposively sampled on the basis of their smoking status.

WP5: users and providers of smoking cessation services within the 15 prisons in Scotland, including SPS and NHS staff, and adult prisoners.

Participants will be excluded from WP2-5 if we are advised by SPS that specific potential participants raise health/safety issues for either researchers/field staff, or the prisoner him/herself, or if they have

communication/health issues which preclude participation. It is anticipated that interviews will be conducted in English, unless arrangements can be made for staff/volunteers to be present to translate.

5. Socio-economic position and inequalities

Around half of those who smoke will die from tobacco-related illnesses, and tobacco-attributable deaths are estimated to be 6.4 million globally in 2015. In Scotland, tobacco use was estimated to be associated with a quarter of deaths (approximately 13,000 per annum) and 56,000 hospital admissions in 2013, with an annual cost to the Scottish Health Service of £300-500m (Scottish Government 2013). A particular challenge for ambitions to achieve smokefree societies is that tobacco and poverty have become “inextricably linked” (Hartwig, Stover et al. 2008) (p5). In Scotland in 2011, more than three times as many (38%) of those living in areas classified as in the most deprived quintile were current smokers compared with the most affluent quintile (12%) and estimates for 2016 suggest little erosion of this disparity (27% vs 10% respectively) (Scottish Government 2013). Health Survey for England data show that 80% of smokers are deprived on at least one of seven commonly used indicators of deprivation (NS-Socio-economic classification, neighbourhood index of deprivation, lone parent, car access, housing tenure, income and unemployment), and the prevalence of smoking rises from 15% in those with no indicators of low SES to 61% of those with all seven (Hiscock, Bauld et al. 2012). In his foreword to Scotland’s current smoking strategy, the Minister for Public Health observed that “perhaps the most troubling aspect of current smoking patterns is the hugely disproportionate impact on Scotland’s most deprived [and]...effective action to reduce smoking prevalence demands a focus on those communities at greatest risk of unequal health outcomes” (Scottish Government 2013) (piii). The Report states “We will not achieve our ambition of a tobacco-free Scotland without addressing the stark socio-economic inequalities in smoking prevalence rates” (p2). When considering ‘Inequalities-Targeted Services’, prisoners are highlighted: “We recognise the real need to work towards protection of offenders from second-hand smoke and in the provision of effective smoking cessation services” (p41). These statements reflect a growing recognition that if inequalities in the harms of smoking are to be reduced, prisoners are an important target population (Public Health England 2015): as well as having disproportionately high substance abuse and mental health problems which are associated with high smoking rates, they are also at increased risk of factors associated with smoking, including having low/no educational qualifications, having experienced poverty, unemployment¹⁶, homelessness, exclusion from school, adverse family and social experiences such as domestic and/or sexual abuse and time in local authority care (World Health Organization Regional Office for Europe 2013).

6. Planned interventions

- a) Harmonisation and increased provision of smoking cessation services for prisoners (including health service provision as provided by Health Boards and other smoking cessation support) across Scottish prisons. It is anticipated that this will increasingly occur during Phases 1-2 of the study.
- b) Implementation of smokefree prison policy on 30 November 2018 and preparatory actions leading up to this event.

7. Proposed outcome measures

Primary outcomes: Current smoking in prisoners; objective indicators of SHS (air quality within prisons [PM_{2.5}]).

Secondary outcomes:

Staff: smoking status; general health and well-being; respiratory symptoms; use of primary health care; self-reported and/or aggregated sickness absence data;

Prisoners: self-perceived smoking status (i.e. extent to which they see themselves as lifelong quitters to estimate impacts post-release); general health and wellbeing; use of primary care; (if available) aggregate medication and health data; acceptance of and compliance with current tobacco control policy.

8. Assessment and follow-up

¹⁶ Recent research also suggests that smokers remain unemployed longer than non-smokers: Prochaska JJ et al. Likelihood of unemployed smokers vs nonsmokers attaining reemployment in a one-year observational study [published online April 11, 2016]. *JAMA Intern Med*. doi:[10.1001/jamainternmed.2016.0772](https://doi.org/10.1001/jamainternmed.2016.0772).

Measurement of all primary and secondary outcomes (including measurement of indicators of SHS exposures) will be repeated post-implementation of smokefree prisons¹⁷ (Phase 3), following the same protocols as at baseline (Phase 1). SHS measurements using the Dylos DC1700 will be undertaken twice in Phase 3 in all prisons, once in the week of implementation (end Nov/early Dec 2018) and once in April/May 2019. Interim process (attitudes/experiences) data (in anticipation of implementation of policy change) will be collected, via staff and prisoner focus groups and interviews data from c. six case study prisons, and repeat staff and prisoner surveys in Phase 2 in April/May 2018, as described above. Anticipated timings are shown in Figure 1a, and in the project timeline and milestones, subject to decisions re policy changes nationally or locally.

9. Proposed sample size:

WP1: We propose around 8 telephone interviews with key personnel in countries which have introduced smokefree prison policies, and face-to-face/telephone interviews (n=c.20) with key staff in prisons introducing smokefree in England and Wales in 2016-2018.

WP2. Survey data will be gathered from c.2700 prison workers (estimated 60% response rate from staff within Scottish prisons of 4500 invited).

Staff salivary cotinine measures: At Phase 1, we anticipate that c30% will be smokers, and perhaps a further 25% live with or travel with a smoker or spend significant non-occupational time in 'smoky' environments (e.g. travel to work with smoking colleagues).

Air quality measures: Six-day air quality measurements (collected using laser particle counters) will be made in each of Scotland's 15 prisons. These will provide an average PM_{2.5} concentration together with the percent of time above specific thresholds such as the WHO guidance limit of 25 µg/m³ and other comparative markers (e.g. average levels measured in pubs prior to implementation of smokefree legislation in Scotland in 2006). Around six 30-minute 'task-based' measurements will also be made in each prison. For comparison, mean PM_{2.5} over a 24 hour period was found to be 33.3 and 4.4 in smoking and non-smoking homes respectively (Semple, Ibrahim et al. 2015). Comparison data will now be collected in the week of implementation of smoke-free prisons and approximately 4-6 months after implementation.

WP3. Staff survey: as above, we will invite all staff to participate in the online staff survey (n=c4500 staff) and aim to gather data from 60%. A 50% response with a true prevalence of smoking of 30% would mean a tolerance of + 1.5%, giving a 95% confidence interval for the estimate of prevalence of 28.5%-31.5%.

Qualitative: Phase 1, 1 FG in each of 15 prisons with a mixture of smoking and non-smoking staff (15 FGs, total n=90 staff, assuming 3 smoking and 3 non-smoking staff members in each group). Phase 3, 1 FG in each of 15 prisons with a mixture of smoking and non-smoking staff (15 FGs, total n=90 staff, assuming 3 smoking and 3 non-smoking staff members in each group).

In Phase 2 semi-structured face-to-face or telephone interviews with 6 staff (including a mix of smokers and non-smokers) in each of 6 case-study prisons (total n=36 in Phase 2). In Phases 2 and 3 interviews with 4 staff in each of 6 case study prisons (total n=24 in Phase 2 and 3).

WP4 Prisoner survey with c20% subsample of prisoners. We assume a sample size of n=1480 prisoners and a true prevalence of smoking of 70%. In the absence of a known figure we again assume a conservative ICC of 0.15. This would provide power of 0.8 to detect a fall in prevalence among male prisoners from 70% to 48%.

Qualitative: 6 45 minute face-to-face semi-structured individual interviews (including a mix of smokers, ex-smokers and never smokers) in each of c.6 'case-study' prisons in Phase 2 (total n=36 in Phase 2). 6-8 individual interviews with prisoners in each of c.6 'case study' prisons in Phase 3 (total n=36 in Phase 3).

WP5 Cessation service (CS): experiences/provision.

Phase 1: Mapping of prison estate and cessation service links - assessment of preparedness and proposed plans for harmonisation - 15 telephone interviews with prison smoking cessation service leads; up to 15 telephone interviews with Health Board smoking cessation service leads.

¹⁷ As noted in the introduction to Version 2.5, the policy to implement smokefree policies across all of Scotland's prisons by 30 November 2018 was announced by SPS on 17.7.17.

Phase 2: Formative and process evaluation in six case study prisons - development and preparation for implementation of smokefree.

- Interviews with up to six staff responsible for delivering and supporting cessation support in each case-study prisons (N=36)

-Interviews/small group discussions with up to nine prisoners who smoke or who are ex-smokers and who have used, are currently using or are planning to use structured smoking cessation support in each case study prison (N=54)

Phase 3: Process and impact evaluation of smoking cessation support delivery post-implementation of smoke-free across the prison estate.

- Interviews with smoking cessation leads; up to 15 interviews with Health Board smoking cessation service leads and 15 with prison health care leads responsible for cessation support.

10. Analysis

10.1 Statistical analysis: Analysis of objective measures of SHS will follow previously published methods (Feyeraband and Russell 1990, Repace, Al-Delaimy et al. 2006, Semple, Ibrahim et al. 2015). SHS levels in the prisons will be assessed using air quality monitors. We will compare mean PM_{2.5} concentrations before and after the implementation of a smokefree policy within each prison over a 6 day period. We will compare each site individually using paired sample t-tests and collectively using a multilevel model including the levels of prison, site, measurement occasion (day and hour) and wave (pre- and post-intervention).

Subject to the anonymised, aggregate data being made available, we will calculate sickness absence rates as the number of hours lost due to sickness absence as a proportion of the total number of hours that could have been worked by prison officers in each prison for each month. We will identify the impact of the smokefree prison policy on sickness absence among prison officers by fitting interrupted time series models to the monthly data. The models will take account of pre-implementation trends, seasonality and the way in which any new policy is introduced (i.e. whether the transition to smokefree is abrupt or gradual). Data permitting, we shall also analyse sickness absence rates by cause, distinguishing smoking-related absences from all other absences. We will continue to explore what other routinely collected data may be used for similar models examining changes over time.

Analyses of quantitative questionnaire data will include: basic descriptive statistics (e.g. proportions agreeing with particular attitudinal items); and examination of change in prisoner/staff smoking prevalence, health and attitudes over time via multilevel logistic regression with the levels of prisons, study phases and prisoners, treating the samples at different waves as independent (repeat cross sections) since there will be considerable movement of prisoners and adjusting for socio-demographic characteristics. Analyses will assess missing data patterns and use multiple imputation¹⁸ to create multiple datasets with as many relevant predictor variables as possible included in the imputation models. Questionnaire data will also be utilised in health economic modelling.

Any change in the prevalence of smoking among prisoners will be assessed using multilevel logistic regression, treating the samples at different waves as independent since there will be considerable movement of prisoners, with the levels of prisons, waves and prisoners. We will adjust for socio-demographic characteristics.

Secondary outcomes collected by survey will be analysed by multilevel linear, logistic or multinomial regression depending on the outcome. In all analyses we will assess missing data patterns and use multiple imputation to create multiple datasets for analysis with as many relevant predictor variables as possible included in the imputation models.

10.2 Qualitative data analysis: Interview and focus group transcripts will be checked for accuracy against the digital recordings, suitably anonymised, and analysed using a structured thematic approach (Ritchie and Lewis 2003) to explore our research questions and any emerging themes. Several members of the research team with qualitative expertise will read anonymised transcripts to support the development of robust analytical frameworks for each of the embedded qualitative studies. Coding frames and codebooks will be developed

¹⁸ We will investigate whether it is possible to use relevant data from the SPS biannual prisoner questionnaire to inform imputation of missing data on smoking

through discussion and close reading of transcripts, and will be tested and discussed to ensure common understandings of key themes before coding to broad themes is undertaken. The content of the themes relevant to the key research questions will be examined in detail by at least two researchers, who will read coded extracts line by line to identify all sub-themes, using suitable methods to enable systematic comparison of data from different participants/groups, noting anticipated and unanticipated themes. Attention will be paid to 'deviant cases' to ensure all perspectives are captured. Nvivo software will be used to assist in the organisation and analysis of the qualitative data.

11. Ethical arrangements

Ethical approval for WPs1-5 will be obtained from Glasgow University College of Social Sciences Ethics Committee. All elements of the study will also be separately approved by the Scottish Prison Service Research Access and Ethics Committee, who were asked to advise whether research within any of the work packages also required approval from an NHS Research Ethics Committee. Given WP5 involves interviews with Health Board smoking cessation managers, advice has been sought from the South East Scotland Research Ethics Service. We have been advised that the research does not need NHS ethical review under the terms of the Governance Arrangements for Research Ethics Committees. Research for WP1 may require additional ethical approval, e.g. for prisons in England and Wales, from HMPPS.

Research involving prisoners, as with other potentially vulnerable groups, raises many issues (Schlosser 2008) and requires particularly careful attention to ethics. Prison governors, prison staff and prisoners will be informed about the study via in-person visits by the PI (Kate Hunt) and/or Helen Sweeting, together with a representative of Health and Wellbeing at SPS (Sarah Corbett and/or Ruth Parker), in advance of any data collection. These will be arranged via SPS, and a key point of contact within each prison will be agreed. Permission for the participation of each prison will be required from the SPS and prison governor. Following these in-person visits, general information leaflets and posters alerting staff and prisoners to the research will be distributed and displayed within each of the prisons by prison staff in advance of data collection, so that staff and prisoners are aware that they may be approached about the study.

Participants in any element of the research will receive an information sheet explaining the aims and objectives of the research that they are being asked to contribute to. This will clearly indicate: why they have been approached; that their participation is voluntary and they are free to withdraw at any time and have no obligation to answer any questions they do not wish to; that all data will be anonymised and stored securely, and only be reported in anonymised form to any parties. They will be given assurances that their answers will be treated in the strictest confidence, explaining the independence of the research team from Scottish Government and the SPS. Only aggregate and anonymised data will appear in any Reports, publication or presentations. In particular, prisoners will be reassured that their participation (or lack of participation) will not in any way influence their care or treatment in prison. Written or audio recorded opt-in consent will be obtained for all qualitative interviews and measures of personal exposure to SHS. As the online staff survey and prisoner survey have no identifying information, it will be explained that completion of the online survey by staff and the paper questionnaire survey by prisoners is deemed to constitute consent, unless other requirements are detailed during ethical review. We will consult with the SPS and unions before finalising the content of the questionnaires.

Prisoner questionnaire data will be entered in-house at SPSU and transcription of qualitative data will be undertaken by a specialist transcription service or university administration staff bound by confidentiality guidelines. Data from questionnaires will be stored in an anonymised format, using questionnaire identification numbers (IDs).

For focus groups and individual qualitative interviews, participant ID numbers and corresponding names will be held in separate files and all will be stored in password protected folders. Individual names will be replaced with pseudonyms in transcribed interview materials; we will follow SPS advice and guidance on anonymisation of prisons. Digital recordings of interviews (made with written participant permission) will be stored securely, and held separately from transcripts and information on participant identity. Care will be taken to prevent deductive disclosure of identities in the use of selected extracts from interviews/focus groups in all project outputs. All quotations will be anonymised, except where there is an explicit request and permission for an

attributed quote. We will follow data service requirements for recording and management of data (<http://ukdataservice>). SPHSU is bound by the MRC's and Glasgow University's rigorous research governance policies. Research data will be destroyed according to current MRC research guidelines. In line with this, the research data will be kept for ten years after the study has been completed, and then destroyed (in accordance with MRC and University of Glasgow policy (audio and electronic documents will be securely deleted and paper copies will be shredded).

Any member of the research team visiting a prison will be required to undergo appropriate security checks, and any member who is expected to have one-to-one encounters with prisoners will be required to complete any safety and security training as recommended by SPS, in addition to training in SPHSU's standard fieldwork safety procedures for research team members in community settings (as required). All research will be carried out in accordance with MRC best practice guidelines (<http://www.mrc.ac.uk/publications/browse/good-research-practice-principles-and-guidelines/>), the ESRC Framework for research ethics (<http://www.esrc.ac.uk/files/funding/guidance-for-applicants/esrc-framework-for-research-ethics-2015/>), and any relevant guidance on research practice from SPS and Health Boards. At all times, safety and prison operational considerations, and any impacts on the delivery of health care in the prisons, will dictate the day-to-day conduct of the research within each prison.

12. Research Governance: University of Stirling will be the nominated sponsor of the research.

An **Operational Research Management Group** will meet ~fortnightly (chaired by Kate Hunt, with Helen Sweeting and other SPHSU/ISM research/support staff involved in other aspects of the overall study).

Project Research Team meetings (all co-Investigators on the NIHR proposal) will occur face-to-face or by teleconferencing at least quarterly.

SPS Research Advisory Group: A Research Advisory Group will be convened by SPS, comprising representatives of the key stakeholders they identify. This will include representatives from stakeholder partners who contributed to SPS' 2015 Tobacco Strategy¹⁹ group (SPS Health and Wellbeing team; Scottish Government Tobacco Control Team; NHS Health Scotland; SPS trade union representatives or similar representative of prison staff) and representatives of SPS Communications branch, SPS Smokefree Prisons Programme Lead (Linda Doward) and Programme Manager (Sarah Corbett) and Research Ethics Committee. It is anticipated that the RAG will meet at least bi-annually during most phases of the overall study, and more often as deemed necessary. This group will ensure that proposed activities are achievable and realistic within the resources available and any operational constraints within the day-to-day good governance of prisons, and that a focus is maintained on achieving the outputs SPS considers a priority. In addition, during Phase 2, and if required Phase 3, the PI (Kate Hunt), Helen Sweeting and other core team members (e.g, appointed researcher Ashley Brown, Dr Sean Semple) will meet c.monthly with core members of the SPS Smokefree Prisons Programme (Linda Doward and/or Sarah Corbett) and attend monthly meetings of SPS Stakeholder Advisory Group for the smokefree team to provide feedback on new results/findings from ongoing analyses.

Study Steering Committee (SSC): The SSC will meet face-to-face at least annually, with additional teleconferences/ meetings as required, chaired by Dr Andrew Fraser, NHS Health Scotland.

13. Project timetable and milestones:

In the original version of this Protocol (Version 1.4) we noted that "All timings beyond the early stages of Phase 1 may be subject to change, depending on the timing of any Government or prison service announcements about the implementation of new policies on smoking in prisons in Scotland. The project timetable and milestones will be reviewed with NIHR, the SPS-convened Research Advisory Group, the Study Steering Committee, and with the Research team in the event of the announcement of a timeframe for the implementation of new policies on smoking in prisons." Following the announcement on 17.7.17, as described in the introduction to this revised Protocol, we have consulted and gained approval from the SPS-convened RAG, our SSC (meeting of 23.12.17) and NIHR.

¹⁹ The membership of the SPS Research Advisory Group changed during 2017 to include those central to the implementation of the smokefree policy.

Dates	Milestone	Workpackage
Feb-Jul 2016	Obtain ethical approval from SPS Research Access and Ethics Committee; check whether NHS ethical permission needed for WP5	All
Jul-Oct 2016	Obtain local Glasgow University ethics for WP1-5	All
<i>1st Sept 2016</i>	<i>Project starts</i>	
Apr-Sept 2016	Initial consultations with SPS Research Advisory Group. In-person visits to all prisons to meet senior staff to discuss research and establish key point of contact in each prison. Finalisation of staff survey questionnaire and topic guides; drafting of prisoner questionnaires.	All
Sep 2016 – May 2018	Collect and analyse interviews for WP1; update literature scoping review	WP1
Sep-Dec 2016	Phase 1 measurement of SHS exposures	WP2
Oct-Dec 2016	Phase 1 staff survey and staff focus groups	WP3
Nov 2016-Apr 2017	Phase 1 prisoner survey	WP4
Oct 2016-Feb 2017	Scoping of smoking cessation provision; Phase 1 interviews with smoking cessation staff	WP5
Dec 2016-Jan 2017	Analysis of Phase 1 staff survey. Prepare summary of Phase 1 staff survey and present to RAG	WP3; WP6
Dec 2016-May 2017	Analysis of SHS exposure data; analysis of prisoner survey; analysis of staff focus groups; analysis of scoping of smoking cessation provision. Prepare summaries and present to RAG, prepare papers and dissemination materials for prison staff and prisoners	WP2-5; WP6
<i>July 2017</i>	<i>Phase 1 complete; Phase 2 commences</i>	
July-Oct 2017	Selection of case study prisons	WP3-5; WP6
Aug 2017-Oct 2018	Staff and prisoner Phase 2 interviews and analysis	WP3-4
Aug 2017-Oct 2018	Phase 2 Interviews with providers/users of smoking cessation services	WP5
April-May 2018	Phase 2 staff survey	WP3
April-May 2018	Phase 2 prisoner survey	WP4
Jan-Nov 2018	Review policy development; analysis of Phase 2 data; prepare summaries and present to SPS' RAG/SAG and smokefree implementation team	WP and All
Jun-Dec 2018	Analysis and dissemination of Phase 1 and Phase 2 data	All
<i>Dec 2018</i>	<i>Phase 3 commences</i>	
Nov/Dec 2018	Phase 3 measurement of SHS exposures in week of implementation	WP2
Dec 2018-May 2019	Phase 3 (post-implementation) interviews with smoking cessation staff	WP5
April-May 2019	Phase 3 (post-implementation) measurement of SHS exposures	WP2
April-May 2019	Phase 3 (post-implementation) staff survey	WP3
April-May 2019	Phase 3 (post-implementation) prisoner survey	WP4
Dec 2018-Nov 2019	Phase 3 (post implementation) staff and prisoner interviews	WP3, WP4
Dec 2018-May 2020	Analysis; modelling of impacts; continuing write up of papers from Phase 2 and Phase 3	WP1-5; WP6
June 2020	Project ends; outcomes paper submitted for publication	
~Sep 2020	Project report submitted	

14. Expertise

- Service and academic public health; social scientific, epidemiological and biomedical research on smoking and smoking cessation (Kate Hunt, Helen Sweeting, Sean Semple, Linda Bauld, Douglas Eadie, Jill Pell, Alastair Leyland, Philip Conaglen)

- Measurement of indoor air pollution including exposure to SHS and other occupational exposures, in prisons and other settings (Sean Semple, Evangelia Demou)
- Evaluating smoking cessation services and the implementation of smokefree policies (Sean Semple, Linda Bauld, Douglas Eadie, Jill Pell, Kathleen Boyd)
- Evaluating complex public health interventions (Kate Hunt, Linda Bauld, Peter Craig, Alastair Leyland, Jill Pell, Kathleen Boyd) and natural experiments (Peter Craig, Alastair Leyland, Jill Pell, Kathleen Boyd)
- Improving health and wellbeing in those with contact with the criminal justice system (Philip Conaglen)
- Design, conduct and analysis of health surveys (Kate Hunt, Helen Sweeting, Jill Pell, Alastair Leyland, Kathleen Boyd, Evangelia Demou, Philip Conaglen); use of routine health data through linkage (Jill Pell, Alastair Leyland, Emily Tweed); use of routine data on prisoner and similar populations (Emily Tweed); advanced statistics and health and economic modelling (Alastair Leyland, Kathleen Boyd); inequalities in health and health behaviour (Kate Hunt, Helen Sweeting, Jill Pell, Alastair Leyland, Philip Conaglen, Emily Tweed); conducting qualitative research in prison settings (Kate Hunt, Douglas Eadie) and qualitative data analysis (Kate Hunt, Helen Sweeting, Linda Bauld, Douglas Eadie); and masculinities, health and health behaviour (Kate Hunt, Helen Sweeting), including in a prison setting.

In addition, research team members have served on relevant national bodies or have direct experience of policy-making and research, including: LB is CRUK's Cancer Prevention champion, with expertise in smoking cessation services and use of evidence to inform policy and was UK government's scientific adviser on tobacco control (2006-10); HS is Chair of the Scottish Tobacco-free Alliance Research Group and a member of ASH Scotland Policy and Development sub-committee; JP is a member of the CLEAN collaboration, a pan-Scotland group evaluating impacts of Scottish smokefree legislation; PCr has policy experience within central government, previously managing research on health services and public health research; PCo represents Scottish Directors of Public Health on the Scottish National Prison Healthcare Network Advisory Board and has responsibility for improving health and being in those in contact with the CJS for NHS Lothian.

15. Partner Collaboration

We will continue to work in close collaboration with representatives from key partner organisations, including the SPS (through Ruth Parker (Head, Health and Wellbeing), Linda Dorward, Sarah Corbett and Tina Everington (Health Improvement Manager)); Scottish Government (Fiona Dunlop, National Tobacco Control Advisor); NHS Health Scotland (Celia Gardiner, Tobacco Programme Manager); and Health Boards. SPS will facilitate access to prisons and resources, subject to approval by their Research and Ethics Committee, to cover any operational implications of the conduct of the research (e.g. escorting researchers/field staff through prison facilities and grounds, distributing study information sheets to appropriate parties, facilitating support for prisoner survey). We will ensure rapid dissemination of all phases of the research to prison management, prison staff and prisoner populations.

16. Funding

Funding for this research has been obtained from the NIHR Public Health Research funding board (Grant no: 15/55/44).

17. References

- Australian Institute of Health and Welfare (2015). The health of Australia's Prisoners 2015. Canberra, Australian Institute of Health and Welfare.
- Awofeso, N. (1999). "Controlling tobacco use within prisons." International Journal of Tuberculosis and Lung Disease **3**: 547-548.
- Ayres, J. G., S. Semple, L. MacCalman, S. Dempsey, S. Hilton, J. F. Hurley, B. G. Miller, A. Naji and M. Petticrew (2009). "Bar workers' health and environmental tobacco smoke exposure (BHETSE): symptomatic improvement in bar staff following smoke-free legislation in Scotland." Occupational and Environmental Medicine **66**(5): 339-346.
- Baybutt, M., C. Ritter and H. Stover (2014). Tobacco use in prison settings: a need for policy implementation. Prisons and Health. S. Enggist, L. Moller, G. Galea and C. Udeson. Geneva, World Health Organisation.
- Beck, G. (1995). "Bullying among young offenders in custody." Issues in Criminological and Legal Psychology **22**: 54-70.
- Binswanger, I. A., E. A. Carson, P. M. Krueger, S. R. Mueller, J. F. Steiner and W. J. Sabol (2014). "Prison tobacco control policies and deaths from smoking in United States prisons: population based retrospective analysis." British Medical Journal **349**.
- Butler, T., R. Richmond, J. Belcher, K. Wilhelm and A. Wodak (2007). "Should smoking be banned in prisons?" Tobacco Control **16**(5): 291-293.
- Carpenter, M. J., J. R. Hughes, L. J. Solomon and T. A. Powell (2001). "Smoking in correctional facilities: a survey of employees." Tobacco Control **10**(1): 38-42.
- Collinson, L., N. Wilson, R. Edwards, G. Thomson and S. Thornley (2012). "New Zealand's smokefree prison policy appears to be working well: one year on." New Zealand Medical Journal **125**(1357): 1-5.
- de Viggiani, N. (2007). "Unhealthy prisons: exploring structural determinants of prison health." Sociology of Health and Illness **29**: 115-135.
- Donahue, J. J. (2009). "Tobacco Smoking Among Incarcerated Individuals: A review of the nature of the problem and what is being done in response." Journal of Offender Rehabilitation **48**: 589-604.
- Eadie, D., S. MacAskill, J. McKell and M. Baybutt (2012). "Barriers and facilitators to a criminal justice tobacco control coordinator: an innovative approach to supporting smoking cessation among offender." Addiction **107** (Suppl 2): 26-38.
- Enggist, S., L. Moller, G. Galea and Udesen (2014). Prisons and Health. Geneva, World Health Organisation.
- Feyerabend, C. and M. Russell (1990). "A rapid gas-liquid chromatographic method for the determination of cotinine and nicotine in biological fluids. ." J Pharm Pharmacol **42**: 450-452.
- Gautam, J., M. Glover, A. Scott and D. Welch (2011). "Smoke-free prisons in New Zealand: maximising the health gain." New Zealand Medical Journal **124**: 100-106.
- Globalsmokefree Partnership (2009). Reducing Tobacco Smoke Exposure in Prisons. Switzerland, Globalsmokefree Partnership, World Heart Federation.
- Hargreaves, I., J. Lewis and T. Speers (2003). Towards a better map: science, the public and the media, Economic and Social Research Council.
- Hartwig, C., H. Stover and C. Weilandt (2008). Report on tobacco smoking in prison.
- Hiscock, R., L. Bauld, A. Amos and S. Platt (2012). "Smoking and socioeconomic status in England: the rise of the never smoker and the disadvantaged smoker." Journal of Public Health **34**(3): 390-396.
- Hyland, A., M. Travers, C. Dresler, C. Higbee and K. Cummings (2008). "A 32 country comparison of tobacco smoke derived particle levels in indoor public places." Tobacco Control **17**(3): 159-165.
- Kouvonen, A., M. Kivimaki, T. Oksanen, J. Pentti, T. Heponiemi, A. Väänänen, M. Virtanen and J. Vahtera (2012). "Implementation of workplace-based smoking cessation support activities and smoking cessation among employees: The Finnish Public Sector Study." Am J Public Health **102**: e56-e62.

Lankenau, S. (2001). "Smoke 'em if you got 'em: cigarette black markets in U.S. prisons and jails." The Prison Journal **81**: 142-161.

MacDonald, L., K. Angus, S. MacAskill and D. Eadie (2010). Rapid Literature Review of Smoking Cessation and Tobacco Control Issues Across Criminal Justice System Settings. Stirling, Institute for Social marketing, University of Stirling.

McInerney, M. (2015). "Smoking ban under spotlight after Melbourne prison riot. Australian experts are split over whether smoking bans are a crucial advance in prison health or "a bridge too far" that can only spark unrest." Retrieved 3 July 2015, from <http://www.bbc.co.uk/news/world-australia-33358585>.

Mulcahy, M., D. Evans, S. Hammond and R. J (2005). "Secondhand smoke exposure and risk following the Irish smoking ban: an assessment of salivary cotinine concentrations in hotel workers and air nicotine levels in bars." Tobacco Control **14**: 384-388.

NHS Health Scotland (2015). Specification for national prisoner smoking cessation service in Scotland. Edinburgh, NHS Health Scotland.

Offender Health Research Network (2014). Smoking in Prisons in England and Wales: An examination of the case for public health policy change.

Pell, J. P., S. Haw, S. Cobbe, D. E. Newby, A. C. H. Pell, C. Fischbacher, A. McConnachie, S. Pringle, D. Murdoch, F. Dunn, K. Oldroyd, P. MacIntyre, B. O'Rourke and W. Borland (2008). "Smoke-free Legislation and Hospitalizations for Acute Coronary Syndrome." New England Journal of Medicine **359**(5): 482-491.

Public Health England (2015). Reducing Smoking in Prisons. Management of tobacco use and nicotine withdrawal. PHE publications gateway number 2014588. London, Public Health England.

Repace, J., W. Al-Delaimy and J. Bernert (2006). "Correlating atmospheric and biological markers in studies of secondhand tobacco smoke exposure and dose in children and adults." J Occup Environ Med **48**(2): 181-194.

Richmond, R., T. Butler, J. Belcher, A. Wodak and K. Wilhelm (2006). "Promoting smoking cessation among prisoners: feasibility of a multi-component intervention." Australian and New Zealand Journal of Public Health **30**: 474-478.

Richmond, R., T. Butler, K. Wilhelm, A. Wodak, M. Cunningham and I. Anderson (2009). "Tobacco in prisons: a focus group study." Tobacco Control **18**(3): 176-182.

Richmond, R. L., T. G. Butler, D. Indig, K. A. Wilhelm, V. A. Archer and A. D. Wodak (2012). "The challenges of reducing tobacco use among prisoners." Drug & Alcohol Review **31**: 625-630.

Ritchie, J. and J. Lewis (2003). Qualitative Research Practice: A Guide for Social Science Students and Researchers. London, Sage.

Ritter, C., C. K. Huynh, J.-F. Etter and B. S. Elger (2012). "Exposure to tobacco smoke before and after a partial smoking ban in prison: indoor air quality measures." Tobacco Control **21**(5): 488-491.

Schlosser, J. (2008). "Issues in interviewing inmates. Navigating the methodological landmines of prison research. ." Qualitative Inquiry **14**: 1500-1525.

Scottish Government (2013). Creating a tobacco-free generation. A tobacco control strategy for Scotland. Edinburgh.

Semple, S., A. E. Ibrahim, A. Apsley, M. Steiner and S. Turner (2015). "Using a new, low-cost air quality sensor to quantify second-hand smoke (SHS) levels in homes." Tobacco Control **24**(2): 153-158.

Semple, S., L. Macalman, A. A. Naji, S. Dempsey, S. Hilton, B. G. Miller and J. G. Ayres (2007). "Bar Workers' Exposure to Second-Hand Smoke: The Effect of Scottish Smoke-Free Legislation on Occupational Exposure." Annals of Occupational Hygiene **51**(7): 571-580.

Semple, S., H. Sweeting, E. Demou, G. Logan, R. O'Donnell and K. Hunt (2017). "Characterising the Exposure of Prison Staff to Second-Hand Tobacco Smoke." Annals of Work Exposures and Health **61**(7): 809-821.

Shaw, D. (2015). "Prison smoking ban could cause 'stability issues'. Banning smoking in prisons in England and Wales could make them more unstable, the Prison Governors Association (PGA) has warned.", from <http://www.bbc.co.uk/news/uk-33617492>.

Sweeting, H. and K. Hunt (2015). Evidence on smoking and smoking restrictions in prisons. A scoping review for Scottish Prison Service's Tobacco Strategy Group.

MRC/CSO Social and Public Health Sciences Unit Occasional paper no. 25. Glasgow, MRC/CSO Social and Public Health Sciences Unit.

Taylor, P., C. Ogden and K. Corteen (2012). " Tobacco Smoking and Incarceration: Expanding the 'Last Poor Smoker' Thesis. ." Internet Journal of Criminology. : 1-19.

Vijayaraghavan, M., K. Messer, M. White and J. Pierce (2013). "The Effectiveness of Cigarette Price and Smoke-Free Homes on Low-Income Smokers in the United States." Am J Public Health **103**: 2276-2283.

World Health Organisation (2007). Protection from exposure to second hand smoke:: policy recommendations. Geneva, World Health Organisation.

World Health Organization Regional Office for Europe (2013). Good governance for prison health in the 21st century: a policy brief on the organization of prison health. Copenhagen, The WHO Regional Office for Europe.

Figure 1a: Revised overview of workpackages in light of known date for implementation of smokefree prisons in Scotland




Figure 1b: Overview of workpackages as in original funding proposal

