



PROTOCOL

Evaluating the integration of the Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) into primary care and its impact on patient treatment and care.

Sponsor:	University of Warwick
Funding Body:	NIHR HS & DR Programme 13/13/16
Ethics Approval date:	London South East Research Ethics Committee REC ref 21/LO/0455 17.6.21
Confidentiality Advisory Group Approval date	21.7.21 (CAG ref: 21/CAG/0089)
Version Number:	V1.3
Date:	21.7.2021

Protocol Amendments:

Amendment No.	Date of Amendment	Date of Approval
---------------	-------------------	------------------

CONTACT NAMES AND NUMBERS

Sponsor:	Ms Carole Harris University of Warwick, Research & Impact Services University House Kirby Corner Road, Coventry, CV4 8UW Tel: 024 765 22746 Email: sponsorship@warwick.ac.uk
Chief Investigator:	Professor Anne-Marie Slowther Division of Health Sciences Warwick Medical School University of Warwick Coventry CV4 7AL Tel: 024 765 74513 Email: a-m.slowther@warwick.ac.uk
Study contact:	Project manager Division of Health Sciences Warwick Medical School Gibbet Hill Campus Coventry CV4 7AL Email: respectpc@warwick.ac.uk
Co-investigators:	Professor Frances Griffiths, University of Warwick Professor Martin Underwood, University of Warwick Professor Gavin Perkins, University of Warwick Professor Jeremy Dale, University of Warwick Professor Paramjit Gill, University of Warwick Dr Claire Hawkes, University of Warwick Dr Rachel Spencer, University of Warwick Dr Hazel Blanchard, GP Forrest Medical Centre, Coventry Dr Chris Turner, University Hospitals Coventry and Warwickshire NHS Trust Chris Bain, PPI representative

Statistician

Dr Angela Nouifally
Division of Health Sciences
University of Warwick
Email: A.Noufaily@warwick.ac.uk

Study Steering Committee:

Chair: Dr Victoria Tzortziou Brown, OBE
Institute of Population Health Sciences
Queen Mary, University of London, Barts and the London School of
Medicine and Dentistry
Email: v.tzortzioubrown@qmul.ac.uk

Dr Juliet Spiller Consultant in palliative medicine, Marie Curie
Hospice, Edinburgh

Dr Sarah Shaw Associate Professor, Nuffield Department of
Primary Care Health Sciences, University of Oxford

Ms Najma Hafeez PPI representative

Mr Nick Darlington PPI representative

Ms Angela Johnson, Manager, Weavers Care Home, Coventry

Mr James Cant, Chief Executive Resuscitation Council UK

For general queries please contact the study team:

Project manager

Division of Health Sciences

The University of Warwick

Gibbet Hill Road

Coventry

CV4 7AL

Email: respectpc@warwick.ac.uk

TABLE OF CONTENTS

PAGE

	TABLE OF CONTENTS	4
	LIST OF ABBREVIATIONS/GLOSSARY	7
1.	BACKGROUND	9
1.1	Existing knowledge	11
1.2	Need for this research	14
1.3	Ethical considerations	14
1.4	Data Collection	17
1.5	Data Management	20
1.5.1	Data storage	20
1.5.2	Data access and quality assurance	20
1.5.3	Archiving	20
1.6	End of the Evaluation Study	20
2.	STUDY SUMMARY	21
2.1	Aims and objectives	21
2.2	Design and theoretical/conceptual framework	22
3.	WORK PACKAGE 1: EXPLORING THE EXPERIENCES OF PATIENTS, CARERS AND CLINICIANS	23
3.1	Research questions	23
3.2	Setting	23
3.3	Recruitment of GP practices and care homes	23
3.4	Recruitment and consent: individual interviews	24
3.4.1	Patient and family/carer interviews	24
3.4.2	GPs interviews and fieldwork conversations with practice staff	26
3.4.3	Nursing and care home manager interviews	26
3.5	Consent process	27
3.6	Withdrawals	27
3.7	Data collection	27
3.7.1	Describing patient characteristics	27
3.7.2	Interview process	28
3.7.3	Patient and family/carer interviews	28
3.7.4	Interviews with GPs and conversations with practice staff	29
3.7.5	Nursing and care home managers	30
3.8	Data Management	30

3.9	Analysis	31
4.	WP TWO: UNDERSTANDING THE WIDER CONTEXT OF RESPECT IMPLEMENTATION (OBJECTIVE 2)	31
4.1	Research question	32
4.2	Focus groups and Interviews	32
4.2.1	Methods.....	32
4.2.2	Recruitment	32
4.2.3	Consent	33
4.2.4	Withdrawals.....	33
4.2.5	Data collection	33
4.2.6	Data Management	33
4.2.7	Analysis	34
4.3	National surveys of public and GPs	34
4.3.1	Method	34
4.3.2	Sample and data collection	34
4.3.3	Data Management	34
4.3.4	Sample size and analysis.....	35
5.	WP THREE. INVESTIGATING CONGRUENCE BETWEEN RESPECT RECOMMENDATIONS AND TREATMENT DECISIONS: AN ANALYSIS OF PATIENT RECORDS (OBJECTIVE 4).....	35
5.1	Research question	35
5.2	Method	36
5.3	Recruitment and consent	36
5.4	Withdrawals.....	36
5.5	Data collection	37
5.6	Data Management.....	37
5.7	Sample size	37
5.8	Analysis	37
6.	WP4 EVIDENCE SYNTHESIS AND STAKEHOLDER CONFERENCE (OBJECTIVES 3 AND 5).....	38
7.	STUDY ORGANISATION AND OVERSIGHT	39
7.1	Sponsor and governance arrangements.....	39
7.2	Ethical approval	39
7.3	Trial Registration.....	40
7.4	Indemnity.....	40
7.5	Project timetable and milestones.....	40
7.6	Administration	41
7.7	Study Management Group (SMG)	41
7.8	Study Steering Committee (SSC).....	41

7.9	Data Monitoring Committee (DMC)	41
7.10	Essential Documentation.....	41
7.11	Financial Support	41
8.	MONITORING AND QA.....	42
9.	PPI INVOLVEMENT	42
10.	DISSEMINATION AND PUBLICATION	42
11.	REFERENCES.....	44

LIST OF TABLES	PAGE
Table 1 Summary of the main approaches by each work package	18
Table 2 Options for data collection depending on COVID-19 guidance	19

LIST OF FIGURES	PAGE
Figure 1 Study Flow Diagram	22
Figure 2 Plan of investigation and timetable	40

LIST OF ABBREVIATIONS/GLOSSARY

Abbreviation	Explanation
ACP	Advance Care Planning
BMA	British Medical Association
BSA	British Social Attitudes Survey
CAG	Confidentiality Advisory Group
CCG	Clinical Commissioning Group
CI	Chief Investigator
CPR	CardioPulmonary Resuscitation
CQC	Care Quality Commission
CRN	Clinical Research Network
DMC	Data Monitoring Committee
DNACPR	Do Not Attempt CardioPulmonary Resuscitation
ECTP	Emergency Care Treatment Plans
ED	Emergency Department
GCP	Good Clinical Practice
GP	General Practitioner
GDPR	General Data Protection Regulation
HRA	Health Research Authority
ICF	Informed Consent Form
ICU	Intensive Care Unit
IRAS	Integrated Research Application System
NICE	National Institute for Health and Care Excellence
NIHR	National Institute for Health Research
NPT	Normalisation Process Theory
PCN	Primary Care Network
PI	Principal Investigator
PPI	Patient & Public Involvement
POLST	Physicians Orders for Life Sustaining Treatment
RCGP	Royal College of General Practitioners
RCUK	Resuscitation Council UK
REC	Research Ethics Committee
ReSPECT	Recommended Summary Plan for Emergency Care and Treatment

R&D	Research and Development
SMG	Study Management Group
SOP	Standard Operating Procedure
SSC	Study Steering Committee
TEP	Treatment Escalation Plan
WCTU	Warwick Clinical Trials Unit
WMS	Warwick Medical School
WP	Work Package

1. BACKGROUND

When a person develops a life threatening condition or has a sudden deterioration in an existing illness rapid treatment decisions are needed. Often there is limited clinical information or information on what the individual's treatment preferences might be. Anticipatory treatment decisions and recommendations may improve this decision making process. Do Not Attempt CardioPulmonary Resuscitation (DNACPR) decisions are the commonest form of anticipatory treatment decisions but concerns exist about their use including:

- failure to consider CPR decisions before a patient deteriorates,
- lack of discussion with patients and/or their family,
- lack of transferability of forms between primary and secondary care and ambulance services(1,2).

A key concern is that a focus on CPR leads to lack of consideration of other treatments that may, or may not, be appropriate(3,4,5).

A more holistic approach to anticipatory decision-making for future treatments is captured in the concept of advance care planning. Advance care planning, defined as a process of formal decision making that aims to help people establish decisions about future care that take effect when they lose capacity(6), is recommended in NICE guidance(7) and in the UK national Gold Standards Framework for end of life care(8). While historically advance care planning has been seen as important for people approaching the end of their life, the Gold Standards Framework website notes that advance care planning can support people at any age or stage of health in understanding and sharing their personal values, life goals, and preferences regarding future medical care(8). While advance care plans include recommendations about specific treatments including CPR they have a broader scope. They can include preferences on place of death and other non-treatment aspects of care. Advance care plans are typically directed by the person although supported by, and often prompted by, health care professionals caring for them. Lund, in 2015, described two broad theoretical approaches to advance care planning,

1. a phenomenological or psychosocial approach that sees advance care plans as patient driven, drafted in the patient's own environment and focuses on broad goals, and
2. a pragmatic or organisationally oriented approach that follows a trigger event, focusses on identifying specific treatment goals in the face of death and is documented and shared(9).

Emergency Care Treatment plans, sometimes called Treatment Escalation Plans, focus on treatment decisions in emergency or acute illness situations. Their aim is to make treatment recommendations that reflect the person's preferences and values, and are reached in discussion with the person or their family. However, the specific recommendations are made by the health care professional and are intended to guide future treating clinicians(10). Emergency care treatment plans would be conceptualised within the pragmatic approach described by Lund et al (9).

Another important distinction between advance care plans and emergency care treatment plans is that advance care plans ultimate focus is on end of life care preferences, an emergency care treatment plans focusses on steps to be taken in the event of an acute pathophysiological deterioration in which recovery is possible although may be unlikely(11). In the USA the Physician's Orders for Life Sustaining

Treatment (POLST) has been adopted in 47 States(12). In the UK examples include a TEP used across primary and secondary care in Devon(13), and an Emergency Health Care Plan as part of the Deciding Right initiative currently active in North East England(14).

In 2016, the Resuscitation Council UK (RCUK) led development of a new approach to support conversations about goals of care, and to provide guidance to clinicians about treatments, to be recommended in an urgent situation when the patient lacks capacity to decide for themselves (9). This initiative was in response to a growing concern regarding the use of DNACPR recommendations and informed by our evidence review(15).

Initially rolled out in Acute Trusts the Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) has now been adopted for use in primary care by several Clinical Commissioning Groups. Seventy percent of counties in England have now fully or partially adopted ReSPECT across primary and secondary care or are working towards adoption, and there is strong support for its uptake in primary care (personal communication). Initial adoption occurred in Acute NHS Trusts. We have just completed an evaluation of its use in acute NHS Trusts (16). However, there may be advantages to having these conversations in primary care. For example patients may have an established relationship with their GP, conversations can occur over an extended period, patients are less sick and more able to engage in discussion, and conversations can be placed in a wider context of advance care planning. A consultant participating in our current study commented to this effect:

“this is quite a serious and significant discussion ... it should either happen when the person is comfortable in their own home or they have gone to see their GP” (17).

However, there are also potential difficulties in moving ReSPECT conversations to primary care; patients and families may be less ready to think about these things until a crisis emerges, GPs may be uncertain about hospital-based interventions, and both may have concerns about the effect of a conversation on the patient doctor relationship. The ReSPECT process aims to support patient involvement in decisions about their care but these conversations may also lead to distress or uncertainty for patients and their family(18). If forms are not completed and reviewed with diligence future treatment decisions may lead to less rather than more appropriate treatment.

The COVID-19 pandemic has precipitated an increased focus on the use of emergency care treatment plans in general and ReSPECT in particular. The high mortality in older frail people and in those with specific co-morbidities, few effective treatments, and restrictions on hospital visiting, focussed attention on the need for discussions about what treatments they would, or would not, want, or would be recommended should they develop COVID. GPs and hospital doctors admitting acutely ill patients were encouraged to have these conversations with patients early and to document the resulting treatment recommendations. This renewed emphasis on anticipatory decision-making was seen as supporting best practice in person centred care, and an opportunity to embed the principles and practice of advance care planning and emergency care treatment planning into clinical practice more generally (19,20). However, there were concerns that the method of implementation might lead to inappropriate use and resulting poor patient care. Doctors were expected to document conversations and treatment recommendations, including decisions about resuscitation and admission to hospital, when face to face contact between GPs and patients or their families was restricted. Examples of DNACPR and ReSPECT forms being completed without a conversation with the patient or their family were reported in the press leading to public concern (21) and statements from professional and

regulatory organisations (22,23). The Care Quality Commission's interim report into its review of Do Not Attempt Cardiopulmonary Resuscitation decisions during the COVID-19 pandemic noted there was confusion and miscommunication about the application of DNACPR decisions, with some evidence of unacceptable and inappropriate DNACPRs being made at the start of the pandemic. It also identified differing views on the extent to which people are now experiencing positive person-centred care and support in relation to this issue(24).The RCGP and BMA have recognised the challenges for GPs in having these conversations during a pandemic and have provided guidance to its members reiterating the requirement for individualised conversations about future treatment decisions (25,26). It is not clear what impact this experience, or the findings of the CQC's final report, will have on the use of advance care plans and emergency care treatment plans, including ReSPECT as the pandemic recedes.

GPs were being encouraged to use ReSPECT prior to COVID and this has accelerated during the pandemic. However, the rapid implementation under pressure of the pandemic risks inappropriate use which may have implications for future implementation. There is therefore a pressing need to explore how it currently does or does not work in primary care, the impact of the pandemic on its implementation, and implications for patients, their families and health care professionals.

1.1 Existing knowledge

In a scoping review to identify studies evaluating implementation of emergency care treatment plans including barriers to implementation, patient and provider experiences, and outcomes we identified four UK studies, three reporting on treatment escalation plans in single hospital sites (27,28,29). All reported that the treatment escalation plan increased appropriate decision-making around treatment escalation and resuscitation, although appropriate was not defined. TEPS were positively evaluated by staff. An evaluation of the first ReSPECT pilot in Scotland included 200 ReSPECT forms completed in a range of settings (hospital (83%), hospice and community), The study found that patients with a ReSPECT form were more likely to die in their preferred place of care and had a reduced chance of readmission within 3 months of hospital discharge. Patients and family (n=15) felt that discussions were open, honest and that they were involved. Staff (n=20, eight with experience of ReSPECT) were generally positive about the process (qualitative responses) (30). Challenges included time pressures, staff reluctance to initiate a conversation or lack of confidence to do so. A study currently being conducted by NIHR Applied Research Collaboration (ARC) West is evaluating use of ReSPECT in nursing homes(31). We are currently completing the first comprehensive evaluation of the ReSPECT process in secondary care(16). We have reported preliminary findings from interviews with secondary care consultants' in two acute NHS Trusts in England. We found that uncertainty about prognosis, constraints of time and external environment, and the need to minimise patient distress, influence prioritisation and content of conversations (17).

All but one of the other identified studies in our literature review related to the Physicians Orders for Life Sustaining Treatment (POLST) intervention or versions thereof and were conducted in North America. A 2015 systematic review of studies of the POLST programme (N=23) found that POLST was most commonly used in older white people who are near the end of life and that clinicians were generally positive about its use. However, they reported a wide range of challenges including lack of knowledge and training, discomfort with discussing issues raised, and problems using it to guide treatment (32). There was little evidence on whether POLST reflects patient or surrogate treatment preferences. The authors called for research to explore patients' and families' experiences of POLST

and assess its effect on care outcomes. We identified 36 further studies published since this review. Two identified barriers to implementation of POLST at the systemic and individual level including

- lack of patient knowledge,
- reluctance to discuss end of life, uncertainty about prognosis,
- lack of time and concern about the physician patient/family relationship,
- lack of infrastructure and
- legislative concerns (33,34).

Successful implementation was facilitated by standardised protocols or decision aids for conversations (35,36,37,38,39,40). Possession of a POLST form specifying limited or comfort level care was associated with reduced likelihood of admission to ICU or days spent in ICU although data were variable depending on reason for hospital admission and underlying medical condition(41,42,43). POLST was also associated with increased referrals to hospice and out of hospital death in people with cancer and Parkinsons Disease (44,45,46). Studies looking at POLST in nursing homes or elderly care facilities found a high level of consistency between POLST recommendations and subsequent treatment decisions(42,47,48,49).

As the literature on POLST is almost exclusively from North America and does not include primary care we considered wider literature on anticipatory decision making. Advance care planning has many synergies with ECTPs particularly in primary care and there is a large international literature on ACP, including some studies in primary care. We therefore reviewed this literature using as a starting point a 2018 overview of systematic reviews that included 80 reviews and over 1600 studies of ACP(52). We then searched PubMed for systematic reviews published since 2016 (cut-off date for the 2018 review) using the search terms advance care plan* AND implementation OR barrier* OR facilitator* OR outcome*. We identified a further 34 reviews. The 2018 review overview found some evidence of benefits including documenting of care preferences, dying in preferred place and health care savings, and that facilitators of effective ACP include provision of information by a knowledgeable person, transfer of ACPs across healthcare settings and moving ACP from a hospital to community setting(50).

Of the 39 reviews identified since 2016, 15 focussed on ACP in specific diseases including cancer, dementia, COPD and heart failure. Key messages from all reviews were similar to those found by Jiminez (50). A holistic, culturally sensitive and systems wide approach was seen as necessary for effective ACP (51,52,53,54) with education of both patients and health care professionals(55,56). Patients and families value honesty and openness in conversations (57,58,59) but there is a reluctance among health care professionals and patients to talk about end of life (60). ACP was felt to bring benefits including a greater sense of peace and less worry, but it could also be disruptive and distressing (18,58,61)). ACP was found to be associated with dying in their preferred place (59), reduction in hospitalisation rates and length of stay and invasive treatment and reduced health care costs (62,63,64). Three reviews looked specifically at culture and ethnicity in relation to ACP (65,66,67). Cultural factors reported as affecting ACP acceptability included religiosity, spirituality, trust in the health care system, social networks, and patient attitudes regarding decision-making. There was variable methodological quality of studies reported by most reviews. One review specifically looked at methodological quality of studies investigating concordance between ACP and end of life care. The proportion of patients who received concordant care varied from 14% to 98% but studies were methodologically poor (68).

We found two systematic reviews that focussed specifically on ACP and general practice (DeVlemenik 2013, (n=15)(69) Risk 2019 (n=54)(70). The 2019 review used an ecosociological framework to classify barriers and enablers to ACP's uptake at the individual, interpersonal, provider and system level. Key barriers identified reflected previous review findings and included lack of patient knowledge and trust, confusion over whose role it was to initiate the conversation, concern over impact in the patient doctor relationship, GP's lack of confidence and training, and lack of organisational support including mechanisms for sharing ACPs across settings. The review also noted the paucity of research on ACP general practice and very few studies included the patient's voice. A 2016 scoping review of implementation studies of ACP in nursing homes (n=16) found variation in implementation strategies and outcomes with barriers to implementation including absence of physicians in the process, reluctance of staff, residents and families to initiate or participate in discussions and legal issues (71).

The literature on ACPs and POLST identified structured tools and protocols as a key enabler for implementation. The ReSPECT process and form, with the supporting educational materials provided on the ReSPECT website, could be characterised as such. However, a review that specifically looked at implementation studies of ACP using Normalisation Process Theory as an analytical framework concluded that structured tools were unlikely to be sufficient (10). The authors suggested that interventions most likely to be successful will need to make ACP workable within complex and time pressured clinical workflows. They developed four propositions about ACP normalisation which are relevant when considering normalisation of ReSPECT;

- Clinical and organizational pressures affect opportunities to initiate and operationalize ACPs.
- Prognostic uncertainty affects clinical decisions to initiate ACP conversations
- Responses of patients' and their families to initiation of conversations are unpredictable and emotionally complex
- Clinical and organisational factors intervene to confound elicited plans and preferences

Findings from our current evaluation of ReSPECT in secondary care support the first three propositions. Propositions two and three in this framework reflect the complexity, uncertainty and emotive nature of ReSPECT conversations. These kinds of conversations rely on a relationship of trust between the patient, their family and the health care professional. The literature on trust suggests that patient trust for their clinician has two elements, trust in their competence and trust that they have the patient's best interests at heart (72). Trust is necessary in situations of uncertainty, and anticipatory treatment decisions have a higher level of uncertainty than contemporaneous decisions as potential events and outcomes are in the future. In addition, at the time the recommendations take effect the patient will not be in a position to contribute to or modify decisions made. The limited empirical evidence on patient and family experiences of ACP suggest that they value honesty and transparency (57,58,659), seen as facilitators of trust, and studies focusing on culture and ethnicity in relation to ACP notes trust in the health care system as a factor in acceptance of ACP (65,66,67). There is mixed evidence on whether continuity of care facilitates development of trust in the patient doctor relationship. It will be important in the proposed study to explore how trust of patients' and their family in clinicians and in the health care system influences the ReSPECT process.

The policy aim for ReSPECT is that it should be normalised into everyday clinical practice. The proposed project will therefore evaluate the use of ReSPECT in primary care in relation to the Normalisation Process Theory framework. In doing so it will specifically consider the interactions between individual

and system levels and include the patient voice, both identified as gaps in the systematic review of ACP in primary care (70).

1.2 Need for this research

From 2015 to 2035 there will be a >150% increase in number of older people with complex multi-morbidity (73). These people are particularly susceptible to sudden significant deterioration in their health resulting in emergency treatments and admission to hospital. Many will be unable to make a decision regarding treatment at the time of the emergency. Half of the increase in emergency admissions between 2013/14 and 2016/17 came from people aged 65 and over, particularly those who are frail (74). The 2.8% of the older population who are care home residents account 7.9% of all emergency admissions for people aged 65 or older (75). In 2016 there were over 1.6 million emergency admissions in the UK for people in the last year of their life (an 8% rise since 2011) costing the NHS £2.5 billion and amounting to around 11 million days in hospital (76). While some emergency admissions in the last year of life are necessary most people prefer to die at home (77,78). Decisions around admission to intensive care are often made with no knowledge of the patient's values or wishes and may be influenced by a range of non-clinical factors leading to over or under treatment (79).

There is an ethical and professional obligation to balance benefits and burdens of treatment from the patient perspective(80) Anticipatory decision making using 'Emergency Care Treatment Plans' (ECTPs) can facilitate person-centred shared decision-making in the acute situation by enabling considered assessment of the benefits and burdens of a range of treatments taking into account what is important to the patient. If the ReSPECT initiative is effective it should improve patient care, fit with current policy on personalised care(81) and allow more effective use of NHS resources.

GPs who took part in focus groups for our evaluation of ReSPECT in Acute Trusts described a willingness to engage in the ReSPECT process, recognised that conversations should be embedded in clinical practice but suggested practical and cultural challenges to achieving this. They agreed evaluation of its use in primary care was needed. An early evaluation of its adoption across primary care, potential enablers and obstacles to implementation, and impact on patient care will provide evidence to inform successful and sustainable implementation, ensuring that potential benefits to patients and the health service are realised.

1.3 Ethical considerations

This study will be conducted in accordance with the NHS Research Governance Framework and the principles of Good Clinical Practice (GCP). As the study may involve some patients who by the nature of their underlying illness lack capacity, the requirements of the Mental Capacity Act (2005) will be observed. The study will comply with relevant Warwick Standard Operating Procedures (SOPs) and all data will be stored securely and held in accordance with Data Protection Act 2018.

This study raises a number of ethical issues given the sensitivity of the subject under investigation. The main ethical issues we have identified are as follows:

1.3.1 Recruitment and conduct of Interviews with patients and family members

ReSPECT discussions are inherently sensitive and emotive conversations. People who have a ReSPECT form completed may often be coming towards the end of their life and any ReSPECT conversation will be precipitated by an acknowledgement that an acute deterioration in health from which recovery

may be unlikely is anticipated. Recruiting participants and conducting interviews will need to be approached with sensitivity. We have identified processes to avoid approaching a patient or member of their family who is in the last stages of life and to identify if a relative is bereaved prior to a follow up interview. The senior research fellow who will conduct the interviews will have either experience of or receive training in conducting sensitive interviews with patients and families and in assessing capacity to consent. Interviews can be stopped at any time should the participant wish to do so or becomes distressed. When a patient lacks capacity to consent to access to their ReSPECT form and medical records we will seek personal consultee agreement from their next of kin.

It is possible that a patient will be identified who had a ReSPECT form completed in hospital and may be unaware or have forgotten that this occurred. This could be distressing if they first become aware or re-aware of a ReSPECT form through a letter about a research study. We will where possible purposively select patients who have had a ReSPECT form completed in primary care to make this less likely. We will also develop a communication plan to ensure that any enquiries to the study team regarding an unknown ReSPECT form completion are responded to sensitively and the patient or their carer directed to the GP for further discussion.

1.3.2 Responding to distress of patients and family members during interviews

Interviews can be stopped at any time should the participant wish to do so or if they become distressed. Participants will be known to the health care professionals within the GP practice who can provide further support should the participant require it. We will also provide contact details for local support services identified through the practice. For bereaved participants we will offer contact details of local and national support services. If the researcher is concerned about a significant harm for an individual participant, for example if the participant discloses a suicidal intention, the researcher will discuss this immediately with either the WP lead or CI who will take action to ensure appropriate support is provided, probably through their GP. WMS has a standard operating procedure for responding to disclosures of this nature by research participants, which we will follow.

1.3.3 Concerns about unprofessional practice or safeguarding arising in the interview

If concern arises about unethical or unsafe clinical practice the researcher will consult the work package lead or CI who will decide if it is necessary to initiate action through normal professional channels, which is likely to be through the relevant GP practice. We think it is unlikely that serious unprofessional conduct will be observed or disclosed in this study. If any disclosures are made the participant (GP, care home staff, patient or family member) involved in the interview where this practice was revealed will be informed that this is happening. The need for a researcher to disclose any evidence of serious professional misconduct or safeguarding concern will be made clear in the relevant participant information sheets.

1.3.4 researcher safety during lone working

Some patient and family interviews may be conducted by the senior research fellow working alone visiting patient's homes. Our process for managing these events will comply with the University of Warwick lone worker guidance and the University of Warwick SOP for risk assessment and monitoring in research studies which specifies conducting a risk assessment and the process for ensuring a 'buddy' system is in place for any lone worker activity. A named person will be the contact

for the researcher when conducting research interviews as a lone worker, with agreed times for contact and steps to take if contact is not made.

1.3.5 verbal consent for informal conversations with practice staff (WP1)

When the researcher initiates or participates in informal conversations with practice staff members, the researcher will obtain and document verbal consent from the participating staff members to write up these conversations, or any part thereof (in quoted or paraphrased form), as field notes, to be used in data analysis. Obtaining verbal consent would allow us to maintain continuous transparency about the fieldwork process and ensure that practice staff members consent to each informal conversation, while minimising disruption to the informal conversation and their routine work. To ensure that participants can easily contact the researcher to withdraw their consent, the researcher will give their University of Warwick business card, which includes their email address and other contact information, to each clinical staff member who participates in an informal conversation.

1.3.6 Accessing medical records and ReSPECT forms (WP1 & 3)

We are seeking approval from the Confidentiality Advisory Group to use section 251 of the NHS Act 2006 to access medical records. Patients (and carers of patients who lack capacity) will be given the opportunity to opt out of this part of the study by the provision of an information leaflet explaining the study. We present the potential risks, mitigations and benefits of this approach below.

The main risk for copying and extracting data from medical records relates to a breach of trust / confidentiality through access to clinical records. We are mitigating the risk by (1) only reviewing sections of the record relevant to the research question (2) collecting the minimum amount of data to address this research question (3) anonymising copied/extracted data (4) making use of routine audit data where possible (5) ensuring staff collecting the data will have a duty of confidentiality through a contract with the hosting GP practice or care home. It is possible (although unlikely, given the existing duty of clinicians to consult patients), that an awareness of this research activity may prompt patients to raise questions about their overall treatment.

The direct benefits for individual research participants are limited to raising a general awareness about ReSPECT amongst clinical staff and patients. There will also be benefits for future patients through a better understanding of how ReSPECT is working in the NHS and social care.

Considering the risks, mitigations and benefits we assess the overall risks from this as low to negligible.

The research requires the research team to access the following information from the patients' clinical records

- i) Information recorded on the ReSPECT form
- ii) Clinical justification for a ReSPECT recommendation
- iii) General information about the patient (full information is provided in section 5.3). The types of information required is e.g. demographic information, frailty index if recorded and

treatment/management decisions for an episode of significant illness within six months of the ReSPECT form completion.

Information will be copied or extracted from the ReSPECT form and GP medical records by NHS practice or research staff.

NHS staff or research nurse/researcher with a research passport/honorary contract with the practice will anonymise the data set before it is securely returned to the central research team at Warwick University for analysis.

A record of who the participant identification numbers have been allocated to will be kept at the research site in order to enable withdrawal of participants after the data has been collected prior to analysis. In the event a patient requests their data not to be included after it has been collected but before the data is analysed, we will treat this as if it were a withdrawal and their data will be removed from the data set.

Our approach seeks to balance respect for the patients right to information in their medical record being treated confidentially, a public interest in obtaining an unbiased sample to achieve a valid research outcome and consideration of practicable alternatives to obtaining consent. This part of the data collection is the subject of our application to the Confidential Advisory Group (CAG).

We consider the use of assumed consent model (sometimes called opt-out consent) as proportionate to (1) the level of risk involved (2) the likelihood of a biased sample with patients who are more likely to have a ReSPECT form completed and acted upon because of significant illness less likely to complete the consent process, (3) the burden to patients / relatives of going through a process of written informed consent.

Information leaflets about the study will be sent to patients / relatives with an accompanying letter from the GP practice. The leaflet will include information about the study, what information is being collected, that it will be anonymised before leaving the NHS site and securely transferred to the central research team for analysis. It will also include contact details for further information and how an individual can opt out of the study. The opt-out approach was developed with patient and public partners and implemented successfully in our previous study evaluating ReSPECT in acute NHS Trusts. previous studies (IRAS Project ID: 204688).

Practice level consent

Practice level approval for participation in the study (and sharing of documents relevant to the system approach to implementing respect) will be provided through site specific approval processes.

1.4 Data Collection

Several modalities will be used for the collection of research data across the different work packages (Table 1). Full information is provided in the detailed description covering each work package.

Table 1 Summary of the main approaches by each work package

	Interviews	Case note review	Survey	Focus group	Audit data
WP 1	X				X
WP 2	X		X	X	
WP 3		X			

Confidentiality: Any researcher(s) from the study research team needing access to patient records to support the data collection at sites will apply for a research passport/letter of access. When reporting the findings of the study, participants (GP practices, care homes, patients and relatives, practice staff, focus group participants) who consent or chose not to opt out of the medical case note review will be assigned a unique participant identification number. All results and findings reported will be anonymised, to ensure no individuals can be identified in the study.

Options for data collection depending on COVID-19 risks and regulations.

We plan to conduct all research activity remotely where possible and where we do not think it impacts on the quality of the research. We have identified two areas where face to face contact between the researcher and others conducting the research (practice and care home staff) or between the researcher and interview participants, may in some circumstances be desirable and where only virtual contact might significantly affect the research quality and opportunity for some people to participate. Therefore, we propose the following approach:

1. **Participant interviews:** Most interviews will be virtual. When government guidance permits meetings between people from different households, we will consider offering face to face interviews as an option to participants. For patients and families, participants will vary in terms of COVID risk status, problems with hearing, vision and dexterity, IT literacy, access to devices and for care home residents, current arrangements for visits. When arranging the interview, we will negotiate with each participant the most appropriate way of conducting the interview. If a participant chooses face to face interview, and this would be compatible with government guidance at that time, we will offer this.
2. **Researcher presence in the practice to support notes extraction and informal conversations with staff:** When government guidance and local GP practice/care home protocols permit entrance of non-practice/non care home individuals to be present for non-essential activities unrelated to clinical/care, we will negotiate with the practice/care home about researcher presence on the site for limited research activity.

Where an activity is face-to-face current public health guidance at that time will be followed e.g. social distancing, hand sanitiser, face mask/visor, cleaning of surfaces and any recording equipment, researcher undertaking lateral flow tests twice weekly. Table 2 shows research activities which may be considered for a face to face approach (in red) and which will be conducted virtually throughout the study, depending on current government guidance at the time.

Table 2 Options for data collection depending on COVID-19 guidance

Research activity	Government guidance permits contact for non- essential activities	Government guidance restricts contact for non- essential activities
GP practice identification	Telephone/secure email	Telephone/secure email
Presentation to potential participant GP practices	Researcher joins a practice meeting via whatever channel currently in use by the practice (face-to-face/video-conference/teleconference)	Researcher joins practice meeting via video-conference or tele-conference
Identify care homes and practice protocols	Telephone/secure email	Telephone/secure email
PPI activity	Video-conference/email	Video-conference/email
GP patient register search	Practice staff using normal precautions for clinical practice Researcher in practice to support	Practice staff using normal precautions for clinical practice Researcher remote support
Interviews with patient/carers living at home	Face to face inside or in secluded private garden OR Video-conference/telephone	Video-conference/telephone
Interviews with patients/carers living in nursing and care home	Face to face in line with care home protocols OR Video-conference/telephone	Video-conference/telephone
Interviews with GPs, nursing and care home manager, faith leaders	Face to face OR Video-conference/telephone	Video-conference/telephone
Conversations with practice staff	Face to face	Telephone
Focus groups	All participants join via video-conferencing	All participants join via video-conferencing
Copying and deidentifying patient records and sending securely to study team	Practice staff using normal precautions for clinical practice	Practice staff using normal precautions for clinical practice
Collecting ReSPECT form copies from care and nursing homes	Research team requests a member of staff of care/nursing home copy respect forms and send to practice	Research team requests a member of staff of care/nursing home copy respect forms and send to practice
Stakeholder conference	Presentations including touch point film/audio recording will be available to participants to view prior to the meeting. Feedback from small groups to all participants will be via written summaries agreed by small groups and posted for all participants to view and post comments.	
	All participants join via video-conferencing	All participants join via video-conferencing

1.5 Data Management

Data collected during the study will be handled and stored in accordance with the 2018 Data Protection Act and Warwick Clinical Trial Unit Standard Operating Procedures.

No personal identifiable data will be transferred between GP sites and the University of Warwick. The detailed data management processes in the descriptions of each work package that follows.

Disclosure of confidential information

If during data collection a participant raises any issue which may jeopardise the safety of the participant, the researcher will follow local safeguarding processes, usually reporting the issue to the GP lead at the practice. Participant information sheets will include information about the disclosure of such information. If the researcher identifies an issue which raises concerns regarding professional misconduct that could result in a significant risk of harm to patients generally the researcher will discuss this with the work package lead. If the work package lead agrees that there is a cause for concern they will inform the GP practice, care home as appropriate in accordance with local policy on raising concerns.

1.5.1 Data storage

All essential documentation and study records will be stored by WMS in conformance with the applicable regulatory requirements and access to stored information will be restricted to authorised personnel. Any paper data forms, field notes, meeting notes, or other documents will be stored in a lockable filing cabinet in a secure room, to which access is restricted to authorised personnel. Electronic data will be stored in a secure area of the computer with access restricted to staff working on the study.

1.5.2 Data access and quality assurance

Study participants will be assigned a unique study identifier. Each site will maintain a confidential and secure list of patient identifiable information (name, date of birth, identification number) for the purposes of audit / quality assurance.

Once the study has been completed the records will be destroyed according to University of Warwick and or local site SOPs. The CI and the study manager (or staff they delegate this role to) will have access to the final study data set from all four work packages. Access requests from both co-investigators and external parties will be considered by the CI. A formal process will be developed by the study team to facilitate such requests and decisions. Any data shared will be anonymised and transferred as per University of Warwick SOPs with data sharing agreements in place.

1.5.3 Archiving

Study documentation and data will be archived for at least ten years after completion of the study. Research sites will archive documentation following their local policies.

1.6 End of the Evaluation Study

The study will officially end on the last day of funding, although dissemination of results will continue beyond that date.

Since this study is not implementing any intervention, it is unlikely to be stopped prematurely, unless funding is ended early. If several or all of the research sites in WP1 and 3 withdraw the ReSPECT process during data collection this could result in these aspects of the study ending prematurely or partially completed, unless replacement sites can be found within the time constraints of the project. The Research Ethics Committee will be notified in writing if the study has been concluded or terminated early.

2. STUDY SUMMARY

2.1 Aims and objectives

This study is a mixed-methods evaluation of the ReSPECT process for adults in primary care to determine how, when, and why it is used, and what effect it has on patient treatment and care.

Objectives

1. To understand how ReSPECT is currently used in primary care from the perspective of patients, their families, clinicians, and care providers
2. To describe the views of patients, the public, and primary and community health care professionals on emergency care treatment plans in general and ReSPECT in particular
3. To identify enablers and obstacles to embedding ReSPECT in primary care practice
4. To explore the impact of ReSPECT on patient treatment decisions
5. To develop a consensus on how ReSPECT should be used in primary care

The study consists of four work packages corresponding to our objectives:

WP1: A qualitative study using interviews with GPs, patients and their families/carers, and managers of care homes to explore how and why ReSPECT conversations occur and recommendations are made, their ethical basis and the experience of patients and their families of the decision-making process.

WP2: Focus groups with health care professionals, and members of patient and community groups to explore their views on the principles and practice of ReSPECT and other forms of anticipatory decision-making. Interviews with faith leaders to explore the extent that the values underpinning the ReSPECT process reflect or are dissonant with specific faith traditions. A national survey of public attitudes to ECTPs and ACP, and a national survey of GPs to explore their experience and views of anticipatory decision-making.

WP3: An analysis of congruence of ReSPECT recommendations and treatment decisions made using an analysis of ReSPECT forms and corresponding patient records.

WP4: A synthesis of key findings from the study and a stakeholder meeting drawing on experience based co-design to identify strategies to support integration of the ReSPECT process to primary care practice.




Figure 1 Study Flow Diagram

2.2 Design and theoretical/conceptual framework

Our overarching theoretical framework is Normalisation Process Theory (NPT). We wish to investigate to what extent ReSPECT is embedded in routine primary care practice and how it is perceived and enacted by health professionals and patients. NPT characterises a set of mechanisms (coherence, cognitive participation, collective action and reflexive monitoring) that influence the embedding of new interventions or processes into clinical practice (82,83).

We will ask how do clinicians, patients and their families:

1. conceptualise ReSPECT (coherence)
2. initiate or engage with the process (cognitive participation)
3. use the process and the documentation (collective action)
4. evaluate the impact of ReSPECT and how it changes behaviour (reflexive monitoring)

NPT has successfully been used in a review of the implementation of ACPs to understand how they are implemented and embedded – normalised – into clinical practice (9). This review developed the following propositions about ACP normalisation:

1. Operational contexts are under pressure. Clinical and organizational pressures and the availability and preparation of staff affect opportunities to initiate and operationalize complex interventions like ACPs.

2. Patient trajectories are uncertain. Prognostic uncertainty is an important factor that affects the clinical decision to initiate discussion of ACPs with patients and their significant others.
3. Negotiations have unpredictable outcomes. Responses of patients and their significant others to the initiation of ACPs are unpredictable and emotionally complex.
4. Advance Care Plans may not be actioned. The operational outcome of ACPs are unpredictable because clinical and organizational factors that intervene to confound elicited plans and preferences.

We will test these propositions during our interviews and focus group work.

3. WORK PACKAGE 1: EXPLORING THE EXPERIENCES OF PATIENTS, CARERS AND CLINICIANS

3.1 Research questions

This interview study will seek to answer the following questions:

1. What patient characteristics are associated with completion of a ReSPECT form in primary care?
2. What are the experiences of patients, their families/carers, GPs and care home managers of being involved in the ReSPECT process?
3. What are the enablers of and obstacles to implementing the ReSPECT process in primary care?

3.2 Setting

Twelve GP practices from across three Clinical Commissioning Group (CCG) areas, or organisational equivalent at the time, in England that have adopted ReSPECT at least 12 months previously. CCGs will be purposively sampled to aim for diversity of geographical location sociodemographic profile, and time since implementation of ReSPECT.

Method

This work package is a qualitative interview study using interviews with GPs, patients and their families/carers, and managers of care homes. We will also collect anonymous data on selected patient demographic data.

3.3 Recruitment of GP practices and care homes

Within each CCG or the equivalent current organisation, we will work with the local primary care clinical research networks (CRNs) and primary care networks (PCNs) to seek expressions of interest from GP practices. We will purposively sample 12 practices from those who express an interest in participation, based on size, geography (rural/urban) ethnic diversity, and socioeconomic status. We will use practice postcodes to select purposively for socioeconomic and ethnicity profile using data from Office for National Statistics. For selected practices, we will present the study to practice staff supported by written information and seek consent from a designated PI for practice participation..

For each of the twelve GP practices, we will seek to identify two care homes where the practice is the primary practice for the home. Care homes will be purposively sampled to include both residential and nursing care homes. We will contact the care homes through the practice and present the study to the

care home staff supported by written information and seek consent from the home manager for participation in the study.

To place our qualitative work in context, in each practice we will identify any practice protocols for initiating/reviewing ReSPECT or other ECTP/advance care planning documents.

3.4 Recruitment and consent: individual interviews

3.4.1 Patient and family/carer interviews

We will recruit up to 48 patient/carers across all sites (12 practices).

Inclusion criteria

- Patients with a ReSPECT form documented in the previous 12 months;
- Age 18 years and over;

Exclusion criteria

- No ReSPECT form recorded;
- Age less than 18 years

From eligible patients (or their carers) who express an interest in participating we will select equally from three groups, which are people:

1. living in a care home
2. living at home, identified as nearing the end of their life (but not in the final stages of a terminal illness)
3. living at home, not identified as nearing the end of life.

All patient or carer facing documentation will be developed in collaboration with our PPI advisory group.

Recruitment through GP practices: A member of practice staff will search the patient register to identify all adults in the practice known to have a ReSPECT form completed in the previous twelve months. We will work with the practice to identify the most appropriate search method. A member of practice staff will check for each person whether they are recorded as lacking capacity and if so whether there is a specified carer or next of kin associated with ReSPECT form completion. The lead GP for ReSPECT/ACP or delegated health professional for the project in the practice will check the list of patients to exclude those who are known to be in the final stages of a terminal illness (expected to die within the next six weeks) or currently in hospital. If a patient record is flagged as the patient having opted out of their data being used for purposes other than direct clinical care the patient will be excluded. People living in participating care homes will not be approached through the practice.

Patients with a ReSPECT form completed in the last six months will be purposively sampled for WP1. An invitation letter with a brief information leaflet will be posted from the practice to identified patients to inform them about the study inviting them to contact the research team if they are interested in taking part. For people whose first language is not English we will offer the opportunity of an interpreter to facilitate the interview. If the person is known to lack capacity in relation to ReSPECT or other health care decisions the person named in relation to the ReSPECT form or the next

of kin/carer recorded in practice records will be sent the information leaflet and letter of invitation. Potential participants will be asked to contact the study team to express interest (post, free phone (text or audio), email) including their contact details and preferred channel of communication with the research team (phone/text/email).

Recruitment through care homes: Participants will also be recruited through two participating care homes linked to the practice. The care home manager will identify any resident registered with the linked practice who has a ReSPECT form and check with the PI in the linked practice that the person is not known to be in the final stages of a terminal illness (expected to die within 6 weeks). For all residents not in this category the care home manager (or delegated staff member) will give a brief information leaflet and expression of interest form either to the resident registered with the practice, or their next of kin if the resident lacks capacity, who has a ReSPECT form. Next of kin will be asked to contact the study team to express interest (post, free phone (text or audio), email) including their contact details and preferred channel of communication with the research team (phone/text/email). We will ask care homes to return expression of interest forms on behalf of their residents (by secure email/post).

Participant selection and consent: Depending on the number of expressions of interest we will purposively sample across practices aiming for diversity of age, gender, main clinical condition, if on end of life care register, usual living arrangements (home/care home with, or without, nursing), place of ReSPECT completion (hospital/GP practice). We estimate that to gain sufficient information power⁽⁸⁴⁾ from second interviews we need to recruit a total of 48 patient/family/carers. This allows for 50% attrition between first and second interviews leaving 8 per Group at second interview.

The researcher will contact those who have expressed an interest in participating to discuss the study and arrange an interview. This will be via their preferred channel of communication or for care homes by phone (with or without care staff support as needed). Prior to the interview, a full information sheet will be sent to the participant. For participants with capacity we will ask them if we can also interview any family member who was involved in the ReSPECT discussion. If they agree, we will provide a separate information sheet and consent form for the family member. If the person is known to lack capacity related to the ReSPECT conversation, the person who took part in the ReSPECT conversation or their next of kin will have been invited. If it becomes clear in the initial discussion with the researcher that the patient lacks capacity to consent to the research the researcher will ascertain if the carer was involved in the ReSPECT form completion and if so seek consent for an interview with the carer. We will, in line with the Mental Capacity Act 2005 section 32, seek personal consultee agreement from the carer/next of kin of any patient who lacks capacity that the person would not object to the carer being interviewed in relation to their ReSPECT form completion.

We will offer patient participants the opportunity to have a family member present for support during the interview. We aim to make this offer to the patient when arranging the interview, so they are able to make their decision independent of their family. We will ask the patient/carer or relevant care home manager to locate, where possible, the ReSPECT form to prompt the patient/carer memory of the ReSPECT process. Consent will be obtained prior to the start of each interview. At the first interview, we will seek permission from the participant to contact them and/or their main carer/next of kin if relevant in six months-time to arrange a second interview.

We will interview each participating patient/carer initially and after 6-9 months. During the second interview we will explore whether their ReSPECT form has been changed in the intervening period, whether they have had any significant illness episodes and if so whether the ReSPECT recommendations were helpful in guiding decisions about their treatment. As many participants may be in the last year of their life some participants may have lost capacity, be in the final stages of their life, or follow-up interviews may be conducted with recently bereaved carers. We will check with the GP before making contact for a second interview and approach this with utmost sensitivity. If a participant has lost capacity we will, in consultation with the GP, consider approaching the carer to see if they would agree to be interviewed. Participants will be reminded of the study and sent a follow up interview information sheet. If they agree to take part in a second interview the researcher will arrange this at a convenient time and location for the participant. Consent will be taken prior to the interview as for the initial interview. We have experience of interviewing bereaved relatives in this process can also be cathartic for the bereaved (85) Participants will receive a shopping voucher (Love2shop voucher) for £20 in payment for their time.

We will give participants the option of spreading the interview over more than one session. This may be a session with several short breaks or, it may be several short sessions held over several days. We want to offer this option as interviews can be tiring for the participant particularly when conducted via telephone or videoconference.

3.4.2 GPs interviews and fieldwork conversations with practice staff

In each practice, we will interview the health professional who completed ReSPECT for patient participants or the GP most closely involved in their care if the form was completed in hospital. An information sheet will be provided and, if the person agrees, a time will be arranged for the interview and consent obtained prior to interview. We will ask the GPs to review the relevant ReSPECT forms (if available), and relevant clinical notes, in advance of the interviews to prompt their memory of the ReSPECT process they carried out.

We will carry out brief conversations (5-10 minutes, not recorded) with other members of the practice team, for example practice nurses, practice manager, receptionist, asking them about their awareness/use of ReSPECT, and how the process runs in the practice. If COVID regulations allow, a researcher will spend two separate days in the practice taking opportunities to speak to practice staff at a time convenient to them. Practice staff will be provided with information leaflets about the study in advance. The researcher will obtain verbal consent and record this in the field notes. Staff will be given the researcher's contact number so they can contact the researcher if they wish to withdraw from the study after the conversation has taken place.

3.4.3 Nursing and care home manager interviews

We will invite the managers (or a senior member of staff) of care homes linked to participating practices to take part in an interview about their general experience of ReSPECT and of how clinical decisions are made for residents with a ReSPECT form. Managers will be sent an invitation letter and information sheet through the GP practice and invited to contact the research team if they are interested in participating. Managers may pass this information to a senior member of staff who is seen as a more appropriate person to provide the relevant perspective during an interview. Those who

express an interest will be contacted by the researcher to discuss the study and arrange an interview. Remote consent (for video or telephone interviews) or written consent (if it is possible to conduct face to face interviews) will be obtained before the interview.

3.5 Consent process

If an interpreter is required for patient or family interviews, we will arrange this through the practice using the NHS contracted interpreter service used for clinical consultations. Written consent will be obtained immediately prior to all face-to-face interviews. For interviews occurring via video or audio call remote verbal consent will be obtained. When the interview time and date is confirmed a consent form will be sent to the participant prior to the interview together with a full information sheet. Before starting the interview, the researcher will answer any questions the participant has, then go through the consent form with the participant. Remote verbal consent will be formally documented by the researcher, including consent to record the interview. The potential participant will be asked to verbally consent to each line on the informed consent form during the video/audio discussion. The researcher will initial and date each line of the consent form and then they will sign and date the completed consent form. The participant will receive a copy of the informed consent form by post or email following the interview.

For conversations with practice staff, the researcher will obtain verbal consent from the participating staff members to write up these conversations, or any part thereof (in quoted or paraphrased form), as field notes, to be used in data analysis. All field notes will anonymise participants and exclude or alter identifying information; pseudonyms will be used throughout, and the staff members' names will not be recorded.

3.6 Withdrawals

Participants can withdraw from the interview study up to three weeks following the interview without prejudice. Beyond three weeks the data will have been transcribed and entered into the analysis process so it will not be possible to extract it from the analysis.

Participants will be given the contact details for the research team who they may contact at any time during this period to inform the team they wish to withdraw consent. Their wishes will be recorded in the Study Master File using their study ID and their data removed from the study.

3.7 Data collection

3.7.1 Describing patient characteristics

We will work with the practice staff to collect pooled anonymous data on;

- practice population demographics
- number of ReSPECT forms recorded
- for people with ReSPECT form: age; ethnicity; eFrailty index (generated from routine practice data); type of residence (care/nursing home or own home/family home) and where ReSPECT form completed (if recorded)

This element of data collection is part of our application to CAG to permit a study researcher to support practice staff.

We will work with individual practices to develop the best identification processes using practice electronic records and other practice systems. A data collection spreadsheet will be completed and held within the practice. Aggregate anonymous data will be transferred securely to the study team at WMS. We will present descriptive statistics on these data.

3.7.2 Interview process

Interview topic guides have been developed with input from our clinical co-investigators and our patient and public advisory group. As we collect data we will review and revise topic guides to ensure we are obtaining the richest possible data whilst remaining sensitive to participants. We will keep the interview open for participants to talk about ReSPECT and advance care planning in whatever order they wish and enable them to talk about issues not covered by our topic guides, whilst aiming to cover all our planned topics. We will audio or video record all interviews with participant consent. Where consent to recording is not given, we will take field notes and expand the field notes directly after the interview. We would like to use brief clips from interview recordings as part of our experience led co-design work at our stakeholder conference. We will seek consent to use clips of interview recordings from participants prior to any interviews. We will check with participants, or next of kin if appropriate, whose recording we wish to use that they are happy with the recording clip prior to using it for the meeting.

3.7.3 Patient and family/carers interviews

First interview: At the start of the interview we will ask if the patient/carer has located the ReSPECT form and if they have, suggest they look at it to remind them of the ReSPECT process. Using the NPT framework to guide our data collection, we will frame our questions to reflect NPT elements. We will ask them:

- What they understand about the ReSPECT process and form (*coherence*)
- Their views on the ReSPECT process, when they consider their ReSPECT form should be reviewed and who should initiate this review, when and how it should be acted on (*cognitive participation*)
- To describe the ReSPECT process from when it was first considered or mentioned through to completion of the form and any updating of the form (*collective action*)
- Thoughts and feelings the process engendered; whether the process changed how they viewed themselves, their health/illness, and the prospect of their death; whether they think it resulted in a change to the care, and treatments they received and how they viewed these; their sense of trust in their clinicians and in the health service more generally including whether they trust that the ReSPECT form will receive attention once completed; how the ReSPECT process could be improved. We will ask whether they have been involved in any other advance care planning discussions and whether COVID-19 has had any impact on their thinking about ReSPECT or advance care planning in general (*Reflexive monitoring*).

Second interview: The researcher conducting the interview will re-read the first interview transcript and make notes to inform their prompts during the second interview. Where a first interview participant has died we will acknowledge this at the start of the interview with their carer. We will ask participants about major changes to the patient's life context and health/illness including any

illness/treatment events since the first interview, including further experience of the ReSPECT process/review. If they have further experience with ReSPECT we will explore this in a similar way to the first interview and prompt for any differences compared to the first interview. Where participants have experienced illness/treatment events we will explore their perception of any influence during this event, of their completed ReSPECT form. With all participants we will explore whether and how their reflections on the ReSPECT process and its impact have changed and what influenced these changes.

3.7.4 Interviews with GPs and conversations with practice staff

GP Interviews will be conducted as described for patients and family/carer interviews.

We will design our GP interviews to deepen our understanding of how GPs operationalise the ReSPECT process within their complex and time pressured clinical workflows using the framework of NPT (10,80). We will ask the GPs to familiarise themselves with the ReSPECT forms of any patient-participants, or other ReSPECT forms they have completed to use as an aide memoire in the interview. In the first part of the interview we will:

- Explore their understanding of the ReSPECT process and form (*coherence*)
- Ask them to give an account of up to 3 ReSPECT processes they were involved in including the ReSPECT process for a patient participant if they were involved in completing the relevant form. We will inform them of this request before the interview so that they can refresh their memory of the case. We will prompt for why they decided to start each process, how they set up the process, their experience of any conversation with patient and family, their experience of form completion, storage and accessibility for future reference, consideration of reviewing the completed ReSPECT form and how the ReSPECT form influences the care they provide for the patient (*cognitive participation*). We will ask the GP not to disclose any personal identifiable information relating to patients during the interview.
- Ask them to describe how the practice organises the ReSPECT process (*collective action*).
- Explore thoughts and feelings the process engendered; whether the process changed how they viewed their patients and themselves; whether they think it resulted in a change to the care and clinical interventions they provide and how they viewed these; any impact on the trust between patients and themselves or the health service more generally including whether they trust that the ReSPECT form will receive attention within general practice and other parts of the health service once completed; how the ReSPECT process could be improved. We will ask whether they have been involved in any other advance care planning discussions and whether COVID-19 has had any impact on their thinking about ReSPECT or advance care planning in general (*reflexive monitoring*).

In the second part of the interview we will ask the GPs to reflect on their experience with ReSPECT using the four propositions developed by May et al (11). We will ask the clinician to what extent it reflects their current experience and what is missing.

During fieldwork, in conversation with practice staff we will explore their understanding of the ReSPECT process and form, their role (if any) in the ReSPECT process, how the practices organises and documents the ReSPECT process, and whether they perceive any impacts on patients or staff.

3.7.5 Nursing and care home managers

Interviews will be conducted as described for GPs, patients and family/carer interviews. The questions will map to the NPT framework and aim to understand their perceptions and experience of the linked general practice's ReSPECT process. We will explore:

- Their understanding of the ReSPECT process and form (*coherence*).
- Their role in the ReSPECT process including whether they initiate the process by suggesting it to a resident/family member or GP, (*cognitive participation*)
- Collective action: how the practices organises the ReSPECT process in relation to their nursing/care home and how the home stores the forms and uses them when the resident is ill (*collective action*).
- Whether their involvement with ReSPECT has changed their thoughts or behaviours and whether they perceive any impacts on their residents or staff of the ReSPECT process. We will ask whether they have been involved in any other advance care planning discussions and whether COVID-19 has had any impact on their thinking about ReSPECT or advance care planning in general (*reflexive monitoring*).

We will draw on existing evidence related to ACP in care homes to further guide out interview questions and prompts (86,87,88).

3.8 Data Management

Data collected during the study will be handled and stored in accordance with GDPR, the 2018 Data Protection Act and WMS Standard Operating Procedures. Prior to participant consent to participation all personal data will be held by the relevant GP practice or care home. The practice will complete a data collection spreadsheet in which each patient identified with a ReSPECT form completed in the previous 12 months will be entered and given a unique identifier code. This database will be used to send invitation letters to patients and their family, and to provide aggregate anonymous data to the Warwick co-ordinating research team. Participants who contact the research team and agree to take part in the study will, with their agreement, have their name and contact details stored securely on the University server for the purposes of contacting them again to arrange an interview.

Consent for interview will be taken and interview data will be collected by researchers from the co-ordinating study team. Consent forms will be stored at Warwick University in the Division of Health Sciences in a locked filing cabinet in a locked room with access limited to the core study team.

Interviews will be recorded, transcribed verbatim and anonymised with each participant being assigned a unique interview ID. Transcripts will be stored in a separate electronic folder to the database of names and contact details of participants. Recordings of interviews will be collected on encrypted devices then securely transferred on the same day to the secure university server. After transfer the recording will be deleted from the recording device. Transfer to any transcription services will be done via a secure system and according to Warwick data transfer SOPs and a data sharing agreement. Any handwritten field notes will be kept in a locked filing cabinet in a locked room in the University. If on site field work is permitted, field notes recorded electronically will be on an encrypted, password

protected laptop while the researcher is at the site and then uploaded to secure university servers the same day.

Transcripts will be uploaded to NVivo software for data management together with any field notes. All transcripts and notes of conversations will be coded; 30% independently by a second researcher. Data analysis will be concurrent with data collection and initial analysis will inform subsequent data collection.

3.9 Analysis

Data analysis will be carried out by the two study senior research fellows. Analysis codes and emerging themes will be discussed at regular analysis meetings with the core research team (FG, AS, CH, RS). Following data collection at the four practices we will hold a data analysis meeting with members of our lay advisory group and our PPI co-investigator. Their input into the analysis will contribute to refining the interview guides for subsequent interviews and ensure that the patient perspective is not lost in our interpretation.

Using framework analysis (89) we will seek to understand how and why the implementation of ReSPECT varies, including identifying dissonance between accounts of different actors: patient, family/carer, GP, nursing/care home manager. We will test and extend existing theory and evidence on the enablers of, and obstacles to, implementing advance care planning in primary care and understand how it applies to the ReSPECT process (69). We will tease out implications for trust between patients and their healthcare providers and within wider society, refining or adding to existing theory (71). We will explore the data for explicit and implicit ethical concerns, using the approach of grounded moral analysis to explicate the ethical dimensions of the ReSPECT process (90). We will identify dilemmas faced within the ReSPECT process and analyse their ethical dimensions. We will describe how participants negotiate the complex issues related to the ReSPECT process in the time-limited context of primary care (9). In particular, to inform future educational materials, we will bring to light how participants seek to simplify aspects of the ReSPECT process through the use of heuristics or simple tools/rules and critically evaluate them.

4. WP TWO: UNDERSTANDING THE WIDER CONTEXT OF RESPECT IMPLEMENTATION (OBJECTIVE 2)

In this work package we will engage the public and health professionals in reflection and debate about the concept and use of Emergency Care Treatment Plans in general and ReSPECT in particular. As the use of ECTPs relates to ethical issues and death, we will engage with representatives of all relevant faith communities. Based on findings and those of WP1, we will develop two questionnaire surveys. One for the public, and one for primary and community health care professionals.

There are two research strands in this WP:

- Focus groups/interviews
- Surveys.

4.1 Research question

What does the wider public and primary and community health and social care professionals think about the concept and use of emergency care treatment plans?

4.2 Focus groups and Interviews

4.2.1 Methods

We will run two focus groups in each of the three primary care areas (six in total), one with members of the public with an interest in healthcare, and one with non-GP health and social care professionals. We will also interview local faith leaders in these areas.

4.2.2 Recruitment

By contacting community groups with an interest in health. we aim to recruit people who are keen to engage in discussion about the ReSPECT process including its complexity, sensitivity, and ethical considerations, and willing to challenge the current approaches where necessary and suggest change. For the focus groups we will speak to each participant in advance of the focus group and send text message or email reminders to reduce no-shows.

Members of the public with an interest in health care:

We will work with the regional Health Watch for the relevant CCG area to identify community groups, including local faith groups, and patient support organisations with which to advertise the study. We will provide a poster about the study, an invitation email and brief information leaflet describing the study for circulation through these groups. We will ask those interested in participating in a focus group or learning more about the study to contact the research team. We will also advertise through GP practice patient groups. From those expressing an interest we will purposively sample to reflect the sociodemographic and ethnic profile of the community covered by the CCG until we have at least 10 people willing to participate.

Health and social care professionals:

We will advertise through local branches of professional organisations relevant to primary and community health and social care in each CCG area and through social media, including organisations for nurses working in the community, paramedics, care workers/managers, community physio- and occupational therapists, and community palliative care professionals. We will ask any health and social care professional working in the community who is interested in participating to contact the research team. From those expressing an interest we will select to ensure diversity of professional background, until we have at least 10 professionals willing and able to attend each focus group.

If we are unable to recruit sufficient numbers of participants for focus groups we will also offer an invitation to individual interviews to the same sample populations.

Faith leaders:

We will interview up to ten faith leaders across the three areas. We will work with the relevant Health Watch in each area to identify the prevailing faith groups through which we will invite a range of faith leaders in the community to participate in an interview. We will provide an information leaflet and invitation letter/email text for this purpose. For those people who contact the research team to express interest the researcher will discuss the study, answer questions and if the person agrees to participate will arrange a time for the interview.

Faith leaders and participants in focus groups/individual interviews will receive a shopping voucher (Love2shop) in payment for their time at INVOLVE rates; currently £20 per hour. Participants will also have travel expenses reimbursed.

4.2.3 Consent

The researcher will contact potential participants, answer any questions about the study and check that they are able to attend the focus group. We will send each participant a more detailed information sheet with a contact telephone number to use if they have any further questions. Two weeks prior to the focus group we will send participants a briefing pack about the ReSPECT programme and a summary of findings from initial analysis of WP1 data. Consent will be obtained by the researcher prior to the focus group, either in person or remote verbal consent obtained if the focus group is to be held online.

4.2.4 Withdrawals

Focus group participants will be able to withdraw their data from the study up to three weeks following data collection (that is prior to analysis), by contacting the study team.

4.2.5 Data collection

Focus groups: An experienced facilitator will conduct each focus group, with a researcher present to take notes. Prior to starting discussion, the facilitator will briefly present what was in the briefing pack as a reminder and to frame the discussion. Focus group members will be prompted to discuss the following questions:

- What are the benefits for patients, carers, the health service, and wider society of the implementation of ReSPECT?
- What are the dis-benefits of ReSPECT?
- What gets in the way of implementing ReSPECT in a way that optimises its use and value, including access to forms and transfer between health and social care organisations?
- How can the dis-benefits of ReSPECT, and challenges to its use be mitigated?

Interviews with faith leaders: These will follow a similar format to the focus groups. We will ask them to consider the discussion topics used for the focus groups. In addition we will:

- Explore how the values embedded in ReSPECT sit with key values of their faith e.g. treating people with dignity, respecting their autonomy, appropriate use of modern medical technologies and prevention of unnecessary harm and suffering.
- Explore areas where the faith's teaching supports the premise of ReSPECT or could cause challenges for people following that faith and any thoughts on how to mitigate these challenges for both patients and clinicians.

4.2.6 Data Management

Focus group and interview data will be managed in the same manner as described for interview data in WP1.

4.2.7 Analysis

We will use Framework analysis (89) to develop themes where there is consistency of opinion in relation to the use of ReSPECT and importantly, themes about which there is some tension or diversity of opinion. This tension or diversity may be within a focus group discussion, within what individuals talk about in relation to their family/community/faith group, or identified through comparing data from different focus groups/interviews. From this we will identify the key issues of consistency and importantly for the survey, issues where there is of tension/diversity. It is these latter issues that will form the focus of the surveys.

4.3 National surveys of public and GPs

4.3.1 Method

Informed by our early qualitative work we will work with our PPI advisory group to identify key questions to measure public awareness and acceptability of ECTPs. Questions are likely to cover use of Emergency Care Treatment Plans or ReSPECT, views on advance decision making (timing, relevant medical conditions, content of decisions/recommendations), and how likely participants are to complete an Emergency Care Treatment Plan.

Again, informed by our qualitative work we will work with our GP co-investigators and PPI advisory group to develop a questionnaire survey to measure the views of GPs nationally regarding the use of Emergency Care Treatment Plans including ReSPECT, in primary care. We will measure their knowledge of Emergency Care Treatment Plans and ReSPECT, views on their use in primary care, and how likely they are to complete an Emergency Care Treatment Plan for their patients.

4.3.2 Sample and data collection

We will outsource the surveys to specific purpose designed survey providers as a cost and time efficient approach to obtain representative data.

Public survey: We will commission the National Centre for Social Research to include our questions in the annual British Social Attitudes Survey (BSA). The BSA is the UK's longest-running survey of public opinion. It is viewed as the authoritative barometer of public attitudes by Government, academia and the media. It provides a high quality nationally representative attitudinal survey with a stratified sample based on postcode and including face-to-face data collection using computer assisted personal interviewing and pen and paper self-completion. New questions are tested in two pilots before use in the survey. We will commission our questionnaire items to be presented to 1000 participants.

GP survey: We will distribute the survey through a specific GP survey platform, medeConnect GP Omnibus, a monthly online survey of UK GPs. The survey provides data on commissioned questions presented to a regionally representative sample of 1000 GPs including partners, salaried non-principals and locums. We will refine our questions, developed from our qualitative data, to include in the GP Omnibus Survey using think aloud interviews with a small number of GPs and pilot the finalised questions with a sample of GPs prior to distribution to ensure the questions generate meaningful data.

4.3.3 Data Management

Anonymised clean data sets from both surveys will be transferred securely from the survey company to Warwick University where they will be stored as password protected files on the university server

accessible only to the core study team. A data sharing agreement will be in place between the survey organisation and the University. No personal data will be transferred. Any concerns participants had about their anonymised information being included in the survey will have been addressed by the organisation collecting the data at the time. Withdrawal will not be possible as only anonymised survey data will be provided to the study team. The data will be analysed by the study team statistician.

4.3.4 Sample size and analysis

We will analyse associations between the primary and secondary survey outcomes using regression models fitted to these outcomes with covariates such as: age, gender, presence/type of chronic disease, functional status (patients), time in practice and practice size (GPs). Our primary outcome is the binary outcome ‘whether they would complete an emergency care treatment plan’ (for themselves if public survey and for their patients if GP survey) for which we will fit a logistic regression model.

For our primary analysis we will dichotomise responses to our 11 point numerical rating score. Scores of <5 we consider as being ‘unwilling’ and a score is ≥ 5 , as being ‘willing’. The minimum sample size available for either survey is 1,000 responses. There is little additional benefit from a greater precision for a larger survey. For each of the surveys, in case of the binary outcome, if 50% (or 60%, 70%, 80% and 90%) of participants consider completing the form, this would ensure 6.2% (or 6.1%, 5.7%, 5% and 3.7% respectively) precision. This is a very good precision for such a study.

In a secondary analysis we will investigate ‘how willing patients are to complete an emergency care plan on a scale of 0 till 10’. We consider this outcome to be continuous and will fit a regression model to investigate associations between the covariates of interests and this outcome. Similar linear models will be fitted to other secondary outcomes. These additional secondary outcomes will be determined as an outcome of our qualitative work and included in the survey questions.

A sample size of 1,000 will give us ample statistical accuracy for these analyses.’

5. WP THREE. INVESTIGATING CONGRUENCE BETWEEN RESPECT RECOMMENDATIONS AND TREATMENT DECISIONS: AN ANALYSIS OF PATIENT RECORDS (OBJECTIVE 4)

5.1 Research question

Our research questions for this WP are:

- What is the quality of completion of ReSPECT forms in primary care?
- To what extent do treatment decisions in an acute or emergency situation reflect the recommendations on a person’s ReSPECT form?

5.2 Method

Retrospective analysis of patient records and ReSPECT forms.

Inclusion criterion

- patients with a ReSPECT form completion recorded in the previous 12 months
- Aged 18 years or over

Exclusion criteria

- Age less than 18 years
- No documented ReSPECT form

5.3 Recruitment

In each of our 12 participating practices we will use the list obtained in WP1 of people with a ReSPECT form recorded. A member of practice staff will identify all adults (18 and over) in the practice with a ReSPECT form completed in the previous twelve months and use the same checking processes as in WP1 prior to mailing invitation letters to all identified patients, or if the patient lacks capacity, the person whose contact details are on the completed ReSPECT form. A purposive sample of patients with a ReSPECT form completed in the last 6 months will also be invited to take part in the interview study (WP1). Participants recruited to the interview study can choose to opt out of the notes evaluation.

The practice will send a letter to the patient, or for patients who lack capacity the family member recorded as their carer in relation to the ReSPECT form or their designated next of kin/carers in the practice record, informing them about the study. Information will include, the purpose of the study, the information that will be collected, and how it will be anonymised before secure transfer to the study team. The letter will include details of the ways in which the person can let the study team know that they do not want their medical records to be used in the study. Each information sheet and response form will have a unique study identification code which will be used by the study team to inform the practice if a person informs them that they do not wish to be included in the medical record review. No personal identifiable information will be transmitted to the central study team.

As in WP1, we will also work with care homes linked to the practice (practice is the primary practice for the home). We will ask the care home manager to give a brief information leaflet and expression of interest form to any resident not involved in WP1 who is registered with the practice, or their next of kin if the resident lacks capacity in relation to ReSPECT, who has a ReSPECT form. We will ensure a cross checking process with the practice to avoid duplication of invitations. Next of kin can contact the study team directly or return an opt out form in a FREEPOST envelope. Care home staff will facilitate the return of an opt out form from residents with capacity.

5.4 Withdrawals

A record of who the participant identification numbers have been allocated to will be kept at the research site in order to enable withdrawal of participants who opt out. No data will be collected from medical records for four weeks after the information letters have been posted to enable time for patients to opt out. In the event a patient requests their data not to be included after it has been collected but before the data is analysed, we will treat this as if it were a withdrawal and their data will be removed from the data set.

5.5 Data collection

A copy of the ReSPECT form and print outs of the patient record since ReSPECT form completion including hospital discharge letters and ED correspondence up to six months post ReSPECT completion will be generated by a member of practice staff or study researcher with a research passport. Each record copy will have personal identifiers removed and be given the participant's unique identifier code. The copy record will be matched with the patient's ReSPECT form (similarly de identified and coded) and then transferred securely to the study team. This element of data collection is subject to our application to the Confidentiality Advisory Group.

We will ask the care home to make a copy of the form, remove all personal identifiers, and transfer this electronically through a secure system to the study team. The participant unique ID code (allocated when the participant was added to the practice held database of patients with ReSPECT forms) will be added to the ReSPECT form so that it can be linked with the patient record.

5.6 Data Management

Relevant sections of patient records will be printed by a member of practice staff or study researcher with a research passport and all personal identifiers removed and a study code assigned. Record print outs will be linked to the relevant ReSPECT form. Each anonymised record will then be transferred securely to the University and stored either in a locked filing cabinet in a locked office (hard copies) or on a password protected secure database on the University server (scanned electronic copies).

5.7 Sample size

The primary outcome in this analysis is the binary outcome 'congruency or not between what is stated on the ReSPECT form and what is actually implemented'. We are only aware of one study that has assessed congruency in a similar manner. Hickman in a study in nursing facilities in the USA found a congruency of 74% between POLST forms and later treatment decisions with regard to choices about comfort care (47) (Multiple other analyses were also performed) This analysis is the closest to our primary analysis of congruence between free text comment on planned care and actual care.

Assuming a percentage congruency of 75%, using a binomial model for binary data, to obtain a 95% confidence interval with 10% precision, we would need 289 completed ReSPECT forms with at least one event in the follow up period. Assuming that out of the recruited participants with completed ReSPECT forms, only 70% will have at least one event in the follow up period, we would need to recruit 413 participants in total with a completed ReSPECT form.

5.8 Analysis

All included ReSPECT forms will be reviewed using the evaluation tool that we have developed in our recently completed study of ReSPECT in secondary care to assess quality of completion of each section of the form.

For all included ReSPECT forms the relevant data (treatment recommendations including CPR decision (Part 4 of the ReSPECT form) will be extracted from the form. The corresponding medical record will be reviewed to identify the first hospital admission or significant treatment event since ReSPECT form completion. A significant treatment event will be defined as an acute episode requiring a medical

treatment decision. Exclusion criteria will be: minor trauma, acute urinary or respiratory infections not classified as pneumonia. The relevant section of the record will be extracted to compare with the ReSPECT recommendations.

The record will be compared with the recommendations in section four of the corresponding ReSPECT form to look for congruence comparing the decisions made during the event (including decisions not to admit to hospital) with:

- Free text recommendations (specific guidance on clinical interventions that may or may not be wanted or clinically appropriate, including being taken or admitted to hospital +/- receiving life support)
- Any CPR decision if cardiac arrest occurred

Two researchers, working independently, will first identify the first significant treatment critical event since form completion, and the first after consent received. This will be done blind to the content of the ReSPECT form. Where there is a disagreement that cannot be resolved a third team member will arbitrate. They will then independently decide if each of these decisions was congruent with the ReSPECT form. With disagreement being resolved by discussion or arbitration by a third reviewer. After the first 20 record reviews the researchers, WP leads and CI will meet to discuss and develop some decision rules for assessing recorded treatment decisions. These will then be applied to future record reviews with further checking and revision if necessary.

We anticipate that it is the first significant treatment event after completion of the ReSPECT form where it is most likely to inform decision making and that this is the most interesting event to study. The practicalities of prospectively identifying, and following up people with newly completed ReSPECT forms to identify all first events would not be possible within the lifetime of this study. Our participants may not be representative of all those who had had a ReSPECT form completed in the previous six months. We will not be able to access the ReSPECT forms of those who have died, left the practice, or are terminally ill. For our primary analysis, we will review the first recorded significant treatment event after completion of the form. We will however also collect data on the first event after recruitment, where a pre-recruitment event was identified, and do a sensitivity analysis using the first significant treatment event after recruitment.

Our primary analysis will be on congruence of free text recommendations and clinical care. We will fit a logistic regression model to the binary primary outcome 'congruency or not between what is stated on the ReSPECT form and what is actually implemented' with covariates such as: age, gender, place of residence, health category and cognitive function. Similar linear regression models will be fitted to our secondary outcome of congruency with DNACPR choices. The analysis of congruence between DNACPR decision and actual CPR decisions will be a secondary analysis.

We will also present simple descriptive statistics.

6. WP4 EVIDENCE SYNTHESIS AND STAKEHOLDER CONFERENCE (OBJECTIVES 3 AND 5)

Findings from our previous work packages will be synthesised in a process of triangulation to look for convergence and divergence within the overarching framework of NPT (how different stakeholders conceptualise, engage with and use the ReSPECT process). We will identify key findings and their associated implications for practice relevant to a) patients and families; b) health care professionals; and c) managers and policy makers. We will present our key findings at a stakeholder conference to be held at Warwick University. Participants invited will include members of patient and carer organisations, community and faith groups, relevant professional organisations, policy makers, and regulatory bodies, and members off the national ReSPECT working group. We will use the principles of experience-based co-design to inform the structure of the meeting (91). The meeting will commence with a formal presentation of the study findings, followed by an invited presentation from a senior member of the Resuscitation Council's ReSPECT sub committee which will outline the current ReSPECT implementation support strategy, including their suggested core audit criteria. Following the presentations, participants, working in facilitated mixed small groups, will listen to an film/audio recording created to illustrate touch points within the ReSPECT process. A touch point is talk that is emotionally significant. Touch points will be identified from across all our interview and focus group data. They will be identified during initial analysis in collaboration with our PPI panel. The audio clips of these touch points will be edited together for the stakeholder conference. At the conference, after listening to the audio the small groups will identify how ReSPECT implementation could be improved. After sharing these initial ideas in plenary, participants will work further in new facilitated mixed small groups to refine the key messages that have emerged from our findings to consider how identified obstacles to implementation of ReSPECT in Primary Care can be overcome. We will specifically consider any identified challenges to involving minority or marginalised groups in the ReSPECT process and suggested mechanisms for successful embedding of ReSPECT in day-to-day primary care practice. Following the meeting the project team will draw on its conclusions to draft a suggested framework to support improved effective implementation of ReSPECT (and emergency care treatment plans more broadly). We will follow the process used successfully in a previous NIHR funded project on decision-making around admission to intensive care(92). Anticipated outputs will be:

- recommendations for training for primary care staff,
- a decision support framework for making emergency care and treatment decisions using ReSPECT, suggested audit criteria for quality assessment of the ReSPECT process,
- information/support materials for patients and families.

A draft report of the conference and suggested recommendations will be sent to all participants for further comment before finalising.

7. STUDY ORGANISATION AND OVERSIGHT

7.1 Sponsor and governance arrangements

The University of Warwick will act as the Sponsor for this study. Warwick standard operating procedures will be followed.

7.2 Ethical approval

All required ethical approval(s) for the trial will be sought using the Integrated Research Application System. The trial will be conducted in accordance with all relevant regulations.

Before enrolling patients into the study, the research team must ensure that the local conduct of the trial has the agreement of the relevant participating organisation as well as overarching HRA approval in place.

Substantial amendments to the protocol will be communicated to all relevant parties (i.e. investigators, sponsor, NIHR, REC, participating sites, local CRN).

Annual reports will be submitted to the REC within 30 days of the anniversary date on which the favourable opinion was given, and annually until the study is declared ended. The REC will be notified of the end of the trial (whether at planned time or prematurely). The CI will submit a final report to the required authorities with the results, including any publications within one year of the end of the study.

7.3 Trial Registration

The study will be eligible for inclusion on the CRN Portfolio.

7.4 Indemnity

NHS indemnity covers NHS staff for any actions performed as part of the study. The University of Warwick provides indemnity for any harm caused to participants by the design of the research protocol and conduct of the research by its staff

7.5 Project timetable and milestones

Tasks		Set up	Year one				Year two			
			Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
Liaison with HS&DR	contracting									
Management group meetings		xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx	xxx
Steering group meetings			x		x		x		x	
Lay advisory group meetings			x	x	x		x		x	
Ethics R&D and HRA approvals										
WP1&3 site recruitment										
WP1&3 site set up										
WP1	Development of study manual & processes									
Interviews	Data collection									
	Analysis									
WP2										
FGs/faith leader interviews	Recruitment									
	Data collection									
	Analysis									
Surveys	Development of questions									
	Data collection									
	Analysis									
WP3	Development of study manual & processes									
	Data collection									
	Analysis									
WP4	Synthesis of findings from WPs1-3									
	Stakeholder meeting									
	Development of suggestions for practice									
Report writing/dissemination										

Figure 2 Plan of investigation and timetable

Key milestones:

- 6 months: 4 sites open for data collection WPs1 & 3
- 12 months: data collection WP1 complete at 8 sites; Surveys initiated
- 18 months: Analysis Wps1-3 complete, WPs 1&3 papers submitted

24 months: Project report, survey paper and overall project paper submitted

7.6 Administration

The study co-ordination will be based at Warwick Medical School, University of Warwick.

7.7 Study Management Group (SMG)

The Study Management Group, consisting of the project staff and co-investigators involved in the day-to-day running of the trial, will meet regularly throughout the project. Significant issues arising from management meetings will be referred to the Study Steering Committee or Investigators, as appropriate.

7.8 Study Steering Committee (SSC)

The study will be guided by a group of respected and experienced personnel and researchers, as well as at least one 'lay' representative. The SSC will have an independent Chairperson. Meetings will be held at six monthly intervals throughout the project. Routine business is conducted by email, post or teleconferencing.

The Steering Committee, in the development of this protocol and throughout the study will take responsibility for:

- Major decisions such as a need to change the protocol for any reason
- Monitoring and supervising the progress of the study
- Reviewing relevant information from other sources
- Informing and advising on all aspects of the study

The full remit and responsibilities of the SSC will be documented in the Committee Charter which will be signed by all members.

7.9 Data Monitoring Committee (DMC)

Since there is no intervention delivered as part of the study, a DMC is not required.

7.10 Essential Documentation

A Study Master File will be set up according to WMS Standard Operating Procedures and held securely at the coordinating centre.

The coordinating centre will provide Investigator Site Files to all sites involved in the study.

7.11 Financial Support

The study has been funded by a grant from the National Institute for Health Research (NIHR) Health Services and Delivery Research programme.

8. MONITORING AND QA

All research staff involved in data collection for WP1, WP2 and WP3 will have had GCP training or University of Warwick research integrity training. Training will also be carried out for research staff who may answer phone calls from patients or legal representatives and need to deal sensitively with their questions.

All interviews will be conducted by the co-ordinating team Research Fellow(s). Consent procedures, interview schedules and a process for recording field notes will be developed and reviewed by researchers, PPI advisory group, and co-investigators responsible for each work package, ensuring a consistent, but flexible approach needed for this type of data collection.

Quality assurance during analysis of qualitative data: coding will be undertaken by independent researchers for 30% of transcripts and any inconsistency discussed to ensure consistency.

Data quality checks will have been done by the BSA and medeConnect GP Omnibus, according to their protocols, prior to transfer of the anonymised data set to University of Warwick for analysis.

Each site will receive an initiation visit where study training will be delivered by the co-ordinating centre researchers. Training will be recorded on a log and stored in the study master file.

After the initial site visits to each practice and care home, the study manager will have regular contact with the sites to identify any problems with compliance with the protocol, training, data collection, or other barriers to progress, and to support sites with the day-to-day management of the study. As well as regular telephone and email contact, and the co-ordinating centre researcher visiting for data collection, a site visit (conducted virtually by video conference) may be arranged if there are particular issues that are best resolved face to face. The study manager will check with each site that all Investigator Site File documents are up to date at least once during the study.

9. PPI INVOLVEMENT

The research was initially developed with the PPI group from our current ReSPECT evaluation study and this group have agreed to continue in this role for this study. Our PPI co-investigator will contribute to the day to day running and organisation of the study, in addition to reviewing patient information resources and contributing to final report writing and dissemination. He also has extensive experience of public engagement and advocacy in his role as Chief Executive of Health Watch Warwickshire. Further PPI input will be provided through two independent members of the Study Steering Committee. Our separate advisory group will be involved at all stages of the study, advising on patient and public facing documentation, contributing to analysis and interpretation of findings and active engagement in our stakeholder conference.

10. DISSEMINATION AND PUBLICATION

The results of the study will be reported first to study collaborators. The main report will be drafted by the study team, and the final version will be agreed by the Study Steering Committee before submission for publication, on behalf of the collaboration.

The success of the study depends on the collaboration of GPs, practice staff, and care home staff, from across the UK. Equal credit will be given to those who have wholeheartedly collaborated in the study.

The study will be reported in accordance with the relevant reporting guidelines (<http://www.equator-network.org>).

The output of this work will have maximal impact through the adoption of a dissemination strategy with three strands. The first will ensure that patients and public are informed of the study results; the second will engage practitioners and health care planners to implement the findings, and the third will involve consulting with policy makers for maximum impact.

Patients and the public: Patient and public understanding of the reasons for and process of emergency care treatment planning is an essential part of establishing the trust necessary to enable implementation of policy and practice. We will produce a plain English summary of the study findings that we will disseminate through patient and community organisations we have identified through the study in addition to national support groups such as Age-UK, Alzheimer's Society, and patient panels of professional organisations. We will work with our PPI group and also seek suggestions for contacts during our focus groups and interviews with faith leaders. We will seek the support of regional Health Watch organisations to disseminate in their areas. Our PPI Co-investigator, Bain, is chief executive of Warwickshire Health Watch and will provide links to other health watch organisations. We will work with actors to develop short videos of patient stories based on our data to be used as an educational and public information resource to be distributed through community groups and general practices. We will also post information about the study and its findings on NHS and University websites and social media.

Practitioners: Through our contacts with the RCUK ReSPECT subcommittee we will disseminate our findings to local and regional ReSPECT implementation. We will work with the RCUK ReSPECT clinical lead and subcommittee to ensure emerging study findings are fed into training and support initiatives as early as possible. We will work with local and regional implementation teams in our study sites to develop brief information/training videos for practice staff and care home staff which can be disseminated through CCGs. We will disseminate our findings to primary care practitioners through presentations at regional and national meetings, web-based resources, and social media. We will submit our key findings to open access, high impact journals with a wide general readership (e.g. BMJ, BJGP, Health Service Journal).

Policy makers: We will continue engagement with key policy makers (NHS England, Department of Health, Clinical Commissioning Groups, Care Quality Commission) during this work with the aim of ensuring the project delivers information of value to any future changes to policy. The project will summarise the key successes and limitations of integrating ReSPECT in primary care. It will assist policy makers by providing an evidence base to inform the need for any changes or refinement to policy. Policy makers and managers will be invited to our stakeholder meeting.

The strategies for dissemination could have the following impacts.

For patients and the public, knowledge about the effects and impacts of emergency care and treatment plans could be used to enable them on a personal level to become more involved in decision-making about these aspects of their care. If emergency care and treatment plans do reduce inappropriate attempts at resuscitation and admission to hospital, it should increase the number of patients who experience a peaceful death in the place of their preference. If patient and relative involvement in

decision-making is improved. The knowledge about patient experience generated by the study could be used by individuals and patient and public organisations to inform to public discussion about anticipatory decision-making for emergency care and treatment.

For clinicians, we will specifically seek out exemplars as best practice to show case how emergency care treatment plans are best used to support ethical decision-making in partnership with patients and relatives. This will help to increase confidence in their use. Knowledge of enablers and potential obstacles to integration of emergency care treatment plans in primary care practice can prompt practitioners, managers and commissioners to consider structural and process changes that would enable them to be used more effectively. We anticipate our findings could be used to generate learning materials that can contribute to RCGP and Marie Curie training on the Daffodil standards for advanced serious illness and end of life care. We have experience of working with e Learning for Health and the Faculty of Intensive Care Medicine to develop e Learning modules based on previous research and will explore similar opportunities with the RCGP.

We anticipate that our findings will inform future development of support and educational resources developed by the RCUK ReSPECT team, and through their work with local and regional implementation leads will directly impact how ReSPECT is operationalised in primary care practice.

The project will contribute evidence to inform policy makers in considering whether and how to develop policy or recommendations on the use ECTPs at a national level.

11. REFERENCES

1. Findlay G , Shotton H, Kelly K, Mason M. Time to intervene? A review of patients who underwent cardiopulmonary resuscitation as a result of an in-hospital cardiorespiratory arrest. A report by the National Confidential Enquiry into Patient Outcome and Death 2012. http://www.ncepod.org.uk/2012report1/downloads/CAP_fullreport.pdf.
2. Perkins GD, Griffiths F, Slowther AM, et al. Do-not-attempt-cardiopulmonary-resuscitation decisions: an evidence synthesis. Southampton (UK): NIHR Journals Library; April 2016..
3. Beach MC, Morrison RS. The effect of do-not-resuscitate orders on physician decision-making. J Am Geriatr Soc. 2002;50:2057-61
4. Henneman EA, Baird B, Bellamy PE, Faber LL, Oye RK. Effect of do-not-resuscitate orders on the nursing care of critically ill patients. Am J Crit Care. 1994;3:467-72.
5. Chen JL, Sosnov J, Lessard D, Goldberg RJ. Impact of do-not-resuscitation orders on quality of care performance measures in patients hospitalized with acute heart failure. Am Heart J. 2008;156:78-84.
6. Hayhoe B. Howe A. Advance care planning under the Mental Capacity Act 2005 in primary care. Br J Gen Pract. 2011;61(589):e537-41
7. National Institute for Health and Care Excellence. NICE guideline 94 Chapter 15 Advance Care Planning. NICE 2018. <https://www.nice.org.uk/guidance/ng94/evidence/15.advance-care-planning-pdf-172397464602>

8. Gold Standards Framework; Advance Care Planning
<https://www.goldstandardsframework.org.uk/advance-care-planning>
9. Lund S, Richardson A, May C. Barriers to advance care planning at the end of life: an explanatory systematic review of implementation studies. *PLoS One*. 2015;10(2):e0116629.
10. Fritz Z, Slowther AM, Perkins GD. Resuscitation policy should focus on the patient, not the decision. *BMJ* 2017 Feb 28;356:j813 (8)
11. May C, Myall M, Lund S, et al. Managing patient preferences and clinical responses in acute pathophysiological deterioration: What do clinicians think treatment escalation plans do?. *Soc Sci Med*. 2020;258:113143. doi:10.1016/j.
12. POLST website <https://polst.org/programs-in-your-state/>
13. <https://www.northdevonhealth.nhs.uk/wp-content/uploads/2015/07/V10-FINAL-Guidance-for-TEP-and-Flowchart.pdf>
14. <https://www.northerncanceralliance.nhs.uk/deciding-right/deciding-right-regional-forms/>
15. Perkins GD, Griffiths F, Slowther AM, et al. Do-not-attempt-cardiopulmonary-resuscitation decisions: an evidence synthesis. Southampton (UK): NIHR Journals Library; April 2016.
16. Perkins G Evaluation of the Recommended Summary Plan for Emergency Care and Treatment. NIHR funded project 15/15/09 <https://www.fundingawards.nihr.ac.uk/award/15/15/09>
17. Eli K, Ochieng C, Hawkes C, et al. Secondary care consultant clinicians' experiences of conducting emergency care and treatment planning conversations in England: an interview-based analysis. *BMJ Open*. 2020;10(1):e031633. Published 2020 Jan 20. doi:10.1136/bmjopen-2019-031633
18. Hall A; Rowland C; Grande G. How Should End-of-Life Advance Care Planning Discussions Be Implemented According to Patients and Informal Carers? A Qualitative Review of Reviews. *Journal of Pain & Symptom Management*. 2019;58(2):311-335
19. https://www.theguardian.com/society/2020/jul/05/is-there-a-better-way-to-die?CMP=Share_AndroidApp_Outlook
20. <https://compassionindying.org.uk/help-the-nhs-with-a-living-will-sunday-times-letter/>
21. Care home residents claim they are being asked to sign letters agreeing they won't go to hospital if they get coronavirus. *Daily Mail* 2020 Apr 16. <https://www.dailymail.co.uk/news/article-8226675/Care-home-residents-asked-sign-letters-agreeing-WONT-hospital-COVID-19.htm>
22. General Medical Council, Nursing and Midwifery Council. Statement on advance care planning during the Covid-19 pandemic, including do not attempt cardiopulmonary resuscitation (DNACPR). April 2020 . <https://www.nmc.org.uk/globalassets/sitedocuments/other-publications/statement-advance-care-planning-dnacpr-gmc-nmc.pdf>
23. British Medical Association, Care Provider Alliance, Care Quality Commission, Royal College of General Practice. Joint statement on advance care planning. March 2020. <https://content.govdelivery.com/accounts/UKCQC/bulletins/283e565>

24. Care Quality Commission. Review of Do not Attempt Cardiopulmonary Resuscitation decisions during the COVID-19 pandemic: Interim report. November 2020
<https://www.cqc.org.uk/sites/default/files/20201204%20DNACPR%20Interim%20Report%20-%20FINAL.pdf>
25. RCGP. Ethical guidance on COVID-19 and primary care
<https://elearning.rcgp.org.uk/mod/page/view.php?id=10557>
26. BMA. COVID-19 toolkit for GPs and GP practices. July 2020
<https://elearning.rcgp.org.uk/mod/page/view.php?id=10557>
27. Fadel MG, Parekh K, Hayden P, Krishnan P. Improving resuscitation decisions: a trust-wide initiative. *BMJ Open Qual.* 2018;7(4):e000268.
28. Shermon E, Munglani L, Oram S, William L, Abel J. Reducing DNACPR complaints to zero: designing and implementing a treatment escalation plan using quality improvement methodology. *BMJ Open Qual.* 2017;6(2):e000011.
29. Stockdale C, Trivedi B, Jerome E, et al. Implementation of a combined Cardiopulmonary Resuscitation and Treatment Escalation Plan document in a District General Hospital. *BMJ Qual Improv Rep.* 2013;2(2):u202653.w1236
30. Fielden L, Lynch D, Petrie K. A quantitative and qualitative evaluation of the ReSPECT Process in Forth Valley. Scotland's first ReSPECT pilot. NHS Forth Valley. 2019
31. <https://arc-w.nihr.ac.uk/research/projects/evaluating-the-recommended-summary-plan-for-emergency-care-and-treatment-respect/>
32. Hickman, S. E., Keevern, E., & Hammes, B. J. (2015). Use of the physician orders for life-sustaining treatment program in the clinical setting: a systematic review of the literature. *Journal of the American Geriatrics Society*, 63(2), 341–350. <http://doi.org/10.1111/jgs.13248>
33. An HJ, Jeon HJ, Chun SH, et al. Feasibility Study of Physician Orders for Life-Sustaining Treatment for Patients with Terminal Cancer. *Cancer Res Treat.* 2019;51(4):1632-1638. doi:10.4143/crt.2019.009
34. Mack DS, Dosa D. Improving Advanced Care Planning through Physician Orders for Life-Sustaining Treatment (POLST) Expansion across the United States: Lessons Learned from State-Based Developments. *Am J Hosp Palliat Care.* 2020;37(1):19-26. doi:10.1177/1049909119851511
35. Torke AM, Hickman SE, Hammes B, et al. POLST Facilitation in Complex Care Management: A Feasibility Study. *Am J Hosp Palliat Care.* 2019;36(1):5-12. doi:10.1177/1049909118797077
36. Dillon E, Chuang J, Gupta A, et al. Provider Perspectives on Advance Care Planning Documentation in the Electronic Health Record: The Experience of Primary Care Providers and Specialists Using Advance Health-Care Directives and Physician Orders for Life-Sustaining Treatment. *Am J Hosp Palliat Care.* 2017;34(10):918-924. doi:10.1177/1049909117693578
37. Lake RE, Franks L, Meisenberg B. Reducing Discrepancy Between Code Status Orders and Physician Orders for Life-Sustaining Therapies: Results of a Quality Improvement Initiative. *Am J Hosp Palliat Care.* 2020;37(7):532-536. doi:10.1177/1049909119899079

38. Mauleon J, Staffileno BA. Integration of an Advance Care Planning Model in Home Health: Favorable Outcomes in End-of-Life Discussions, POLST Rates, and 60-day Hospital Readmissions. *Home Healthc Now*. 2019;37(6):337-344.
39. Hanson LC, Zimmerman S, Song MK, et al. Effect of the Goals of Care Intervention for Advanced Dementia: A Randomized Clinical Trial. *JAMA Intern Med*. 2017;177(1):24-31. doi:10.1001/jamainternmed.2016.7031
40. McGough NN, Hauschildt B, Mollon D, Fields W. Nurses' knowledge and comfort levels using the Physician Orders for Life-sustaining Treatment (POLST) form in the progressive care unit. *Geriatric Nursing (New York, N.Y.)*. 2015 Jan-Feb;36(1):21-24. DOI: 10.1016/j.gerinurse.2014.09.001.
41. Ballou JH, Dewey EN, Zonies DH. Elderly patients presenting to a Level I trauma center with Physician Orders for a Life-Sustaining Treatment form: A propensity-matched analysis. *J Trauma Acute Care Surg*. 2019;87(1):153-160. doi:10.1097/TA.0000000000002321
42. Lee RY, Brumback LC, Sathitratanacheewin S, et al. Association of Physician Orders for Life-Sustaining Treatment With ICU Admission Among Patients Hospitalized Near the End of Life [published online ahead of print, 2020 Feb 16]. *JAMA*. 2020;323(10):950-960. doi:10.1001/jama.2019.22523
43. Kelly C, Vranas, Amber L. Lin, Dana Zive, Susan W. Tolle, Scott D. Halpern, Christopher G. Slatore, Craig Newgard, Robert Y. Lee, Erin K. Kross, Donald R. Sullivan, The Association of Physician Orders for Life-Sustaining Treatment With Intensity of Treatment Among Patients Presenting to the Emergency Department, *Annals of Emergency Medicine*, 10.1016/j.annemergmed.2019.05.008, (2019).
44. Pedraza S, Culp S, Knestrick M, Falkenstine E, and Moss A (2017). "Association of Physician Orders for Life-Sustaining Treatment Form Use With End-of-Life Care Quality Metrics in Patients With Cancer." *Journal of Oncology Practice*. 13(10):e881-e888. doi:10.1200/JOP.2017.022566
45. RJ Schmidt, BB Weaner, D Long (2015). "The Power of Advance Care Planning in Promoting Hospice and Out-of-Hospital Death in a Dialysis Unit" *Journal of Palliative Medicine*. 18(1)62-66. doi:10.1089/jpm.2014.0031
46. Tuck KK, DM Zive, TA Schmidt, J Carter, J Nutt, and EK Fromme (2015). "Life-sustaining treatment orders, location of death and co-morbid conditions in decedents with Parkinson's disease" *Parkinsonism & Related Disorders* 21(10):1205-1209. doi:10.1016/j.parkreldis.2015.08.021
47. Hickman SE, Nelson CA, Moss AH, Tolle SW, Perrin NA, Hammes BJ. The consistency between treatments provided to nursing facility residents and orders on the Physician Orders for Life-Sustaining Treatment form. *J Am Geriatr Soc*. 2011;59(11):2091-2099.
48. Hammes BJ, Rooney BL, Gundrum JD, Hickman SE, Hager N. The POLST program: a retrospective review of the demographics of use and outcomes in one community where advance directives are prevalent. *J Palliat Med*. 2012;15(1):77-85
49. Alvin H. Moss, Dana M. Zive, Evan C. Falkenstine, Erik K. Fromme, Susan W. Tolle, Physician Orders for Life-Sustaining Treatment Medical Intervention Orders and In-Hospital Death Rates: Comparable Patterns in Two State Registries, *Journal of the American Geriatrics Society*, 10.1111/jgs.14273, 64, 8, (1739-1741), (2016).

50. Jiminez G, Tan WS, Virk AK, Low CK, Car J, Ho AHY. Overview of systematic reviews of advance care planning: Summary of evidence and Global lessons. *Journal of pain and symptom management* 2018;56(3):436-59
51. Combes S, Nicholson CJ, Gillett K, Norton C. Implementing advance care planning with community-dwelling frail elders requires a system-wide approach: An integrative review applying a behaviour change model. *Palliat Med.* 2019;33(7):743-756. doi:10.1177/0269216319845804
52. Gilissen J, Pivodic L, Smets T, et al. Preconditions for successful advance care planning in nursing homes: A systematic review. *Int J Nurs Stud.* 2017;66:47-59. doi:10.1016/j.ijnurstu.2016.12.003
53. Ke LS, Huang X, Hu WY, O'Connor M, Lee S. Experiences and perspectives of older people regarding advance care planning: A meta-synthesis of qualitative studies. *Palliat Med.* 2017;31(5):394-405. doi:10.1177/0269216316663507
54. Batchelor F, Hwang K, Haralambous B, et al. Facilitators and barriers to advance care planning implementation in Australian aged care settings: A systematic review and thematic analysis. *Australas J Ageing.* 2019;38(3):173-181. doi:10.1111/ajag.12639
55. Solis GR, Mancera BM, Shen MJ. Strategies used to facilitate the discussion of advance care planning with older adults in primary care settings: A literature review. *J Am Assoc Nurse Pract.* 2018;30(5):270-279.
56. Blackwood DH, Walker D, Mythen MG, Taylor RM, Vindrola-Padros C. Barriers to advance care planning with patients as perceived by nurses and other healthcare professionals: A systematic review. *J Clin Nurs.* 2019;28(23-24):4276-4297. doi:10.1111/jocn.15049
57. Johnson CE, McVey P, Rhee JJ, et al General practice palliative care: patient and carer expectations, advance care plans and place of death—a systematic review *BMJ Supportive & Palliative Care* Published Online First: 25 July 018. doi: 10.1136/bmjspcare-2018-00
58. Zwakman M, Jabbarian LJ, van Delden J, et al. Advance care planning: A systematic review about experiences of patients with a life-threatening or life-limiting illness. *Palliat Med.* 2018;32(8):1305-1321. doi:10.1177/0269216318784474
59. Owen L, Steel A. Advance care planning: what do patients want?. *Br J Hosp Med (Lond).* 2019;80(5):263-267. doi:10.12968/hmed.2019.80.5.263
60. Hemsley B, Meredith J, Bryant L, et al. An integrative review of stakeholder views on Advance Care Directives (ACD): Barriers and facilitators to initiation, documentation, storage, and implementation. *Patient Educ Couns.* 2019;102(6):1067-1079. doi:10.1016/j.pec.2019.01.007
61. Fahner JC, Beunders AJM, van der Heide A, et al. Interventions Guiding Advance Care Planning Conversations: A Systematic Review. *J Am Med Dir Assoc.* 2019;20(3):227-248. doi:10.1016/j.jamda.2018.09.014
62. Gleeson A, Noble S, Mann M. Advance care planning for home health staff: a systematic review [published online ahead of print, 2019 Feb 26]. *BMJ Support Palliat Care.* 2019; doi:10.1136/bmjspcare-2018-001680

63. Arruda LM, Abreu KPB, Santana LBC, Sales MVC. Variables that influence the medical decision regarding Advance Directives and their impact on end-of-life care. *Einstein (Sao Paulo)*. 2019;18:eRW4852. Published 2019 Oct 10. doi:10.31744/einstein_journal/2020RW4852
64. Khandelwal N, Kross EK, Engelberg RA, Coe NB, Long AC, Curtis JR. Estimating the effect of palliative care interventions and advance care planning on ICU utilization: a systematic review. *Crit Care Med*. 2015;43(5):1102-1111. doi:10.1097/CCM.0000000000000852
65. Hong M, Yi EH, Johnson KJ, Adamek ME. Facilitators and Barriers for Advance Care Planning Among Ethnic and Racial Minorities in the U.S.: A Systematic Review of the Current Literature. *J Immigr Minor Health*. 2018;20(5):1277-1287. doi:10.1007/s10903-017-0670-9
66. McDermott E, Selman LE. Cultural Factors Influencing Advance Care Planning in Progressive, Incurable Disease: A Systematic Review With Narrative Synthesis. *J Pain Symptom Manage*. 2018;56(4):613-636. doi:10.1016/j.jpainsymman.2018.07.00 De
67. Rahemi Z, Williams CL. Does ethnicity matter—Cultural factors underlying older adults’ end-of-life care preferences: A systematic review. *Geriatric Nursing Volume 41, Issue 2, March–April 2020, Pages 89-97*
68. Johnson SB, Butow PN, Kerridge I, Bell ML, Tattersall MHN. How Well Do Current Measures Assess the Impact of Advance Care Planning on Concordance Between Patient Preferences for End-of-Life Care and the Care Received: A Methodological Review. *J Pain Symptom Manage*. 2018;55(2):480-495. doi:10.1016/j.jpainsymman.2017.09.008
69. De Vleminck A, Houttekier D, Pardon K, et al. Barriers and facilitators for general practitioners to engage in advance care planning: a systematic review. *Scand J Prim Health Care*. 2013;31(4):215-226. doi:10.3109/02813432.2013.854590
70. Risk J, Mohammadi L, Rhee J, et al Barriers, enablers and initiatives for uptake of advance care planning in general practice: a systematic review and critical interpretive synthesis *BMJ Open* 2019;9:e030275. doi: 10.1136/bmjopen-2019-03027
71. Flo, E., Husebo, B.S., Bruusgaard, P. et al. A review of the implementation and research strategies of advance care planning in nursing homes. *BMC Geriatr* 16, 24 (2016). <https://doi.org/10.1186/s12877-016-0179->
72. Calnan M, Rowe R. Trust in Health Care: An agenda for future research. London The Nuffield Trust 2004
73. Kingston A, Robinson L, Booth H, Knapp M, Jagger C. Projections of multimorbidity in the older population in England to 2035: estimates from the Population Ageing and are Simulation (PACSim) Model. *Age and Ageing*. 2018;47:374-80
74. National Audit Office. Reducing Emergency Admissions: Report by the Comptroller and Auditor General. Department of Health and Social Care. NHS England. 2018. <https://www.nao.org.uk/wp-content/uploads/2018/02/Reducing-emergency-admissions-Summary.pdf>
75. Arne Wolters, Filipe Santos, Therese Lloyd, Creina Lilburne and Adam Steventon. Emergency admissions to hospital from care homes: how often and what for? The Health Foundation. 2019

<https://www.health.org.uk/sites/default/files/upload/publications/2019/Emergency-admissions-from-care-homes-IAU-Q02.pdf>

76. Marie Curie. Emergency Hospital Admissions briefing. 2018
<https://www.mariecurie.org.uk/blog/emergency-admissions-report/183441>
77. Arnold E, Finucane AM, Oxenham D Preferred place of death for patients referred to a specialist palliative care service *BMJ Supportive & Palliative Care* 2015;5:294-29.
78. Fried TR, Van Doorn C, O’Leary JR, Tinetti ME, Drickamer MA. Older persons’ preferences for site of terminal care. *Annals of Internal Medicine*. 1999; 131(2): 109–12.
79. Bassford C, Griffiths F, Svantesson M, Ryan M, Krucien N et al. Developing an intervention around referral and admissions to intensive care: a mixed-methods study. Southampton. NIHR Journals Library. 2019
80. General Medical Council. Consent: Patients and doctors making decisions together. GMC. 2012
81. NHS England. Universal Personalised Care: Implementing the comprehensive model. 2019.
<https://www.england.nhs.uk/wp-content/uploads/2019/01/universal-personalised-care.pdf>
82. May C. Towards a general theory of implementation. *Implement Sci*. 2013 Feb 13;8:18.
83. Gallacher K May C, Montori VM, MAir FS. Understanding patients’ experiences of treatment burden in chronic heart failure using normalisation process theory. *Annals of Family Medicine*. 2011;9:235-243
84. Malterud K, Siersma VD, Guassora AD. Sample size in qualitative interview studies: guided by information power. *Qualitative Health Research*. 2016;26(13):1753-60
85. Bentley B, O’Connor M. Conducting research interviews with bereaved family carers: when do we ask? *J Palliat Med*. 2015 Mar;18(3):241-5. doi: 10.1089/jpm.2014.0320. Epub 2014 Dec 17. PMID: 25517136; PMCID: PMC4348368.
86. Ramsbottom K, Kelley M. Developign strategies to improve advance care planning in long term care homes. Giving voice to residents and their family members. *International Journal of palliative care*. 2014; | <https://doi.org/10.1155/2014/358457>
87. Stewart F, Goddard C, Schiff R, Hall S. Advanced care planning in care homes for older people: a qualitative study of the views of care staff and families. *Age Ageing*. 2011;40(3):330-335. doi:10.1093/ageing/afr006
88. Stone L, Kinley J, Hockley J. Advance care planning in care homes: the experience of staff, residents, and family members. *Int J Palliat Nurs*. 2013;19(11):550-557. doi:10.12968/ijpn.2013.19.11.550
89. Gale, N.K., Heath, G., Cameron, E. et al. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 13, 117 (2013) doi:10.1186/1471-2288-13-117
90. Davies, R., Ives, J. & Dunn, M. A systematic review of empirical bioethics methodologies. *BMC Med Ethics* 16, 15 (2015)

91. Robert G, Cornwell J, Locock L, Purushotham A, Sturmey G, Gager M. Patients and staff as codesigners of healthcare services. *BMJ* : 2015;350:g7714
92. Bassford C, Griffiths F, Svantesson M, Ryan M, Krucien N, Dale J, Rees S, Rees K, Ignatowicz A, Parsons H, Flowers N, Fritz Z, Perkins G, Quinton S, Symons S, White C, Huang H, Turner J, Brooke M, McCreedy A, Blake C, Slowther A. Developing an intervention around referral and admissions to intensive care: a mixed-methods study. Southampton (UK): NIHR Journals Library; 2019 Nov. PMID: 31765112.