

The effects of computerised decision support systems on nursing and allied health professional performance and patient outcomes: a systematic review and user contextualisation

Carl Thompson^{1*}, Teumzghi Mebrahtu¹, Sarah Skyrme¹,
Karen Bloor³, Deidre Andre², Anne Maree Keenan¹,
Alison Ledward¹, Huiqin Yang¹ and Rebecca Randell⁴

¹School of Healthcare, University of Leeds, Leeds, UK

²Library Services, University of Leeds, Leeds, UK

³Department of Health Sciences, University of York, York, UK

⁴Faculty of Health Studies, University of Bradford, Bradford, UK

*Corresponding author

Disclosure of interests

Full disclosure of interests: All authors have completed the unified competing interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare (1) no financial support for the submitted work from anyone other than their employer; (2) no financial relationships with commercial entities that might have an interest in the submitted work; (3) no spouses, partners, or children with relationships with commercial entities that might have an interest in the submitted work; and (4) no non-financial interests that may be relevant to the submitted work.

Primary conflicts of interest: Carl Thompson, Rebecca Randell and Alison Ledward are, or have been, members of the NIHR Health Services Delivery and Research funding committee.

Published June 2023

DOI: 10.3310/GRNM5147

This report should be referenced as follows:

Thompson C, Mebrahtu T, Skyrme S, Bloor K, Andre D, Keenan AM, *et al.* Effects of computerised decision support systems on nursing and allied health professional performance and patient outcomes: a systematic review and user contextualisation *Health Soc Care Deliv Res* 2023. <https://doi.org/10.3310/GRNM5147>

Abstract

Effects of computerised decision support systems on nursing, midwife and allied health professionals' performance and patient outcomes: a systematic review and user contextualisation

Carl Thompson^{1*}, Teumzghi Mebrahtu¹, Sarah Skyrme¹,
Karen Bloor³, Deidre Andre², Anne Maree Keenan¹,
Alison Ledward¹, Huiqin Yang¹ and Rebecca Randell⁴

¹School of Healthcare, University of Leeds, Leeds, UK

²Library Services, University of Leeds, Leeds, UK

³Department of Health Sciences, University of York, York, UK

⁴Faculty of Health Studies, University of Bradford, Bradford, UK

*Corresponding author c.a.thompson@leeds.ac.uk

Background: Computerised decision support systems (CDSS) are widely used by nurses and allied health professionals but their effect on clinical performance and patient outcomes is uncertain.

Objectives: Evaluate the effects of clinical decision support systems use on nurses', midwives' and allied health professionals' performance and patient outcomes and sense-check the results with developers and users.

Eligibility criteria: Comparative studies (randomised controlled trials (RCTs), non-randomised trials, controlled before-and-after (CBA) studies, interrupted time series (ITS) and repeated measures studies comparing) of CDSS versus usual care from nurses, midwives or other allied health professionals.

Information sources: Nineteen bibliographic databases searched October 2019 and February 2021.

Risk of bias: Assessed using structured risk of bias guidelines; almost all included studies were at high risk of bias.

Synthesis of results: Heterogeneity between interventions and outcomes necessitated narrative synthesis and grouping by: similarity in focus or CDSS-type, targeted health professionals, patient group, outcomes reported and study design.

Included studies: Of 36,106 initial records, 262 studies were assessed for eligibility, with 35 included: 28 RCTs (80%), 3 CBA studies (8.6%), 3 ITS (8.6%) and 1 non-randomised trial, a total of 1318 health professionals and 67,595 patient participants. Few studies were multi-site and most focused on decision-making by nurses (71%) or paramedics (5.7%). Standalone, computer-based CDSS featured in 88.7% of the studies; only 8.6% of the studies involved 'smart' mobile or handheld technology. Care processes – including adherence to guidance – were positively influenced in 47% of the measures adopted. For example, nurses' adherence to hand disinfection guidance, insulin dosing, on-time blood sampling, and documenting care were improved if they used CDSS. Patient care outcomes were statistically – if not always clinically – significantly improved in 40.7% of indicators. For example, lower numbers of falls and

ABSTRACT

pressure ulcers, better glycaemic control, screening of malnutrition and obesity, and accurate triaging were features of professionals using CDSS compared to those who were not.

Evidence limitations: Allied health professionals (AHPs) were underrepresented compared to nurses; systems, studies and outcomes were heterogeneous, preventing statistical aggregation; very wide confidence intervals around effects meant clinical significance was questionable; decision and implementation theory that would have helped interpret effects – including null effects – was largely absent; economic data were scant and diverse, preventing estimation of overall cost-effectiveness.

Interpretation: CDSS can positively influence selected aspects of nurses', midwives' and AHPs' performance and care outcomes. Comparative research is generally of low quality and outcomes wide ranging and heterogeneous. After more than a decade of synthesised research into CDSS in healthcare professions other than medicine, the effect on processes and outcomes remains uncertain. Higher-quality, theoretically informed, evaluative research that addresses the economics of CDSS development and implementation is still required.

Future work: Developing nursing CDSS and primary research evaluation.

Funding: This project was funded by the National Institute for Health and Care Research (NIHR) Health and Social Care Delivery Research programme and will be published in *Health and Social Care Delivery Research*; 2023. See the NIHR Journals Library website for further project information.

Registration: PROSPERO ¹ [number: CRD42019147773].

List of abbreviations

AHP	allied health professional/profession as defined and outlined at www.england.nhs.uk/ahp/role/	INR	international normalised ratio (a measure of time for blood to clot)
CBA	controlled before-and-after studies	ITSS	interrupted time series study
CDSS	computerised/computerized decision support system	NASSS	non-adoption, abandonment, scale-up, spread, sustainability (framework)
CNIO/CAHPIO	Chief Nursing Information Officer; Chief Allied Health Profession Information Officer	NMAHP	nurses, midwives and allied health professions
CPOE	computerised physician order entry	NPT	normalisation/normalization process theory
EHR	electronic health record	NRCTs	non-randomised controlled trials
ER/D	emergency room/department	PPI	Patient Public Involvement
HRQOL	health-related quality of life	RCTs	randomised controlled trials

Plain language summary

Computerised decision support systems (CDSS) are software or computer-based technologies providing advice to professionals making clinical decisions – for example, which patients to treat first in emergency departments. CDSS improve some doctors' decisions and patients' outcomes, but we don't know if they improve nurses', midwives' and therapists' or other staff decisions and patient outcomes. Research into, and health professionals' use of, technology – for example, in video consultations – has grown since the last relevant systematic review in 2009.

We systematically searched electronic databases for research measuring how well nurses, midwives and other therapists/staff followed CDSS advice, how CDSS influence their decisions, how safe CDSS are, and their financial costs and benefits. We interviewed CDSS users and developers and some patient representatives from a general practice to help understand our findings.

Of 35 relevant studies – from 36,106 initially found – most (71%) focused on nurses. Just over half (57%) involved hospital-based staff, and three-quarters (75%) were from richer countries like the USA or the UK. Research quality had not noticeably improved since 2009 and all studies were at risk of potentially misleading readers. CDSS improved care in just under half (47%) of professional behaviours, such as following hand-disinfection guidance, working out insulin doses, and sampling blood on time. Patient care – judged using outcomes like falls, pressure ulcers, diabetes control and triage accuracy – was better in 41% of the care measured. There wasn't enough evidence to judge CDSS safety or the financial costs and benefits of systems.

CDSS can improve *some* nursing and therapist decisions and *some* patient outcomes. Studies mostly measure different behaviours and outcomes, making comparing them hard. Theories explaining or predicting how decision support systems might work are not used enough when designing, implementing or evaluating CDSS. More research into the financial costs and benefits of CDSS and higher-quality evidence of their effects are still needed. Whether decision support for nurses, midwives and other therapists reliably improves decision-making remains uncertain.

Introduction

Until recently, healthcare professionals relied on peer-to-peer, paper-based, or standalone guidelines, and limited computer technology to support their clinical judgements and decisions.^{2,3} Since the late 1980s, claims that computerised support at the point of care has potential to improve treatment or management have increased – notably in medicine.⁴ The degree to which the *potential* of computerised support for decisions is *actually* realised is unclear.

In this synopsis we bring together the findings of an evidence synthesis of comparative research into the effects of CDSS on the clinical performance, behaviours and outcomes associated with the work and decisions of nurses, midwives and allied health professionals (NMAHPs, for example, physiotherapists, occupational therapists and paramedics – see www.england.nhs.uk/ahp/role/ for definitions and scope). Our aim is an accessible overview of the synthesis and associated stakeholder engagement. To improve accessibility, we have abridged some of the reporting of our results and methods.

Background

The target for CDSS: decision-making by nurses, midwives and allied health professionals

Historically, decisions about the delivery and organisation of healthcare were assumed to be the province of doctors. Whilst medical dominance has proven remarkably resistant to challenge, ‘decision-rich’ areas such as the prescribing of medications,⁵ the initiation of critical care outreach in acute care, nutritional management and rehabilitation planning offer the chance for professions other than medicine to formally use their decisions to shape the delivery of healthcare, how care processes are experienced and the clinical outcomes that result.

Alongside formal decisions in healthcare such as assigning a diagnosis, prescribing a treatment, or offering a prognosis, the *realpolitik* of healthcare delivery relies on a range of informal judgements, decisions, and negotiated positions between a range of professionals – often with fluid and overlapping roles.⁶ Technology has encroached into healthcare decision-making, purporting to offer support, information and recommendations to help shape professional decisions.⁷

In this synopsis we focus on nurses, midwives and allied health professionals (NMAHPs). Why? Because their work, demographic composition, educational levels, and socio-economic positions often differ from medicine and doctors, but they contribute to a complex, fluid, and – crucially – negotiated division of labour in healthcare.⁸ Authority within this division of labour stems in part from the power to exercise clinical judgement, clinically reason and make or shape decisions. If we assume that work in healthcare is based on – and reflects – reasoned judgements and decisions, then it follows that different professionals in multidisciplinary teams will face different uncertainties, judgements and decisions. The support needed for tackling differing uncertainties may also be different.

Research into the decisions and decision-making of NMAHPs is relatively scarce compared to studies focusing on doctors and medical reasoning, although eminent decision scientists have studied nursing decisions since the 1960s.⁹ Researchers have also described and typologised ‘nursing’ decisions.³ Some scholars point out that there is no de facto reason why nurses – and by implication, other health professionals – should be treated as possessing their own, unique, decision-making cognition, even if decisions when viewed in context appear different.¹⁰ Others have extended well-established descriptive and prescriptive theories of generic professional decision-making to incorporate forms of knowledge and knowing (such as ‘reflection-in-action’) associated with particular groups – notably nurses.¹¹

At the heart of attempts to describe, model and theorise clinical decision-making are two core constructs:

- judgements – the weighing-up or evaluation of clinical, research or other information
- decisions – choosing between discrete options.

The most parsimonious models of decision-making bring together judgement, choice and evaluation in a 'feedback loop'.¹² Consequently, decision-making rarely feels like a discrete event made up separate 'stages' to the decision-maker. Despite the difficulties of 'holistic' decision-making as experienced by decision-makers, there is value in separating it into component elements; in part because the characteristics of decisions (1) determine the style of clinical reasoning best suited to a decision (intuition vs. rational information-processing), but also (2) shape the likelihood of using different forms of decision support.^{3,11} The perceived time available to make choices, the perceived structure of a choice, and the need to show how you got to a judgement or decision (i.e. a choice's visibility) can increase or decrease the chances of using technology-delivered support.^{3,11}

Support for decision-making often comes in the form of technologies – ranging from paper-based aids such as printed guidelines or research summaries, to web, app and computer-based decision support systems. More commonly, support can also come from informal resources such as a colleague's advice, or, at the extreme, a professional's own internalised resources in the form of experience, knowledge recalled from training, or just gut instinct or 'intuition'. It is the application of computer technology to judgements and decisions that is our focus in this synopsis.

What are computerised decision support systems?

Computerised decision support systems (CDSS) are software- or computer-based technologies that offer patient-specific recommendations based on either research, expert opinion, machine learning/artificial intelligence or combinations of these, and designed to influence the clinical decision-making of health professionals.^{13–15} CDSS access patient information from practitioners, healthcare staff, patients' manual data entry or queries of electronic medical records before research or expert knowledge is assessed to provide computer-generated recommendations delivered to the clinician via a computer/tablet, mobile-phone screen or electronic medical record. Clinicians can then choose whether to use these recommendations. Examples of decision support used by NMAHPs include: assessing fall risk and preventative behaviours;¹⁶ pressure-ulcer management;¹⁷ selecting interventions for managing musculoskeletal disorders;¹⁸ screening for childhood language disorders;¹⁹ depression screening²⁰ and, on a whole-system scale, choices faced in clinical pathways for primary care triage and prioritisation (<https://digital.nhs.uk/services/nhs-pathways>).

CDSS come in two main forms: (1) knowledge-based and (2) non-knowledge-based.²¹ Knowledge-based CDSS use logical 'IF-THEN-ELSE' rules to evaluate information provided directly by a clinician or drawn from an electronic health record. These are then matched to a computerised knowledge base (in many cases expert opinion or national/international clinical practice guidelines) to provide assessments/management options/probabilities or actionable recommendations or outputs.^{21,22} These forms of CDSS automate information-gathering and provide advice in line with guidelines. Examples of this type of CDSS are drug prescription/alert tools and emergency and out-of-hours telephone calls used for triaging patients. Non-knowledge-based CDSS use machine learning and artificial intelligence rather than flowchart-style rules or logic to support clinicians' decision-making.²¹ Typical examples of this type of CDSS are predictive risk models for assessing the prognosis of a disease outcome.²³ CDSS based around artificial intelligence and/or machine learning are less common than rule-based systems in NMAHP work.

CDSS systems can stand alone, integrated into, or at least capable of interacting with, wider digital infrastructure in health systems such as electronic health records (EHRs) or computerised physician order entry (CPOE – computer-based systems that automate instructions, with standardised, legible, and complete orders). They can be hosted via a computer, tablet or smartphone, and have web-based/local or 'app' interfaces. CDSS can present information on host devices or via the integrated EHR/CPOE system.

Why look at CDSS for NMAHPs?

NMAHPs make decisions that could benefit from digital support. Unwarranted variations in practice and outcome, for patients with seemingly similar issues and facing similar decisions and uncertainties, exist. These uncertainties make a synthesis of empirical research timely and useful.²⁴⁻²⁶ New ways of working and support for these new roles for NMAHPs feature in many health systems. Opportunity costs associated with digital technology for learners and educators exist: professional preparation for and continuing professional development of digitally competent clinicians able to use new technologies effectively require time.²⁷

We had three main research-based motivations for the synthesis. First, clinical decision support systems will only be useful if they improve clinicians' decision performance (for example, more accurate diagnoses and prognoses), improve patient health outcomes (e.g. morbidity, mortality, fewer adverse events), and offer perceived value for money for health services.^{28,29} We do not know if any of these are true for NMAHPs.

Second, a previous review of studies on CDSS use by nurses found only limited impact on performance and health outcomes.³⁰ The review is more than 13 years old and digital technology and the research evidence base has developed significantly. The effect of CDSS on allied health professionals (AHPs) has not been reviewed systematically. Systematic reviews of studies on the impact of CDSS on healthcare delivery generally suggest they can improve practitioner performance in specific areas of decision-making such as diagnosis (4/10 systems), disease management (23/37 systems) and drug-dosing or prescribing systems (19/29 systems).⁴ The impact on patient outcomes is more equivocal, with only 13% of systems (7/32) reporting improvements.⁴ Reviews focusing on specific areas of clinical practice such as prescribing and drug dosing³¹ or clinical subdomains such as neonatal care³² offer very limited conclusions, because the underpinning evidence is either absent,³² low quality and/or narrow in scope.³¹

Third, existing reviews often neglect the fact that whilst multi-disciplinary team members may all be involved in delivering healthcare, their decisions reflect their role in the division of labour and so are likely to differ. Extant reviews often contain an implicit rationale that doctors' decisions alone are the main mechanism for improving healthcare processes and outcomes.⁴

How are clinical decision support systems supposed to improve decision-making?

Clinical decision support systems work by providing high-quality relevant useful information delivered when it is required to decision-makers.¹³ The main generative mechanism by which CDSS aid NMAHPs decision-making is the combination of CDSS-generated information/suggestions with existing nurse or AHP knowledge. Thus, CDSS augment or supplement clinician decision-making rather than replacing it. CDSS are a key means of encouraging concordance with guideline-based care to reduce unwarranted variations in practice.³³

Examples of decisions supported by CDSS include:³⁴

- Recognising patient deterioration – CDSS can increase situational awareness or incorporation of relevant clinical and research-based information in reasoning, and tailoring of local or national guidance.
- Determining patients with conditions that merit the application of clinical guidelines – CDSS improves the consistency of judgements and adherence to guideline recommendations and reduces (unwarranted) variation.
- Triaging patients, often in the emergency department or primary care, to determine priority cases – the CDSS improves the reliability of judgements and simplifies choices by reducing the 'noise' in the situation and amplifying the appropriate 'signals' to encourage more appropriate decisions.

As with any health technology, CDSS will only improve care and health if actually used by nurses and AHPs in their decision-making. Whatever the quality of the underlying knowledge base, decision rules,

analyses or algorithms, if unimplemented, or implemented badly, will not improve decision quality and patient benefit is less likely. Unfortunately, CDSS implementation and use by NMAHPs is rarely straightforward, and can be suboptimal.³⁵⁻³⁷

CDSS can create the potential for harms as well as benefits.³⁸ These include fragmentation or disruption of work and workflow; alert fatigue; deskilling and the consequences on decisions of poor-quality or incorrect knowledge in the data used for inference or analysis. Additionally, CDSS may rely on a user's computer literacy – something that is highly variable in nurses and AHPs. Systems can incur opportunity costs for clinicians as well as those charged with maintaining and supporting technology in health systems.¹⁵ CDSS can also widen existing inequalities in access to high-quality care; for example, where effective CDSS are located only in prestigious teaching hospitals and associated with improved access to services, then patients who do not have access to teaching hospitals will be disadvantaged.^{39,40}

Thus, there are three main mechanisms by which CDSS 'work' in the context of decision-making by NMAHPs: successfully combining high-quality or novel CDSS information and clinician knowledge; improving quality of care processes and – by implication – outcomes, by improving the appropriateness of recommendations, management/treatment choices, accuracy of predictions or diagnoses; and successful implementation and use by clinicians.

Theoretical framework

We used theory in three ways. First, we drew on existing reviews and meta syntheses of characteristics of CDSS associated with improved outcomes and performance^{41,42} to test the hypothesis that possessing these characteristics would positively influence CDSS aimed at NMHAPs. Second, we used Normalisation Process Theory (NPT) as a lens through which we viewed the results of the included evidence – and their background, design, discussion and/or process evaluation/descriptions – to explore and explain the ways in which CDSS manage (or not) to become embedded and routine as a part of normal, taken-for-granted, practice.⁴³ NPT provided our focus for the implementation of CDSS: the ways that CDSS are used in their social context as a form of collective action by practitioners.⁴³ Third, NPT informed our approach to coding the qualitative responses of intended CDSS users/recipients in our stakeholder engagement/sense-check exercise (see "Calibration' interviews' and 'PPI' sections).

We considered other theoretical approaches. The NASSS framework⁴⁴ (non-adoption, abandonment, scale-up, spread, sustainability) had similar a priori abilities to highlight the ways in which technologies are taken up or abandoned, but fewer people have used it in a decision support context. Actor Network Theory⁴⁵ recognises that interactions between humans and technology can shift over time and are often 'negotiated'. Applications of the theory, beyond using it as a general explanatory framework, would have entailed knowledge of the actor-networks, technologies and contexts that were often missing from study reports and beyond the scope and resources of our planned calibration exercise. NPT offered a practical, pragmatic, validated means of examining 'what people do' and 'how they work' to adopt and sustain CDSS in NMAHP work. In using this framework, this part of the study will add to the ≈130+ evaluations of varied interventions that have made explicit use of the theory.⁴⁶ It constitutes a middle-range theory of socio-technical change⁴⁷ and a theoretical framework for understanding CDSS as complex interventions.⁴³

NPT can give a perspective on CDSS, both as a technology and as a set of practices related to that technology.⁴⁸ Whilst policy and government push the case for new technologies to deliver healthcare improvement (c.f. www.gov.uk/government/news/matt-hancock-launches-tech-vision-to-build-the-most-advanced-health-and-care-system-in-the-world) the empirical literature continues to highlight an implementation gap.⁴⁸ In using NPT we sought to address aspects of adoption (alongside our 'core' systematic review) sometimes downplayed in similar CDSS reviews.^{4,41,49}

NPT centres on four core constructs:⁵⁰ ‘coherence’ – the extent to which an intervention is understood as meaningful, achievable and desirable; ‘cognitive participation’ – the enrolment of those actors necessary to deliver the intervention; collective action – the work that brings the intervention into use; and ‘reflexive monitoring’ – the ongoing process of adjusting the intervention to keep it in place. These four core constructs were used to frame our sense-check interviews with CDSS leaders, implementers and developers.

Aims and objectives

We sought to examine the impact on performance and patient outcomes associated with CDSS purporting to support the decisions and judgements of NMAHPs. To achieve this aim we had two objectives, to:

1. evaluate the clinical effectiveness and cost-effectiveness of CDSS on NMAHPs performance and patient outcomes
2. critically examine our findings in the light of interviews with people who design, implement and use CDSS systems, and to ‘calibrate’ our findings with reference to unpublished accounts.

Methods

To address our first objective, we undertook a systematic review¹ of studies comparing professionals using CDSS to those not using CDSS. Our second objective was addressed using qualitative interviews with individuals and groups seeking to encourage use of CDSS or who use or encounter them in services.

Literature searching

With an information specialist, we developed a search strategy designed to find studies focusing on CDSS and the healthcare professionals we were interested in: nurses, midwives and allied health professionals.

We ran the search strategy on multiple electronic databases and resources twice: October 2019 and February 2021. Specific databases searched included: MEDLINE (Ovid), Embase Classic+Embase (Ovid), PsycINFO (Ovid), HMIC (Ovid) Health Management Information Consortium, AMED (Allied and Complementary Medicine) (Ovid), CINAHL, Cochrane Central Register of Controlled Trials (Cochrane Database of Systematic Reviews, Wiley), Social Sciences Citation Index Expanded (Clarivate), ProQuest Dissertations & Theses Abstracts & Index, ProQuest ASSIA (Applied Social Science Index and Abstract), ClinicalTrials.gov, World Health Organisation International Clinical Trials Registry (ICTRP), Health Services Research Projects in Progress (HSRProj), OpenClinical (www.OpenClinical.org), OpenGrey (www.opengrey.eu), Health.IT.gov, Agency for Healthcare Research and Quality (www.ahrq.gov).

No date of publication and language restrictions were applied to the search. See [Appendix 1](#) for full strategy and terms.

Deciding which studies to include or exclude

Between them, six of the research team screened all the titles and abstracts retrieved. Two of the team (CT and TM) used Cochrane Collaboration Effective Practice Organisation of Care Review Group criteria⁵¹ and the study aims and objectives to decide if studies were relevant. We restricted our review to studies which compared CDSS-use to non-use, evaluated using designs less likely to lead to biased conclusions:

- randomised controlled trials (RCTs)
- non-randomised trials (NRCT)
- controlled before-and-after (CBA) studies
- interrupted time series (ITS) and repeated measures studies.




FIGURE 1 Research Pathway Diagram including PRISMA flow chart of study selection. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71.

Participants

We included studies evaluating the effects of CDSS use by NMAHPs, qualified or in training, and working in primary or secondary care. We had a long list of allied health professional categories, but in the end only paramedics, dieticians and physiotherapists were the focus of the comparative evaluations included.

Interventions

The intervention in the review was the use of any form of CDSS to aid clinical decision-making.

Comparator

The comparator was usual care: clinical practice where clinical decision-making is unsupported by CDSS. Studies must have compared care, treatment, diagnosis or management using CDSS with care, treatment or management without CDSS. We excluded CDSS aimed at diagnostic judgements where the evaluation was only against a defined reference standard. We included studies of CDSS aimed at diagnostic judgements where clinical performance with and without the CDSS featured.

Outcomes

Our primary outcome was the adherence of nurses and AHPs to evidence-based recommendations. Secondary outcomes included diagnostic accuracy, time to judgement, adverse events, health professional satisfaction, patients' health-related quality of life and costs.

Data extraction

Data on study characteristics and outcomes were independently extracted by two reviewers (TM, CT) using the Cochrane Collaboration's EPOC standard data-collection form.⁵² A third reviewer (RR) was available to resolve disagreements if needed; none occurred.

We extracted data on:

1. methods: study design, location, study setting, and date of study
2. participants: number, mean age (age range), gender, inclusion criteria, exclusion criteria of patients and providers
3. interventions: intervention components, comparison, presence of characteristics known to increase effectiveness in CDSS generally
4. outcomes: main and other outcomes specified and collected, time points reported
5. study funder.

Quality assessment

Study quality was assessed using the criteria outlined in the Cochrane Handbook for Systematic Reviews of Interventions Section 8.5⁵³ and EPOC guide.⁵⁴ TM and CT assessed studies for risk of bias. Each potential source was judged as high, low, or unclear. An overall 'Risk of bias'⁵³ assessment was set: high – a serious bias likely to decrease certainty in the results; moderate – a risk that could plausibly raise doubts about conclusions; low – risks were unlikely to alter the results.

Data synthesis

We explored heterogeneity between CDSS systems and outcomes to determine whether meta-analysis was feasible. Heterogeneity between studies in the nature of the interventions, target groups, and outcomes measures in our initial pre-searches meant a narrative approach to synthesising findings was most appropriate. Studies were grouped and summarised by clinical similarity, for example, topics studied, type of CDSS, types of health professionals involved, patient group, outcomes reported and study design.

Intervention effects were estimated using risk difference for dichotomous data and mean differences for continuous data. We calculated 95% confidence intervals where possible.⁵³ Where absolute risks were not reported, these were generated from study information. Risk difference values and 95% confidence intervals were then calculated using the absolute risk values of the comparative groups.

TABLE 1 Headline characteristics of included studies

Author and year	Country	Design	Setting	Number of sites	Study duration	HPs involved	Outcomes	Interventions
Beeckman <i>et al.</i> 2013 ⁸⁹	Belgium	RT	Nursing homes	4	5 months	Nurses and physios	Risk of pressure ulcers; HP knowledge and attitude	Pre-vPlan (a six-step clinical practice to reduce pressure ulcers using CDSS) A standard protocol (a hard copy with no implementation strategy) of reducing pressure ulcers
Bennet <i>et al.</i> 2016 ⁶¹	UK	ITS	Emergency department, district general hospital	1	1 year	Nurses	Triage prioritization; pain assessment and management; management of neutropenic sepsis	Triage CDSS [intervention period] Triage CDSS [pre-intervention period]
Blaha <i>et al.</i> 2009 ⁷⁵	Czech Republic	RT	ICU post elective cardiac surgery, university hospital	1	48 hours	Nurses	Intensive care glycaemic control/diabetes	Intervention (CDSS-model predictive control algorithm) Control-1 (paper based-Matias protocol) Control-2 (paper based-Bath protocol)
Byrne 2005 ⁸³	USA	CBA	Nursing homes	90	33 months	Nurses	Falls and pressure ulcer reduction (assessment and prevention)	CDSS use CDSS non-use
Canbolat <i>et al.</i> 2019 ⁷⁶	Turkey	Non-RT	ICU university general hospital	1	22 months	Nurses (and physicians)	ICU glycaemic control	CDSS use Usual care
Cavalcanti <i>et al.</i> 2009 ⁷⁷	Brazil	Clustered RT	ICU general hospital	5	19 months	Nurses	ICU glycaemic control	Intervention (CDSS use computer-assisted insulin protocol) Control-1 (Leuven protocol) Control-2 (conventional treatment)

TABLE 1 Headline characteristics of included studies (continued)

Author and year	Country	Design	Setting	Number of sites	Study duration	HPs involved	Outcomes	Interventions
Cleveringa <i>et al.</i> 2008 ⁷¹	Netherlands	Clustered RT	Primary care practices	26	1 year	Nurses (and physicians)	Management and prevention of diabetes (and CV risk factors)	CDSS use Usual care
Cleveringa <i>et al.</i> 2010 ⁷²	Netherlands	Clustered RT	Primary care practices	26	1 year	Nurses	Management and prevention of diabetes (and CV risk factors)	Same as Cleveringa <i>et al.</i> 2008 but a cost effectiveness study.
Cortez 2014 ⁶⁶	USA	Clustered RT	Academic medical centre oncology clinics	4	11 weeks	Nurses	Management of cancer symptoms	Intervention (drop down boxes) Control (no drop-down boxes)
Dalaba 2015 ⁸⁶	Ghana	CBA	Primary care health centres (midwifery)	12	2 years	Nurses	Maternal care	CDSS use Usual care (CDSS non-assisted)
Dowding <i>et al.</i> 2012 ⁹⁰	USA	ITS	General hospitals	29	6 years	Nurses	Risk assessment, falls and pressure ulcer prevention	CDSS use Usual care (CDSS non-assisted)
Duclos <i>et al.</i> 2015 ⁸⁴	France	Clustered RT	Paediatric wards in a university hospital	6	2 years	Dieticians	Nutritional care in malnourished children	CDSS use Usual care (CDSS non-assisted)
Dumont <i>et al.</i> 2012 ⁷⁸	USA	RT	ICU wards in a regional referral hospital	1	4 months	Nurses	Glycaemic control	CDSS use paper protocol (modified Portland protocol)
Dykes <i>et al.</i> 2010 ^{91,92}	USA	Clustered RT	Urban hospitals	4	6 months	Nurses	Fall prevention	CDSS use Usual care
Dykes <i>et al.</i> 2020 ⁹²	USA	ITS	Academic medical centres	3	42 months	Nurses	Fall prevention	Pre-intervention period Post-intervention period
Fitzmaurice <i>et al.</i> 2000 ⁶⁷	UK	RT	Primary care/general practice	12	1 year	Nurses	Oral anticoagulation care	CDSS use (nurses) CDSS non-assisted physicians

continued

TABLE 1 Headline characteristics of included studies (continued)

Author and year	Country	Design	Setting	Number of sites	Study duration	HPs involved	Outcomes	Interventions
Forberg <i>et al.</i> 2016 ⁸⁵	Sweden	Clustered RT	Paediatric university hospital	12	3 months	Nurses	Management of peripheral venous catheters in paediatrics	CDSS use Usual care (CDSS non-assisted)
Fossum <i>et al.</i> 2011 ⁹³	Norway	CBA	Nursing homes	15	2 years	Nurses	Preventative behaviours and management of nutrition	CDSS use Usual care (CDSS non-assisted)
Geurts <i>et al.</i> 2017 ⁷³	Netherlands	RT	University paediatric hospital	1	2 years	Nurses	Management of (re) hydration in children	Nurse-led CDSS Usual care
Hovorka <i>et al.</i> 2007 ⁷⁹	Czech Republic	RT	Cardiac surgery, university hospital	1	48 hours	Nurses	Glycaemic control	CDSS use Usual care (CDSS non-assisted)
Kroth <i>et al.</i> 2006 ⁶⁸	USA	RT	University hospital	1	9 months	Nurses	Body temperature assessment	CDSS use Usual care (CDSS non-assisted)
Lattimer <i>et al.</i> 1998 ⁶⁴	UK	RT	Primary care practices	1	1 year	Nurses & physicians	Emergency call assessment	Nurses with CDSS Control (doctors with no CDSS)
Lattimer <i>et al.</i> 2000 ⁶⁵	UK	RT	Primary care practices	1	1 year	Nurses & physicians	Cost analysis of emergency call assessments	Nurses with CDSS Control (doctors with no CDSS)
Lee <i>et al.</i> 2009 ⁷⁴	USA	RT	University trainee -school of nursing	1	8 months	Nurses	Obesity management	CDSS use Usual care (CDSS non-assisted)
Lv <i>et al.</i> 2019 ⁹⁴	China	RT	Community healthcare centres	4	1 year	Nurses	Chronic asthma management	CDSS use Usual care
Mann <i>et al.</i> 2011 ⁸⁰	USA	RT	Surgical military hospital ICU	1	6 days	Nurses	Glycaemic control in burn intensive care patients	CDSS use Usual care (paper-based protocol)
McDonald <i>et al.</i> 2017 ⁸⁸	USA	RT	Nursing care homes	1	2 months	Nurses	Management of chronic medical condition	CDSS use Usual care

TABLE 1 Headline characteristics of included studies (continued)

Author and year	Country	Design	Setting	Number of sites	Study duration	HPs involved	Outcomes	Interventions
Paulson <i>et al.</i> 2020 ⁹⁵	Norway	RT	University hospital	1	10 months	Nurses	Management of malnutrition	CDSS use Usual care
Plank <i>et al.</i> 2006 ⁸¹	Mixed (Austria, Czech Republic, UK)	RT	University hospitals	3	48 hours	Nurses	Glycaemic control	Intervention (CDSS-model predictive control (MPC)) Control (Routine Treatment Protocol (RTP))
Rood <i>et al.</i> 2005 ⁶⁹	Netherlands	RT	Surgical ICU in a teaching hospital	1	10 weeks	Nurses	Glycaemic control	Intervention (CDSS based guideline) Control (paper-based guideline)
Roukema <i>et al.</i> 2008 ⁸⁷	Netherlands	RT	Children's hospital	1	27 months	Nurses	Management of children with fever without apparent source	Nurses (CDSS use) Physicians (CDSS non-assisted)
Sassen <i>et al.</i> 2014 ⁷⁰	Netherlands	RT	University research centre	recruited Online	17 months	Nurses and physios	professionals' behaviour	Intervention (CDSS use) Control (no CDSS use)
Snooks <i>et al.</i> 2014 ⁶²	UK	RT	Emergency ambulance services	13	1 year	Paramedics	Assessment and management of falls	CDSS (used hand-held tablet computers for decisions) Usual care (no CDSS use)
Vadher <i>et al.</i> 1997 ⁸²	UK	RT	Cardiovascular medicine, general hospital	1		A nurse and trainee doctors	oral anticoagulant control	Intervention (nurse with CDSS) Control (trainee doctor without CDSS)
Wells 2013 ⁶³	UK	RT	Emergency ambulance services	13	1 year	Paramedics	Emergency fall assessment and management	CDSS (used hand-held tablet computers for decisions) Usual care (no CDSS use)

61. Bennett P, Hardiker N. A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. *Studies in Health Technology and Informatics* 2016;225:53-7.

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://dx.doi.org/10.1371/journal.pone.0106436>.

63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice. Swansea University (United Kingdom), 2013.
64. Lattimer V, George S, Thompson F, et al. Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;**317**(7165):1054–9.
65. Lattimer V, Sassi F, George S, et al. Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;**320**(7241):1053–7.
66. Cortez S. Measuring active clinical decision support influence on nursing research utilization. University of Phoenix, 2014.
67. Fitzmaurice DA, Hobbs FD, Murray ET, et al. Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;**160**(15):2343–8.
68. Kroth PJ, Dexter PR, Overhage JM, et al. A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *Amia Annual symposium proceedings AMIA symposium* 2006:444–8.
69. Rood E, Bosman RJ, van der Spoel JI, et al. Use of a Computerized Guideline for Glucose Regulation in the Intensive Care Unit Improved Both Guideline Adherence and Glucose Regulation. *Journal of the American Medical Informatics Association* 2005;**12**(2):172–80. doi: 10.1197/jamia.M1598.
70. Sassen B, Kok G, Scheepers J, et al. Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;**16**(10):e211.
71. Cleveringa FGW, Gorter KJ, Van Donk MD, et al. Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;**31**(12):2273–5.
72. Cleveringa FGW, Weising PMJ, Van Den Donk M, et al. Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;**33**(2):258–63.
73. Geurts D, de Vos-Kerkhof E, Polinder S, et al. Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;**176**(2):173–81.
74. Lee NJ, Chen ES, Currie LM, et al. The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, et al. Comparison of Three Protocols for Tight Glycemic Control in Cardiac Surgery Patients. *Diabetes Care* 2009;**32**(5):757. doi: 10.2337/dc08-1851.
76. Cambolat O, Kapucu S, Kilickaya O. Comparison of Routine and Computer-Guided Glucose Management for Glycemic Control in Critically Ill Patients. *Critical Care Nurse* 2019;**39**(4):20–27. doi: 10.4037/ccn2019431.
77. Cavalcanti AB, Silva E, Pereira AJ, et al. A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;**24**(3):371–78. doi: <https://doi.org/10.1016/j.jccr.2009.05.005>.
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;**21**(2):106–15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.
79. Hovorka R, Kremen J, Blaha J, et al. Blood Glucose Control by a Model Predictive Control Algorithm with Variable Sampling Rate Versus a Routine Glucose Management Protocol in Cardiac Surgery Patients: A Randomized Controlled Trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;**92**(8):2960–64. doi: 10.1210/jc.2007-0434.
80. Mann EA, Jones JA, Wolf SE, et al. Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, et al. Multicentric, Randomized, Controlled Trial to Evaluate Blood Glucose Control by the Model Predictive Control Algorithm Versus Routine Glucose Management Protocols in Intensive Care Unit Patients. *Diabetes Care* 2006;**29**(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical and Laboratory Haematology* 1997;**19**(3):203–07.

83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
84. Duclos A, Touzet S, Restier L, et al. Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *European journal of clinical nutrition* 2015;**69**(7):769–75. doi: 10.1038/ejcn.2014.288.
85. Forberg U, Unbeck M, Wallin L, et al. Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implementation Science* 2016;**11**. doi: 10.1186/s13012-016-0375-9.
86. Dalaba MA, Akweongo P, Aborigo RA, et al. Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;**10**(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, et al. Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *Journal of the American medical informatics association: JAMIA* 2008;**15**(1):107–13. doi: 10.1197/jamia.M2164.
88. McDonald MV, Feldman PH, Barron-Vaya Y, et al. Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;**22**(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, et al. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;**50**(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;**19**(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].
91. Dykes PC, Carroll DL, Hurley A, et al. Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;**304**(17):1912–18.
92. Dykes PC, Burns Z, Adelman J, et al. Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA network open* 2020;**3**(11):e2025889-e89.
93. Fossum M, Alexander GL, Ehnfors M, et al. Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;**80**(9):607–17. doi: <https://dx.doi.org/10.1016/j.jimedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, et al. A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of advanced nursing* 2019;**75**(11):3058–67.
95. Paulsen MM, Paur I, Gjestland J, et al. Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. *Clinical Nutrition* 2020;**39**(12):3607–17.

TABLE 2 Risk of bias assessment of RCTs, non-RCTs and CBA studies using the Effective Practice Organisation of Care (EPOC) tool

Author and year	Risk of bias domains and scores										Overall bias score	
	Random sequence generation	Allocation concealment	Baseline outcome measurements similar	Baseline outcome measurements similar	Baseline characteristics similar	Incomplete outcome data	Knowledge of the allocated interventions prevented during the study	Protection against contamination	Selective outcome reporting	Other bias		
Beckman <i>et al.</i> 2013 ⁸⁹	Low	High	Low	Low	Low	Unclear	High	Low	Low	Low	Low	High
Blaha <i>et al.</i> 2009 ⁷⁵	Unclear	Unclear	Low	Low	Unclear	Low	Low	Unclear	Low	Low	Low	Unclear
Byrne 2005 ⁸³	High	High	Low	Low	High	Unclear	Unclear	Low	Low	Low	High	High
Canbolat <i>et al.</i> 2019 ⁷⁶	High	High	Unclear	Unclear	High	Unclear	Unclear	High	Low	Low	Unclear	High
Cavalcanti <i>et al.</i> 2009 ⁷⁷	Low	Low	Unclear	Unclear	High	Low	Unclear	Unclear	Low	Low	Low	High
Cleveringa <i>et al.</i> 2008 ⁷¹	Low	Low	Low	Low	High	Unclear	Unclear	Low	Low	Low	Low	High
Cleveringa <i>et al.</i> 2010 ⁷²	Unclear	Low	Low	Low	High	Low	Unclear	Low	Low	Low	Low	High
Cortez 2014 ⁶⁶	Unclear	Low	High	High	Low	Low	Low	Low	Low	Low	Low	High
Dalaba <i>et al.</i> 2015 ⁸⁶	High	High	High	High	High	Unclear	Unclear	Low	Low	Low	Low	High
Duclos <i>et al.</i> 2015 ⁸⁴	Low	Low	High	High	High	Low	Unclear	Unclear	Low	Low	Low	High
Dumont <i>et al.</i> 2012 ⁷⁸	Unclear	Low	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Low	Low	High	High
Dykes <i>et al.</i> 2010 ⁹¹	Unclear	Low	Low	Low	Unclear	Low	High	High	Low	Low	Low	high
Fitzmaurice <i>et al.</i> 2000 ⁸⁷	Low	Unclear	Low	Low	High	Low	Low	Unclear	Low	Low	Low	High

TABLE 2 Risk of bias assessment of RCTs, non-RCTs and CBA studies using the Effective Practice Organisation of Care (EPOC) tool (continued)

Risk of bias domains and scores										
Author and year	Random sequence generation	Allocation concealment	Baseline outcome measurements similar	Baseline outcome characteristics similar	Incomplete outcome data	Knowledge of the allocated interventions adequately prevented during the study	Protection against contamination	Selective outcome reporting	Other bias	Overall bias score
Forberg <i>et al.</i> 2016 ⁸⁵	Low	Unclear	Low	Low	Low	Unclear	High	Low	Low	High
Fossum <i>et al.</i> 2011 ⁹³	High	High	Low	Unclear	Low	Unclear	Low	Low	Low	High
Geurts <i>et al.</i> 2016 ⁷³	Low	Low	Unclear	Low	Low	Low	High	Low	High	High
Hovorka <i>et al.</i> 2007 ⁷⁹	Low	Low	Unclear	Unclear	Unclear	Low	High	Low	Low	High
Kroth <i>et al.</i> 2006 ⁶⁸	Low	Unclear	Unclear	High	Low	Low	Low	Low	Low	High
Lattimer <i>et al.</i> 1998 ⁶⁴	Low	Low	Unclear	Unclear	Unclear	Low	Low	Low	Low	Unclear
Lattimer <i>et al.</i> 2000 ⁶⁵	Unclear	Unclear	Unclear	High	Unclear	Low	Low	Low	Low	Unclear
Lee <i>et al.</i> 2009 ⁷⁴	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High	Low	Low	High
Lv <i>et al.</i> 2019 ⁹⁴	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	High	Low	Low	High
Mann <i>et al.</i> 2011 ⁸⁰	Low	Unclear	Unclear	Unclear	Unclear	Unclear	High	Low	Low	High
McDonald <i>et al.</i> 2017 ⁸⁸	Low	Low	Unclear	Low	Low	High	Unclear	Low	High	High
Paulson <i>et al.</i> 2020 ⁹⁵	Low	Low	Unclear	Low	High	Low	Unclear	Low	Low	High
Plank <i>et al.</i> 2006 ⁸¹	Unclear	Unclear	Unclear	High	Low	Low	High	Low	Low	High
Rood <i>et al.</i> 2005 ⁶⁹	Low	Unclear	Unclear	Unclear	Unclear	Unclear	High	Low	Low	High

continued

TABLE 2 Risk of bias assessment of RCTs, non-RCTs and CBA studies using the Effective Practice Organisation of Care (EPOC) tool (continued)

Risk of bias domains and scores										
Author and year	Random sequence generation	Allocation concealment	Baseline outcome measurements similar	Baseline outcome characteristics similar	Incomplete outcome data	Knowledge of the allocated interventions adequately prevented during the study	Protection against contamination	Selective outcome reporting	Other bias	Overall bias score
Roukema <i>et al.</i> 2008 ⁸⁷	Low	Low	Unclear	Unclear	Unclear	Unclear	High	Low	Low	High
Sassen <i>et al.</i> 2014 ⁷⁰	Unclear	Low	Low	Low	High	High	Low	Low	Low	High
Snooks <i>et al.</i> 2014 ⁶²	Low	Low	Unclear	Unclear	Unclear	Low	Low	Low	Low	Unclear
Vadher <i>et al.</i> 1997 ⁸²	Low	Unclear	Unclear	Low	Unclear	Low	High	Low	High	High
Wells 2013 ⁶³	Low	Unclear	Unclear	Low	Unclear	Low	High	Low	Low	High

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://doi.org/10.1371/journal.pone.0106436>.

63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice. Swansea University (United Kingdom), 2013.

64. Lattimer V, George S, Thompson F, *et al.* Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;317(7165):1054–9.

65. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;320(7241):1053–7.

66. Cortez S. Measuring active clinical decision support influence on nursing research utilization. University of Phoenix, 2014.

67. Fitzmaurice DA, Hobbs FD, Murray ET, *et al.* Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;160(15):2343–8.

68. Kroth PJ, Dexter PR, Overhage JM, *et al.* A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *Amia Annual Symposium proceedings AMIA symposium* 2006:444–8.

69. Rood E, Bosman RJ, van der Spoel JJ, *et al.* Use of a Computerized Guideline for Glucose Regulation in the Intensive Care Unit Improved Both Guideline Adherence and Glucose Regulation. *Journal of the American Medical Informatics Association* 2005;12(2):172–80. doi: 10.1197/jamia.M1598.

70. Sassen B, Kok G, Scheepers J, *et al.* Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;16(10):e211.

71. Cleveringa FGW, Gorter KJ, Van Donk MD, *et al.* Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;31(12):2273–5.

72. Cleveringa FGW, Weising PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;33(2):258–63.

73. Geurts D, de Vos-Kerkhof E, Polinder S, et al. Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;**176**(2):173–81.
74. Lee NJ, Chen ES, Currie LM, et al. The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, et al. Comparison of Three Protocols for Tight Glycemic Control in Cardiac Surgery Patients. *Diabetes Care* 2009;**32**(5):757. doi: 10.2337/dc08-1851.
76. Canbolat O, Kapucu S, Kilickaya O. Comparison of Routine and Computer-Guided Glucose Management for Glycemic Control in Critically Ill Patients. *Critical Care Nurse* 2019;**39**(4):20–7. doi: 10.4037/ccn2019431.
77. Cavalcanti AB, Silva E, Pereira AJ, et al. A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;**24**(3):371–8. doi: <https://doi.org/10.1016/j.jcrc.2009.05.005>.
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;**21**(2):106–15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.
79. Hovorka R, Kremen J, Blaha J, et al. Blood Glucose Control by a Model Predictive Control Algorithm with Variable Sampling Rate Versus a Routine Glucose Management Protocol in Cardiac Surgery Patients: A Randomized Controlled Trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;**92**(8):2960–64. doi: 10.1210/jc.2007-0434.
80. Mann EA, Jones JA, Wolf SE, et al. Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, et al. Multicentric, Randomized, Controlled Trial to Evaluate Blood Glucose Control by the Model Predictive Control Algorithm Versus Routine Glucose Management Protocols in Intensive Care Unit Patients. *Diabetes Care* 2006;**29**(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical and laboratory haematology* 1997;**19**(3):203–07.
83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
84. Duclos A, Touzet S, Restier L, et al. Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *European journal of clinical nutrition* 2015;**69**(7):769–75. doi: 10.1038/ejcn.2014.288.
85. Forberg U, Unbeck M, Wallin L, et al. Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implementation Science* 2016;**11** doi: 10.1186/s13012-016-0375-9.
86. Dalaba MA, Akweongo P, Aborigo RA, et al. Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;**10**(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, et al. Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *Journal of the American medical informatics association: JAMIA* 2008;**15**(1):107–13. doi: 10.1197/jamia.M2164.
88. McDonald MV, Feldman PH, Barron-Yaya Y, et al. Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;**22**(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, et al. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;**50**(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
91. Dykes PC, Carroll DL, Hurlley A, et al. Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;**304**(17):1912–18.
93. Fossum M, Alexander GL, Ehnfors M, et al. Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;**80**(9):607–17. doi: <https://dx.doi.org/10.1016/j.jimedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, et al. A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of advanced nursing* 2019;**75**(11):3058–67.
95. Paulsen MM, Paur I, Gjestland J, et al. Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. *Clinical Nutrition* 2020;**39**(12):3607–17.

Missing data

We contacted investigators of primary studies to verify study characteristics and obtain missing outcome data where only study abstract or results were presented in published manuscripts. Missing summary data were computed from other reported statistics wherever possible.

Investigating the effects of CDSS characteristics on outcomes

For each included study, we abstracted information on 16 system characteristics associated with effectiveness in CDSS for each study.⁴¹ We classified each as present or absent (the predictor variable). A categorical dependent variable of either 'success' (CDSS better than usual care in at least one of the outcomes reported in each study) or 'failure' (usual care better than CDSS in one of the outcomes reported) was created for each of the 35 included studies. To evaluate whether CDSS-generated outcomes were associated with these characteristics, logistic regression models were constructed using the approach advocated by Firth for generating robust standard errors.⁵⁵ We set a 5% significance level and 95% confidence intervals for each CDSS characteristic.

'Calibration' interviews

We sought to access the reported experience and perceptions of key staff involved in the implementation and use of CDSS in services to sense-check our synthesis results and aid presentation. We (prior to the COVID-19 pandemic) planned a national online survey of UK NHS Chief (Nursing/AHP) Informatics Officers, but attempts at recruitment using NHSE email-based lists and forums and social media were disappointing. The COVID-19 pandemic and redeployment of key staff meant our original approach was not feasible. We eventually identified six key CDSS leaders in a range of organisations and with links to policy as well as delivery: two acute NHS Trusts (one of which was a large teaching hospital); one mixed acute and community semi-rural NHS Trust; an academic health science network lead with links to a large district general hospital-style Trust; a senior policy-level NHS lead for CDSS; a clinical academic with strategic and operational leadership role in a large urban hospital. Their implementation of CDSS varied from 20 years ago (the large teaching hospital) with most in the last three years.

Whilst two of the leaders highlighted specific system-user professional roles (such as 'nurses') or by clinical area (such as 'renal') the rest indicated that a wide multidisciplinary staff base were the intended users – including nurses and AHPs. The systems involved were intended to support a wide range of decisions; for example, disease management, detection, diagnosis, generating treatment options, forecasting/prognosis and triage.

We conducted individual virtual interviews, lasting around 40 minutes to one hour, via Zoom or telephone with our six CDSS leaders using an interview schedule developed to address the four main concepts of NPT (coherence, cognitive participation, collective action, reflexive monitoring) and contextualised for CDSS – as the 'innovation' or new way of working in NPT – and drawing on our main findings as prompts for discussion/sense-checking. Analysis of transcribed data and notes was abductive⁵⁶ and thematic⁵⁷ following the process outlined by Braun and Clarke.⁵⁸ An initial codebook was generated based on NPT constructs and sub-constructs and text read and coded. We used matrices⁵⁹ with NPT constructs as columns, text from each participant as rows before comparison between participants and across columns in a version of metacoding.⁶⁰ Two of the team developed sub-themes from the initial codes of the four NPT key concepts. This was a small scale 'pragmatic' qualitative analysis aimed at helping understand uncertain review findings; we did not carry out inter-coder reliability checks and other qualitative-analytic techniques.

TABLE 3 Risk of bias assessment of interrupted time series studies using the Effective Practice Organisation of Care (EPOC) tool

Author and year	Risk of bias domains and scores							Overall bias
	Intervention independent of other changes	Shape of the intervention effect pre-specified	Intervention unlikely to affect data collection	Knowledge of the allocated interventions adequately prevented during the study	Incomplete outcome data adequately	Selective outcome reporting	Other bias	
Bennet 2016 ⁶¹	High	Low	Low	Low	Low	Low	Low	High
Dykes <i>et al.</i> 2020 ⁹⁶	High	Low	Low	Low	Low	Low	Low	High
Dowding <i>et al.</i> 2012 ⁹⁰	High	Low	Low	Low	Low	Low	Low	High

61. Bennett P, Hardiker N. A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. *Studies in health technology and informatics* 2016;**225**:53–7.

90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;**19**(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].

96. Dykes PC, Burns Z, Adelman J, *et al.* Evaluation of a Patient-Centered Fall-Prevention Tool Kit to Reduce Falls and Injuries: A Nonrandomized Controlled Trial. *JAMA Network Open* 2020;**3**(11):e2025889–e89. doi: 10.1001/jamanetworkopen.2020.25889.

TABLE 4 Summary of patient care process results

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Adherence to guidelines							
Dumont <i>et al.</i> 2012 ⁷⁸	CDSS use	Nurses (OA = 44)	141 adults	Deviations from the protocol, out of 10 (mean (SD))	4 months = 0.39(1.0) 4 months = 3.0(4.3)	-	Mean difference: -2.61 (-4.5 to -0.71)
Forberg <i>et al.</i> 2016 ⁸⁵	Paper protocol	Nurses	159 adults				
	CDSS use	108 nurses	Not applicable	Nurses adherence to guidelines on disinfection of hands	Baseline = 97/108 3 months = 93/105	-1.2%	6.7% (4.9 to 8.5)
	CDSS non-use	103 nurses	Not applicable		Baseline = 96/103 3 months = 87/102	-7.9%	
	CDSS use			Nurses adherence to guidelines on usage of disposable gloves (n/N)	Baseline = 80/108 3 months = 76/105	-1.7%	-1.4% (-2.2 to -0.5)
	CDSS non-use				Baseline = 71/103 3 months = 70/102	-0.3%	
	CDSS use			Nurses adherence to guidelines on daily inspection of peripheral venous catheters (PVC) site (n/N)	Baseline = 58/108 3 months = 58/103	2.6%	-5.2% (-7.1 to -3.3)
Rood <i>et al.</i> 2005 ⁶⁹	CDSS non-use				Baseline = 47/102 3 months = 55/102	7.8%	
	CDSS-based GL	ICU nurses	66 adults	Adherence to insulin dose advice (n/N)	10 weeks = 1818/2352	-	22% (19 to 25)
	Paper-based GL	ICU nurses	54 adults		10 weeks = 1667/2597	-	
	CDSS-based GL	ICU nurses	66 adults	Adherence to the guideline for taking blood samples on time (n/N)	10 weeks = 945/2352	-	4.7% (2.0 to 7.4)
Vadher <i>et al.</i> 1997 ⁸²	Paper-based GL	ICU nurses	54 adults		10 weeks = 922/2597	-	
	CDSS	1 nurse	87 adults	Dose advice 'acceptance' in patients with therapeutic range 2-3	Post-test = 188/214	-	28% (20.4 to 35.5)
	Control	3 trainee doctors	90 adults		Post-test = 145/242	-	

TABLE 4 Summary of patient care process results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
	CDSS	1 nurse		Dose advice 'acceptance' in patients with therapeutic range 3–4.5 (n/N)	Post-test = 160/239	-	-6.2% (-14.7 to 2.2)
	Control	3 trainee doctors			Post-test = 150/205	-	
	CDSS	1 nurse		Interval advice 'acceptance' (%) in patients with therapeutic range 2–3	Post-test = 170/230	-	23.9% (15.6 to 32.2)
	Control	3 trainee doctors			Post-test = 133/266	-	
	CDSS	1 nurse		Interval advice 'acceptance' (%) in patients with therapeutic range 3–4.5	Post-test = 129/239	-	3.9% (-5.4 to 13.3)
	Control	3 trainee doctors			Post-test = 101/202	-	
Patient assessment, diagnosis, and treatment practices							
Bennett <i>et al.</i> 2016 ⁶¹	CDSS use period			Pain assessment	Post-test = 97.7%	-	62.7% (59.6 to 65.8)
	CDSS non use				Pre-test = 35%		
	CDSS use			IV antibiotics in 1 hour for sepsis	Post-test = 5.6%	-	-5.9% (-8.3 to -3.5)
	CDSS non use				Pre-test = 11.5%		
Duclos <i>et al.</i> 2015 ⁶⁴	CDSS	Dieticians	667 children	Investigation of malnutrition aetiology	Post-test = 284/667		21.2% (15.9 to 26.5)
	Usual care	Dieticians	477 children		Post-test = 102/477		
	CDSS	Dieticians	667 children	Managed by a dietitian	Post-test = 305/667		12% (6.3 to 17.7)
	Usual care	Dieticians	477 children		Post-test = 161/477		

continued

TABLE 4 Summary of patient care process results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
	CDSS	Dieticians	667 children	prescribed refeeding protocol	Post-test = 230/667		-4.5% (-10.2 to 1.2)
	Usual care	Dieticians	477 children		Post-test = 186/477		
Geurts <i>et al.</i> 2017 ⁷³	CDSS	Nurses	113 children	Patient consultation time(min)-median (IQR)	Post-test = 136(108)	-	3 minutes
	Usual care	Nurses	109 children		Post-test = 133(92)		
	CDSS	Nurses	113 children	Electrolyte level test	Post-test = 15/113	-	-7.8% (-17.7 to 2.1)
	Usual care	Nurses	109 children		Post-test = 23/109		
	CDSS	Nurses	113 children	Acid-base balance test	Post-test = 13/113	-	-3.2% (-12.1 to 5.7)
	Usual care		109 children		Post-test = 16/109		
	CDSS	Nurses	113 children	Oral rehydration solution (nasogastric tube)	Post-test = 17/113	-	6.7% (-1.6 to 15.2)
	Usual care	Nurses	109 children		Post-test = 9/109		
	CDSS	Nurses	113 children	IV rehydration given	Post-test = 0/113	-	-1.8% (-4.4 to 0.7)
	Usual care	Nurses	109 children		Post-test = 2/109		
	CDSS	Nurses	113 children	Other liquid given	Post-test = 18/113	-	-11.6% (-22.4 to -0.8)
	Usual care	Nurses	109 children		Post-test = 30/109		
Roukema <i>et al.</i> 2008 ⁸⁷	CDSS use	Nurses	74 children	Time spent in ED (minutes), median (IQR)	27 months = 138 (77)	-	15 minutes
	Control	Nurses	90 children		27 months = 123 (96)		
	CDSS use	Nurses	74 children	Time spent in ED for lab test (minutes), median (IQR)	27 months = 140 (68)	-	-20 minutes
	Control	Nurses	90 children		27 months = 160 (98)		
Snooks <i>et al.</i> 2014 ⁶²	CDSS	17 paramedics	436 adults	Mean length of episode of care (minutes)	CDSS vs. control	-	-5.7 min (-38.5 to 27.2) ^b
	Control	19 paramedics	343 adults				

TABLE 4 Summary of patient care process results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Wells 2013 ⁶³	CDSS	22 paramedics	436 adults	Respiratory rate recorded, %	1 year = 405/436	-	-1.2% (-4.7 to 2.2)
	Control	20 paramedics	341 adults		1 year = 321/341		
	CDSS	22 paramedics	436 adults	Pulse rate recorded	1 year = 414/436	-	0.9% (-3.9 to 2.0)
	Control	20 paramedics	341 adults		1 year = 327/341		
	CDSS	22 paramedics	436 adults	Consciousness recorded	1 year = 405/436	-	-5.1% (-7.9 to -2.2)
	Control	20 paramedics	341 adults		1 year = 334/341		
Kroth et al. 2006 ⁶⁸	CDSS use	164 nurses	Not applicable	Proportion of erroneously recorded temperatures	9 months = 248/45823	-	-0.8% (-0.9 to -0.6)
	Control	173 nurses	Not applicable		9 months = 575/44339		
Documenting of events							
Dowding et al. 2012 ⁸⁰	CDSS use	Nurses		Fall documentation ratio	Post-CDSS use vs. pre-CDSS use period	-	1.4 (0.03 to 73.7) ^b
	CDSS non-use	Nurses					
	CDSS use			Hospital acquired pressure ulcer (HAPU) risk documentation ratio	Post-CDSS use vs. pre-CDSS use period		9.1 (1.95 to 42.5) ^b
	CDSS non-use						
Paulson et al. 2020 ⁸⁵	CDSS use	Nurses	44 adults	Documentation of nutritional intake compared to requirements	10 months = 37/44		80% (67 to 92)
	Usual care	Nurses	50 adults		10 months = 2/50		
	CDSS use	Nurses	44 adults	Documentation of a nutritional care plan	10 months = 31/44		54.4% (37.6 to 71.3)
	Usual care	Nurses	50 adults		10 months = 8/50		
	CDSS use	Nurses	44 adults	Documentation of nutritional treatment	10 months = 36/44		23.8% (6 to 41.6)
	Usual care	Nurses	50 adults		10 months = 29/50		

continued

TABLE 4 Summary of patient care process results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Patient referrals							
Snooks <i>et al.</i> 2014 ⁶²	CDSS	17 paramedics	436 adults	Patients referred to falls service	1 year = 42/436 1 year = 17/343		4.7% (1.1 to 8.3)
a calculated from reported information unless stated otherwise; b as reported by study authors.							
61. Bennett P, Hardiker N. A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. <i>Studies in health technology and informatics</i> 2016;225:53–7.							
62. Snooks HA, Carter B, Dale J, <i>et al.</i> Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. <i>PLoS ONE</i> 2014;9(9):e106436. doi: https://dx.doi.org/10.1371/journal.pone.0106436 .							
63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice. Swansea University (United Kingdom), 2013.							
68. Kroth PJ, Dexter PR, Overhage JM, <i>et al.</i> A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. <i>Amia Annual symposium proceedings AMIA symposium</i> 2006:444–8.							
69. Rood E, Bosman RJ, van der Spoel JJ, <i>et al.</i> Use of a Computerized Guideline for Glucose Regulation in the Intensive Care Unit Improved Both Guideline Adherence and Glucose Regulation. <i>Journal of the American Medical Informatics Association</i> 2005;12(2):172–80. doi: 10.1197/jamia.M1598 .							
73. Geurts D, de Vos-Kerkhof E, Polinder S, <i>et al.</i> Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. <i>European Journal of Pediatrics</i> 2017;176(2):173–81.							
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. <i>Am J Crit Care</i> 2012;21(2):106–15. doi: https://dx.doi.org/10.4037/ajcc2012956 .							
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. <i>Clinical and laboratory haematology</i> 1997;19(3):203–07.							
84. Duclos A, Touzet S, Restier L, <i>et al.</i> Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. <i>European journal of clinical nutrition</i> 2015;69(7):769–75. doi: 10.1038/ejcn.2014.288 .							
85. Forberg U, Unbeck M, Wallin L, <i>et al.</i> Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. <i>Implementation Science</i> 2016;11. doi: 10.1186/s13012-016-0375-9 .							
87. Roukema J, Steyerberg EW, van der Lei J, <i>et al.</i> Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. <i>Journal of the American medical informatics association: JAMIA</i> 2008;15(1):107–13. doi: 10.1197/jamia.M2164 .							
90. Dowling DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. <i>J Am Med Inform Assoc</i> 2012;19(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].							
95. Paulsen MM, Paur I, Gjestland J, <i>et al.</i> Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. <i>Clinical Nutrition</i> 2020;39(12):3607–17.							

TABLE 5 Summary of patient care outcome results

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Glycaemic control							
Blaha <i>et al.</i> 2009 ⁷⁵	CDSS (eMPC)	ICU nurses	40 adults	Entire study time in target range (blood glucose) (mmol/l)	After 48 hrs = 46%	-	Versus Mathias: 7.8% (-13.7 to 29.4)
	Mathias protocol		40 adults		After 48 hrs = 38.2%	-	Versus Bath 6.3% (-3.9 to 16.5)
	Bath-protocol		40 adults		After 48 hrs = 39.7%		
Canbolat <i>et al.</i> 2019 ⁷⁶	CDSS (eMPC)	ICU nurses	40 adults	Entire study mean blood glucose (SE) (mmol/l)	Baseline = 8.1(0.6) 48 hrs = 5.9(0.2)	-2.2 mmol/l	Versus Mathias: -1 mmol/l
	Mathias protocol		40 adults		Baseline = 7.9(0.4) 48 hrs = 6.7(0.1)	-1.2 mmol/l	Versus Bath: -0.7 mmol/l
	Bath-protocol		40 adults		Baseline = 8.0(0.2) 48 hrs = 6.5(0.2)	-1.5 mmol/l	
Cavalcanti <i>et al.</i> 2009 ⁷⁷	CDSS (automated BG control)	Nurses	33 adults	Occasions for BG out of target (120 to 180 mg/dl) range	22 months = 2101/5789	-	-21.8% (-23.7 to -20.0)
	Standard protocol	Physicians	33 adults		22 months = 2977/5122		
	CDSS (automated BG control)			Occasions for BG out of target range due to insulin treatment	22 months = 745/5789	-	-28.1% (-29.7 to -26.5)
Cavalcanti <i>et al.</i> 2009 ⁷⁷	Standard protocol				22 months = 2099/5122		
	CDSS (computer-assisted insulin protocol)	ICU nurses	56 adults	Mean blood glucose (mmol/dl)	19 months = 125	-	Versus Leuven -2.1 mmol/dl
	Control (Leuven protocol)	ICU nurses	58 adults		19 months = 127.1	-	Versus conventional -33.5 mmol/dl
	Control (conventional treatment)	ICU nurses	53 adults		19 months = 158.5		

continued

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Cleveringa <i>et al.</i> 2008 ⁷¹	CDSS (computer-assisted insulin protocol)	ICU nurses	56 adults	Patients with hypoglycaemia	19 months = 12/56	-	Versus Leuven -20% (-36.6 to -3.4) Versus conventional 17.6% (5.7 to 29.5)
	Control (Leuven protocol)	ICU nurses	58 adults		19 months = 24/58		
	Control (conventional treatment)	ICU nurses	53 adults		19 months = 2/53		
	CDSS use in diabetic patients	Nurses	1699 adults	A1C < 7%	Baseline = 60.8% 1 year = 68%	7.2%	4.6% (2.7 to 6.5)
	Usual care	Nurses	1692 adults		Baseline = 61.6% 1 Year = 64.2%	2.6%	
	CDSS use in diabetic patients		1699 adults	Systolic BP < 140	Baseline = 41% 1 year = 53.9%	12.9%	10.2% (7.9 to 12.5)
	Usual care		1692 adults		Baseline = 39.5% 1 year = 42.2%	2.7%	
	CDSS use in diabetic patients		1699 adults	Total cholesterol < 4.5 mmol/l	Baseline = 36.2% 1 year = 49.0%	10.5%	3.7% (1.2 to 6.2)
	Usual care		1692 adults		Baseline = 38.5% 1 year = 45.3%	6.8%	
	Hovorka <i>et al.</i> 2007 ⁷⁹	CDSS (eMPC)	ICU nurses	30 adults	Proportion in target range (4-6.1 mmol/l)	48 hrs = 60.4%	-
Usual care		ICU nurses	30 adults		48 hrs = 27.5%		
CDSS (eMPC)				Entire study mean blood glucose (mmol/l) (SD)	48 hrs = 6.2 (1.1) 48 hrs = 7.2 (1.1)		-1 mmol/l
CDSS (eMPC)				Time in target range (hours)	48 hrs = 14.5 48 hrs = 6.6		7.9 hrs

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a	
Mann <i>et al.</i> 2011 ⁸⁰	CDSS use	ICU nurses	18 adults	Occasions glucose range on target (80 to 110 mg/dl)	72 hrs = 47%	-	6% (-7.7 to 19.7)	
	Paper protocol	ICU nurses	18 adults		72 hrs = 41%	-		
	CDSS use	ICU nurses		Occasions over target range (over 110 mg/dl)	72 hrs = 49%	-	-5% (-18.8 to 8.8)	
	Paper protocol	ICU nurses			72 hrs = 54%	-		
	CDSS use			Occasions under target (under 80 mg/dl) range	72 hrs = 4.5%	-	-0.3% (-2.1 to 1.5)	
	Paper protocol				72 hrs = 4.8%	-		
Plank <i>et al.</i> 2006 ⁸¹	CDSS (MPC) use	ICU nurses	Not reported	Occasions within the target glycaemic range (80–110 mg/dl)	48 hrs = 52%	-	33% (20.5 to 45.4)	
	Usual care	ICU nurses	Not reported		48 hrs = 19%	-		
	CDSS (MPC) use	ICU nurses	Not reported	Improvement glycaemic control for 48 hours	48 hrs = 65%	-	40% (27.4 to 52.6)	
	Usual care	ICU nurses	Not reported		48 hrs = 25%	-		
	CDSS (MPC) use		Not reported	Occasions over the target glycaemic range (>110 mg/dl)	48 hrs = 46%	-	-31% (-43.7 to -18.2)	
	Usual care		Not reported		48 hrs = 77%	-		
	CDSS (MPC) use		Not reported	Average glucose (mg/dl)	48 hrs = 117 mg/dL	-	-14 mg/dl	
	Usual care		Not reported		48 hrs = 131 mg/dL	-		
	Blood coagulation management							
	Fitzmaurice <i>et al.</i> 2000 ⁶⁷	CDSS use	Nurses	122 adults	Proportion of tests in range	Baseline = 223/366 1 year = 732/1181	1.1%	-1.9% (-3.1 to -0.7)
	CDSS non-use	Physicians	245 adults		Baseline = 264/480 1 year = 986/1700	3%		

continued

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
	CDSS use	Nurses		International Normalised Ratio (INR) results within range point prevalence	Baseline = 74/118 1 year = 86/121	8.4%	-2.6% (-5.3 to -0.1)
	CDSS non-use	Physicians			Baseline = 129/244 1 year = 157/245	11%	
	CDSS use	Nurses		Time spent within INR target range	Baseline = 64/113 1 year = 76/110	12%	7% (-0.7 to 14.7)
	CDSS non-use	Physicians			Baseline = 99/174 1 year = 143/230	5%	
Antenatal and peripartum care							
Dalaba et al. 2015 ⁸⁶	CDSS use	Nurses	Not reported	Antenatal complications per 1000 attendance	Before = 9 After = 12	0.3%	0.3% (-0.03 to 0.6)
	CDSS non-use	Nurses	Not reported		Before = 16 After = 16	0%	
	CDSS use			Delivery complications per 1000 attendances	Before = 107 After = 96	-0.9%	2.4% (1.1 to 3.7)
	CDSS non-use				Before = 133 After = 100	-3.3%	
Managing patients with chronic co-morbid diseases							
McDonald et al. 2017 ⁸⁸	CDSS use	165 nurses	2550 adults	Medication regimen complexity index <24.5	Post-test = 158/2550	-	0% (-1.1 to 1.1)
	Usual care	335 nurses	5369 adults		Post-test = 333/5369	-	
	CDSS use	165 nurses	2550 adults	Emergency-room use	Post-test = 421/2550	-	-0.2 (-1.9 to 1.6)
	Usual care	335 nurses	5369 adults		Post-test = 897/5369	-	
	CDSS use	165 nurses	2550 adults	Hospitalisation	Post-test = 502/2550	-	-1.4% (-3.3 to 0.5)
	Usual care	335 nurses	5369 adults		Post-test = 1133/5369	-	
Lv et al. 2019 ⁹⁴	CDSS use	Nurses	70 children	Asthma exacerbations (median and inter-quartile range)	Baseline = 9(3) 1 year = 3(2)	-	
	Usual care	Nurses	73 children		Baseline = 9 (4) 1 year = 4(2)	-	

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Outpatient obesity screening							
Lee <i>et al.</i> 2009 ⁷⁴	CDSS use	13 nurses	807 adults	Encounters with obesity related diagnosis	8 months = 91/807	-	10.3% (8.0 to 12.5)
	Usual care	16 nurses	997 adults		8 months = 10/997		
	CDSS use	13 nurses	807 adults	Encounters with missed obesity-related diagnosis	8 months = 51/208	-	-4.1.9% (-48.8 to -35.1)
	Usual care	16 nurses	997 adults		8 months = 440/662		
Fall and pressure ulcer management							
Beeckman <i>et al.</i> 2013 ⁸⁹	CDSS (Pre-vPlan)	65 nurses and physios	225 adults	Pressure-ulcer prevention	Day 1 = 15/58 Day 120 = 41/65	37.2%	2.3% (-11.0 to 15.6)
	Standard protocol	53 nurses and physios	239 adults		Day 1 = 16/63 Day 120 = 41/68	34.9%	
	CDSS (Pre-vPlan)	65 nurses and physios	225 adults	Prevalence of pressure ulcer	Day 1 = 34/225 Day 120 = 16/225	-8%	-6.3% (-10.2 to -2.4)
	Standard protocol	53 nurses and physios	239 adults		Day 1 = 39/239 Day 120 = 35/239	-1.7%	
Byrne 2005 ⁸³	CDSS use	89 nurses	Not reported	Fall rate	Before = 0.312 After = 0.318	0.6%	3.1%
	CDSS non-use		Not reported		Before = 0.315 After = 0.29	-2.5%	
	CDSS use		Not reported	Pressure-ulcer rate	Before = 0.085 After = 0.088	-0.3%	-0.6%
	CDSS non-use		Not reported		Before = 0.091 After = 0.094	0.3%	
Dowding <i>et al.</i> 2012 ⁹⁰	CDSS use			Fall rate	Post-CDSS use vs. pre-CDSS use period	-	0.91 (0.75 to 1.12) ^b
	CDSS non-use						
	CDSS use			HAPU ratio	Post-CDSS use vs. pre-CDSS use period	-	0.47 (0.25 to 0.85) ^b
	CDSS non-use						

continued

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Dykes <i>et al.</i> 2010 ⁹¹	CDSS use	Nurses	5160 adults	Fall rate difference (per 1000 patient days)	CDSS use vs. usual care		-1.16 (-2.16 to -0.17) ^b
	Usual care	Nurses	5104 adults				
Dykes <i>et al.</i> 2020 ⁹²	UDSS use	Nurses	19,283 adults	Fall rate difference (per 1000 patient days)	Post-CDSS use vs. pre-CDSS use period		-0.15 (-0.04 to -0.25) ^b
	CDSS non-use	Nurses	17,948 adults				
Fossum <i>et al.</i> 2011 ⁹³	CDSS use	Nurses	367 adults	Prevalence of pressure ulcers	Before = 16/167 After = 23/200	1.9%	4.2% (0.2 to 8.2)
	CDSS non-use	Nurses	274 adults		Before = 17/150 After = 11/122	-2.3%	
Triaging							
Bennett <i>et al.</i> 2016 ⁶¹	CDSS use period	Nurses	400 adults	Correct triage prioritisation	Post-test = 85.2%	-	24.7% (18.8 to 30.6)
	CDSS non-use	Nurses	400 adults		Pre-test = 60.5%		
Lattimer <i>et al.</i> 1998 ⁶⁴	CDSS	Nurses	Not applicable	Calls managed with telephone advice from GP	Post-test = 1109/7184	-	-34.2% (-35.6 to -32.8)
	Usual care	Physicians	Not applicable		Post-test = 3629/7308		
Lattimer <i>et al.</i> 2000 ⁹⁷	CDSS	Nurses		Patient attended primary care centre	Post-test = 1177/7184	-	-10% (-11.4 to -8.8)
	Usual care	Physicians			Post-test = 1934/7308		
	CDSS	Nurses		Patient visited at home by duty GP	Post-test = 1317/7184	-	-5.5% (-6.9 to -4.2)
	Usual care	Physicians			Post-test = 1745/7308		
Snooks <i>et al.</i> 2014	CDSS	Nurses		Total admissions within 3 days	1 year = 428/7184	-	-0.98% (-1.8 to -0.2)
	Usual care	Physicians			1 year = 507/7308		
Snooks <i>et al.</i> 2014	CDSS	Paramedics	436 adults	Patients left at scene without conveyance to emergency department	1 year = 183/436	-	5.2% (-1.7 to 12.1)
	Control	Paramedics	343 adults		1 year = 126/343		

TABLE 5 Summary of patient care outcome results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
	CDSS		436 adults	Patients with further emergency admission to hospital or death	1 year = 69/436	-	1.5% (-3.5 to 6.6)
	Control		343 adults		1 year = 49/343		
	CDSS			Patients with ED attendance/emergency admission to hospital/death	1 year = 92/436	-	3.3% (-2.3 to 8.9)
	Control				1 year = 61/343		
	CDSS			Patients who reported >1 further fall	1 year = 135/236	-	-6.8% (-16.3 to 2.7)
	Control				1 year = 112/175		
Quality of life and patients' satisfaction							
Cleveringa <i>et al.</i> 2010 ⁷²	CDSS use			Life-years gained	CDSS vs. usual care		0.14 (-0.12 to 0.40) ^b
	Usual care						
	CDSS use			Healthy years (QALYs, discounted)	CDSS vs. usual care		0.037 (-0.066 to 0.14) ^b
	Usual care						
Snooks <i>et al.</i> 2014 ⁶²	CDSS	Paramedics	239 adults	Quality of life (SF12 MCS), mean (SD)	1 year = 41.9(10.3)		-1 (-3.1 to 1.1)
	Control	Paramedics	177 adults		1 year = 42.9(10.9)		
	CDSS	Paramedics	239 adults	Quality of life (SF12 PCS), mean (SD)	1 year = 29(8)		-1 (-2.6 to 0.6)
	Control	Paramedics	177 adults		1 year = 30(8.5)		
	CDSS	Paramedics	228 adults	Patient satisfaction (QC Technical), mean (SD)	1 year = 97.8(10.7)		-0.4 (-2.4 to 1.6)
	Control	Paramedics	165 adults		1 year = 98.2(9.4)		

a calculated from reported information unless stated otherwise; b as reported by study authors.

61. Bennett P, Hardiker N.A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. *Studies in health technology and informatics* 2016;225:53-7.

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://doi.org/10.1371/journal.pone.0106436>.

64. Lattimer V, George S, Thompson F, *et al.* Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;317(7165):1054-9.

67. Fitzmaurice DA, Hobbs FD, Murray ET, et al. Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;**160**(15):2343–8.
71. Cleveringa FGW, Gorter KJ, Van Donk MD, et al. Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;**31**(12):2273–5.
72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, et al. Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;**33**(2):258–63.
74. Lee NJ, Chen ES, Currie LM, et al. The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, et al. Comparison of Three Protocols for Tight Glycemic Control in Cardiac Surgery Patients. *Diabetes Care* 2009;**32**(5):757. doi: 10.2337/dc08-1851.
76. Canbolat O, Kapucu S, Kilickaya O. Comparison of Routine and Computer-Guided Glucose Management for Glycemic Control in Critically Ill Patients. *Critical Care Nurse* 2019;**39**(4):20–7. doi: 10.4037/ccn2019431.
77. Cavalcanti AB, Silva E, Pereira AJ, et al. A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;**24**(3):371–8. doi: <https://doi.org/10.1016/j.jcrc.2009.05.005>.
79. Hovorka R, Kremen J, Blaha J, et al. Blood Glucose Control by a Model Predictive Control Algorithm with Variable Sampling Rate Versus a Routine Glucose Management Protocol in Cardiac Surgery Patients: A Randomized Controlled Trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;**92**(8):2960–4. doi: 10.1210/jc.2007-0434.
80. Mann EA, Jones JA, Wolf SE, et al. Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, et al. Multicentric, Randomized, Controlled Trial to Evaluate Blood Glucose Control by the Model Predictive Control Algorithm Versus Routine Glucose Management Protocols in Intensive Care Unit Patients. *Diabetes Care* 2006;**29**(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
86. Dalaba MA, Akweongo P, Aborigo RA, et al. Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;**10**(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
88. McDonald MV, Feldman PH, Barron-Vaya Y, et al. Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;**22**(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, et al. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;**50**(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowling DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;**19**(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].
91. Dykes PC, Carroll DL, Hurlley A, et al. Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;**304**(17):1912–18.
92. Dykes PC, Burns Z, Adelman J, et al. Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA network open* 2020;**3**(11):e2025889-e89.
93. Fossum M, Alexander GL, Ehnfors M, et al. Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;**80**(9):607–17. doi: <https://dx.doi.org/10.1016/j.ijmedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, et al. A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of advanced nursing* 2019;**75**(11):3058–67.
97. Lattimer V, Sassi F, George S, et al. Cost analysis of nurse telephone in out of hours primary care: Evidence from a randomised controlled trial: BMJ BMJ. *British Medical Journal* 2000;**320**(7241):1053–7.

PPI

In conjunction with our PPI co-applicant (AL) we invited eight members of a single GP practice's Patient Participation Group to a virtual meeting in early 2021. Our PPI co-applicant (AL) hosted the meeting, supported by one of the research team. Participants were sent a description of the purpose and use of CDSS several days before the meeting and asked to consider issues related to patient care and experience of consultations. The practice had a CDSS system embedded into its EHR system. Advanced practitioners, practice nurses and the practice physiotherapist accessed the EHR and CDSS system both during and outside consultations or treatment. In the meeting participants were presented with some of the uncertainties that the team felt were unaddressed by the included studies and offered the chance to ask new questions. The meeting resulted in 12 frequently asked questions (FAQs) that a patient faced with a nurse or AHP using a CDSS might ask. After the meeting, participants were asked to vote via email on the five FAQs they identified as 'most important'.

The project team and the primary research had dedicated PPI expertise from our co-applicant and team member Alison Ledward, an experienced partner in health research and with a background in education and social work. The virtual stakeholders were from a semi-rural area of southern England with only small pockets of socio-economic deprivation. Of note is the almost complete absence of PPI and information related to diversity, inclusion and equality in the research study reports synthesised ([Figure 1](#)).

Findings from the systematic review

From 36,106 initially identified publications, we screened the full text of 262 papers to arrive at our final synthesis of 35 studies. The included studies (see [Table 1](#)) were mainly randomised controlled trials (RCTs) ($n = 28$, 80%) with the other 20% a mix of controlled before-and-after (CBA) studies, interrupted time series (ITS) and a single non-randomised trial (NRCT). Eighty-three per cent ($n = 29$) of the included studies were published after our previous systematic review of decision support in 2007;³⁰ most examined the effects on hospital staff (57%) and Western healthcare systems (USA, UK, Netherlands, Czech Republic and Norway provided the backdrop for 75% of the studies). A single study reported theory to inform the design of the intervention and/or implementation. Just less than a third of the studies (28%) had a published protocol to compare the reported study against.

Who are the users of evaluated CDSS?

Overwhelmingly, evaluations focused on single disciplines using CDSS compared to similar professionals making unaided decisions: nurses ($n = 25$, 71%) and paramedics (6%). Fewer evaluations compared CDSS supported nurses to CDSS unsupported doctors, or a multidisciplinary mix of nurses and physiotherapists in intervention and control groups.

What about the CDSS systems?

Most CDSS come as standalone computer-based systems (89%, $n = 31$) with less than 10% accessed via mobile technology or the web. All the CDSS were 'knowledge-based' (see earlier typology) and whilst single-function systems (such as disease management) were the norm, there were examples of multi-function CDSS (for example, diagnosis and management).

- Triage – five studies in emergency care⁶¹⁻⁶³ and primary care.^{64,65}
- Disease management – five studies: managing cancer symptoms,⁶⁶ oral anticoagulation,⁶⁷ temperature monitoring,⁶⁸ blood glucose monitoring,⁶⁹ and optimising shared decision-making for self-management.⁷⁰
- Diagnosing and managing disease – four studies; diagnosing and treating diabetes;^{71,72} recognising and acting on clinical dehydration in acute gastroenteritis;⁷³ and screening, automated diagnosis, and care-planning for people with obesity.⁷⁴
- Drug dosing – eight studies: mainly in blood glucose control in intensive and emergency environments⁷⁵⁻⁸¹ and oral anticoagulant regimens in hospital cardiovascular patients.⁸²

- Reminder systems – three studies used CDSS for reminders on disease prevention,⁸³ disease diagnosis,⁸⁴ and disease management.⁸⁵ Three others used reminders for multiple functions: disease prevention and management,⁸⁶ disease diagnosis and management,⁸⁷ and disease diagnosis reminder/alert along with disease diagnosis and management.⁸⁸

Are evaluations of CDSS for nurses, midwives and AHPs biased?

With the exception of three RCTs (classed as 'unclear'), all the studies' risks of generating biased conclusions were 'high' (Table 2). The threat of bias did not diminish over time. In RCTs, NRCTs, and CBA studies, sources of bias encountered included no randomisation (13%) or unclear randomisation (27%); unclear (38%) or not done (17%) allocation; only a third of studies ($n = 10$) reported similar baseline measures of outcome and a single study only adjusted their analysis for any differences. Seventeen studies did not specify baseline outcome measurements in their report (57%) and whilst baseline characteristics of providers and patients were similar in around a third of the studies, they differed in another third, and in a further third only patient characteristics were reported – despite the fact that the decisions supported were made primarily by professionals. Half of the 32 randomised studies did not specify missing data and 15 of 32 (47%) studies failed to specify whether CDSS users had knowledge of how they had been allocated to intervention and control groups. In 60% of evaluations (18 studies), 'contamination' was likely or could not be ruled out. More positively, there was no evidence of selective reporting of outcomes. For the three evaluations based around an interrupted time series (Table 3), whilst confounding was an issue, there were no issues with selective outcome reporting, missing data or lack of clarity about when the 'interruption' happened.

The impact of CDSS on performance and outcomes

A broad range of outcomes are used in evaluations of CDSS: 119 outcomes, with 111 different measures. They can be grouped into five areas: (1) care processes, (2) care outcomes, (3) professional knowledge, beliefs and behaviour, (4) safety and (5) economic costs and consequences.

Care processes

There were 34 process outcomes reported. CDSS improved just less than half of these (16/34, 47%) in four evaluations^{69,78,82,85} (Table 4). Conversely, outcomes were worse or no different for 53% of process outcomes (18/34) (Table 5).

CDSS had mixed effects on guideline adherence – sometimes within the same study. In the only trial which took into account baseline and follow-up data,⁸⁵ nurses in *both* arms showed *lower* adherence to hand disinfection (CDSS = -1.2%, Control = -7.9%) and disposable-glove guidance (CDSS = -1.7%, Control = -0.3%), but improved daily inspection of peripheral venous catheters (CDSS = 2.6%, Control = 7.8%). Compared to their non-CDSS-using colleagues, CDSS-using nurses were slightly better at adhering to hand-disinfection guidelines (risk difference = 6.7%; 95% CI: 4.9 to 8.5%) but worse at adhering to policies on disposable gloves (risk difference = -1.4%; 95% CI: -2.2 to -0.5%) and inspection of peripheral venous catheter sites (risk difference = -5.2%; 95% CI: -7.2 to -3.3%).

In trials that did not take into account baseline values,^{69,78,82} CDSS-supported nurses adhered more to guidelines on insulin dosing (risk difference = 22%; 95% CI: 19 to 25%), blood sampling on time (risk difference = 4.7%; 95% CI: 2.0 to 7.4%), and deviated less from protocols (mean score difference out of 10 = -2.6; 95% CI: -4.5 to -0.71)^{69,78} and were more accepting of recommended medication doses than trainee doctors.⁸²

Assessing and treating patients

Six studies^{63,68,73,84,87,61} examined 18 indicators of the quality of patient assessment and treatment. In single studies, CDSS-using nurses assessed pain more readily in emergency department patients (62.7% higher than non CDSS-users [95% CI: 59.6 to 65.8%]) and investigated more paediatric malnutrition aetiology by 21.2% (95% CI: 15.9 to 26.5%), but were slower to provide IV antibiotics within an hour of sepsis onset (5.9% slower, 95% CI: -8.3 to -3.5%). They were no more likely to order laboratory tests

TABLE 6 Summary of health professionals' knowledge, beliefs, and behaviour results

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Mean or risk difference (95% CI) ^a
Beckman <i>et al.</i> 2013 ⁶⁹	CDSS (Pre-vPlan)	65 nurses and physios	225 adults	Positive knowledge change	Baseline = 28/65 5 months = 26/50	8.9%	6.5% (0.8 to 13.2)
	Standard protocol	53 nurses and physios	239 adults		Baseline = 21/53 5 months = 16/38	2.4%	
	CDSS (Pre-vPlan)	65 nurses and physios	225 adults	Positive attitude change	Baseline = 48/65 5 months = 42/50	10.2%	12.7% (5.9 to 19.5)
Cortez 2014 ⁶⁶	Standard protocol	53 nurses and physios	239 adults		Baseline = 39/53 5 months = 27/38	-2.5%	
	CDSS (drop-down boxes)	26 nurses	NA	Research utilisation	Baseline = 35% 11 weeks = 38%	3%	9% (3.3 to 14.7)
Dumont <i>et al.</i> 2012 ⁷⁸	Control	24 nurses	NA		Baseline = 19% 11 weeks = 13%	-6%	
	CDSS use	Nurses (OA = 44)	141 adults	Nurses' satisfaction, out of 10 (mean [SD])	4 months = 8.4(1.4)	-	3.6 (2.4 to 4.8)
	Paper protocol	Nurses	159 adults		4 months = 4.8(2.4)	-	
	CDSS use			Perception of how often needed to deviate from the protocol, out of 10 (mean [SD])	4 months = 2.7(2.2)	-	-4.7 (-6.1 to -3.3)
Sassen <i>et al.</i> 2014 ⁷⁰	Paper protocol				4 months = 7.4(2.4)	-	
	CDSS use	42 nurses and physios	Not reported	Behaviour, mean (SD)	Baseline = 4.5 (1.02) 17 months = 4.6 (0.85)	0.1 (0.93)	0.1 (-0.32 to 0.53)
	Control	27 nurses and physios	Not reported		Baseline = 4.8 (0.69) 17 months = 4.8 (0.82)	0 (0.75)	
	CDSS use	42 nurses and physios		Intention, mean (SD)	Baseline = 6.3 (1.0) 17 months = 6.1 (1.1)	0.2 (1.05)	0.3 (-0.22 to 0.82)
Control	27 nurses and physios			Baseline = 5.9 (1.15) 17 months = 6.0 (0.91)	-0.1(1.05)		

continued

TABLE 6 Summary of health professionals' knowledge, beliefs, and behaviour results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Mean or risk difference (95% CI) ^a
	CDSS use	42 nurses and physios		Attitude, mean (SD)	Baseline = 6.3 (0.44) 17 months = 6.3 (0.56)	0.0 (0.05)	-0.1 (-0.13 to -0.07)
	Control	27 nurses and physios			Baseline = 6.2 (0.69) 17 months = 6.3 (0.68)	0.1 (0.09)	
	CDSS use	42 nurses and physios		Perceived behavioural control, mean (SD)	Baseline = 4.7 (0.79) 17 months = 5.0 (0.73)	0.3 (0.77)	-0.1 (-0.49 to 0.29)
	Control	27 nurses and physios			Baseline = 4.9 (0.87) 17 months = 5.3 (0.8)	0.4 (0.85)	
	CDSS use	42 nurses and physios		Subjective norms, mean (SD)	Baseline = 5.5 (0.55) 17 months = 5.6 (0.63)	0.1 (0.59)	0 (0.34 to 0.34)
	Control	27 nurses and physios			Baseline = 5.6 (0.93) 17 months = 5.7 (0.76)	0.1 (0.84)	
	CDSS use	42 nurses and physios		Moral norms, mean (SD)	Baseline = 6.0 (0.63) 17 months = 6.2 (0.7)	0.2 (0.67)	0.1 (-0.21 to 0.41)
	Control	27 nurses and physios			Baseline = 6.2 (0.59) 17 months = 6.3 (0.55)	0.1 (0.57)	
	CDSS use	42 nurses and physios		Barriers, mean (SD)	Baseline = 3.1 (1.17) 17 months = 3.2 (1.12)	0.1 (1.14)	0.3 (-0.23 to 0.83)
	Control	27 nurses and physios			Baseline = 2.8 (1.01) 17 months = 2.6 (0.96)	-0.2 (0.98)	

a calculated from reported information unless stated otherwise.

66. Cortez S. *Measuring Active Clinical Decision Support Influence on Nursing Research Utilization*. University of Phoenix; 2014.

70. Sassen B, Kok G, Schepers J, et al. Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;16(10):e211.

78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;21(2):106-15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.

89. Beeckman D, Clays E, Van Hecke A, et al. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;50(4):475-86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.

TABLE 7 Summary of adverse events results

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Change of value within a group ^a	Risk difference (95% CI) ^a
Cleveringa <i>et al.</i> 2010 ⁷²	CDSS use in diabetic patients	Nurses	1699 adults	Cardiovascular events occurring	CDSS vs. usual care	-	-11% (-18 to -4) ^b
Fitzmaurice <i>et al.</i> 2000 ⁶⁷	Usual care	Nurses	1692 adults				
	CDSS nurse	Nurses	224 adults	Serious adverse reaction events	1 year = 3 (1.3%)		-5.7% (-10.1 to -1.2)
	CDSS non-use	Physicians	143 adults		1 year = 10 (7%)		
	CDSS nurse	Nurses	224 adults	Deaths	1 year = 3 (1.3%)		-5% (-9.2 to -0.7)
	CDSS non-use	Physicians	143 adults		1 year = 9 (6.3%)		
Snooks <i>et al.</i> 2014 ⁶²	CDSS	17 Paramedics	436 adults	Patients dying	1 year = 19/436 (4.4%)	-	1.2% (-1.5 to 3.8)
	Control	19 Paramedics	343 adults		1 year = 11/343 (3.2%)		

a calculated from reported information unless stated otherwise.

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://doi.org/10.1371/journal.pone.0106436>.

67. Fitzmaurice DA, Hobbs FD, Murray ET, *et al.* Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;160(15):2343-8.

72. Cleveringa FGW, Weising PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;33(2):258-63.

TABLE 8 Summary of analysis results (odds ratio and 95% CI) for the effect of features on effectiveness of CDSS

Factor	CDSS better than usual care ^a	Usual care better than CDSS ^a
Some of study's authors are also systems' developers	0.60 (0.1 to 3.5)	0.23 (0.01 to 4.6)
System provides advice automatically within practitioner's workflow	0.71 (0.16 to 3.2)	0.39 (0.08 to 2.1)
System provides advice at time of care	0.85 (0.18 to 3.9)	0.10 (0.01 to 1.95)
Advice presented in electronic charting or order entry systems	-	-
Provides advice for patients	1.12 (0.04 to 29.9)	1.22 (0.04 to 33.1)
Requires reason for over-ride	1.12 (0.04 to 29.9)	1.22 (0.04 to 33.1)
System facilitates or automates recommended actions	0.30 (0.04 to 2.1)	0.36 (0.02 to 7.5)
Advice is evidence based	2.8 (0.13 to 60.2)	0.49 (0.02 to 10.5)
Critiquing function	-	-
Practitioner does not enter data into system	-	-
Modern system (study published after 2000)	0.26 (0.01 to 5.4)	0.59 (0.07 to 4.8)
Advice or reminders provided directly to patients	1.12 (0.04 to 29.9)	1.22 (0.04 to 33.1)
Trained users	3.08 (0.46 to 20.7)	1.1 (0.2 to 6.0)
Local users were consulted during creation of recommendations	1.94 (0.08 to 44.2)	0.71 (0.03 to 16.4)
System presents its reasoning	-	-
System cites research evidence	-	-

^a Firth's logistic regression.

Note

-, perfectly collinear.

TABLE 9 Features associated with CDSS effectiveness by study

Factors associated with effectiveness of a CDSS																		
Author and year	Some of study's authors are also system's developers	System provides advice automatically within practitioner's workflow	System provides advice at time of care	Advice presented in electronic charting or order entry systems	Provides advice for patients	Requires reason for over-ride	System facilitates or automates recommended actions	Advice is evidence based	Critiquing function	Practitioner does not enter data into system	Modern system (study published after 2000)	Advice or reminders provided directly to patients	Trained users	Local users were consulted during creation of recommendations	System presents its reasoning	System cites research evidence	CDSS better than the usual care in at least one reported outcome ^a	CDSS worse than the usual care in at least one reported outcome ^a
Beckman 2013 ⁸⁹	✓	✓	✓								✓						✓	
Bennet 2016 ⁶¹											✓						✓	
Blaha 2009 ⁷⁵	✓	✓									✓							
Byrne 2005 ⁸³							✓				✓							
Canbolat 2019 ⁷⁶	✓	✓	✓								✓						✓	
Cavalcanti 2009 ⁷⁷	✓	✓									✓						✓	
Cleveringa 2008 ⁷¹											✓							
Cleveringa 2010 ⁷²											✓						✓	
Cortez 2014 ⁶⁶											✓							
Dalaba 2015 ⁸⁶											✓							✓
Dowding 2012 ⁹⁰											✓						✓	
Duclos 2015 ⁸⁴	✓	✓	✓				✓				✓						✓	

continued

TABLE 9 Features associated with CDSS effectiveness by stud (continued)

Factors associated with effectiveness of a CDSS																						
Author and year	Some of study's authors are also system's developers	System provides advice automatically within practitioner's workflow	System provides advice at time of care	Advice presented in electronic charting or order entry systems	Provides advice for patients	Requires reason for over-ride	System facilitates or automates recommended actions	Advice is evidence based	Critiquing function	Practitioner does not enter data into system	Modern system published after 2000	Advice or reminders provided directly to patients	Trained users	Local users were consulted during creation of recommendations	System presents its reasoning evidence	System cites research evidence	CDSS better than the usual care in at least one reported outcome ^a	CDSS worse than the usual care in at least one outcome ^a				
Mann 2011 ⁸⁰											✓							✓				
McDonald 2017 ⁸⁸	✓		✓				✓				✓											
Paulsen 2020 ⁹⁵	✓	✓	✓		✓						✓	✓								✓		
Plank 2006 ⁸¹	✓										✓									✓		
Rood 2005 ⁶⁹	✓						✓				✓									✓		
Roukema 2008 ⁸⁷	✓		✓								✓											
Sassen et al. 2014 ⁷⁰											✓		✓							✓	✓	
Snooks 2014 ⁶²											✓										✓	
Vadher 1997 ⁸²																					✓	
Wells 2013 ⁶³											✓										✓	✓

a outcomes reported as reaching statistical significance.

61. Bennett P, Hardiker N. A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. *Studies in health technology and informatics* 2016;225:53-7.

62. Snooks HA, Carter B, Dale J, et al. Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;4(9):e106436. doi: <https://dx.doi.org/10.1371/journal.pone.0106436>.

63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice. Swansea University (United Kingdom), 2013.

64. Lattimer V, George S, Thompson F, et al. Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;**317**(7165): 1054–9.
65. Lattimer V, Sassi F, George S, et al. Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;**320**(7241):1053–7.
66. Cortez S. Measuring active clinical decision support influence on nursing research utilization. University of Phoenix, 2014.
67. Fitzmaurice DA, Hobbs FD, Murray ET, et al. Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;**160**(15):2343–8.
68. Kroth PJ, Dexter PR, Overhage JM, et al. A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *Amia Annual Symposium proceedings AMIA symposium* 2006:444–8.
69. Rood E, Bosman RJ, van der Spoel JJ, et al. Use of a Computerized Guideline for Glucose Regulation in the Intensive Care Unit Improved Both Guideline Adherence and Glucose Regulation. *Journal of the American Medical Informatics Association* 2005;**12**(2):172–80. doi: 10.1197/jamia.M1598.
70. Sassen B, Kok G, Schepers J, et al. Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;**16**(10):e211.
71. Cleveringa FGW, Gorter KJ, Van Donk MD, et al. Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;**31**(12):2273–5.
72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, et al. Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;**33**(2):258–63.
73. Geurts D, de Vos-Kerkhof E, Pollinder S, et al. Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;**176**(2):173–81.
74. Lee NJ, Chen ES, Currie LM, et al. The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, et al. Comparison of Three Protocols for Tight Glycemic Control in Cardiac Surgery Patients. *Diabetes Care* 2009;**32**(5):757. doi: 10.2337/dc08-1851.
76. Canbolat O, Kapucu S, Kilickaya O. Comparison of Routine and Computer-Guided Glucose Management for Glycemic Control in Critically Ill Patients. *Critical Care Nurse* 2019;**39**(4):20–7. doi: 10.4037/ccn2019431.
77. Cavalcanti AB, Silva E, Pereira AJ, et al. A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;**24**(3):371–78. doi: <https://doi.org/10.1016/j.jcrc.2009.05.005>.
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;**21**(2):106–15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.
79. Hovorka R, Kremen J, Blaha J, et al. Blood Glucose Control by a Model Predictive Control Algorithm with Variable Sampling Rate Versus a Routine Glucose Management Protocol in Cardiac Surgery Patients: A Randomized Controlled Trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;**92**(8):2960–64. doi: 10.1210/jc.2007-0434.
80. Mann EA, Jones JA, Wolf SE, et al. Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, et al. Multicentric, Randomized, Controlled Trial to Evaluate Blood Glucose Control by the Model Predictive Control Algorithm Versus Routine Glucose Management Protocols in Intensive Care Unit Patients. *Diabetes Care* 2006;**29**(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical and laboratory haematology* 1997;**19**(3):203-07.
83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
84. Duclos A, Touzet S, Restier L, et al. Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *European journal of clinical nutrition* 2015;**69**(7):769-75. doi: 10.1038/ejcn.2014.288.

85. Forberg U, Unbeck M, Wallin L, et al. Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implementation Science* 2016;11. doi: 10.1186/s13012-016-0375-9.
86. Dalaba MA, Akweongo P, Aborigo RA, et al. Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;10(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, et al. Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *Journal of the American Medical Informatics Association: JAMIA* 2008;15(1):107–13. doi: 10.1197/jamia.M2164.
88. McDonald MV, Feldman PH, Barron-Yaya Y, et al. Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;22(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, et al. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;50(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;19(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].
91. Dykes PC, Carroll DL, Hurley A, et al. Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;304(17):1912–18.
92. Dykes PC, Burns Z, Adelman J, et al. Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA network open* 2020;3(11):e2025889–e89.
93. Fossum M, Alexander GL, Ehnfors M, et al. Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;80(9):607–17. doi: <https://dx.doi.org/10.1016/j.jimedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, et al. A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of advanced nursing* 2019;75(11):3058–67.
95. Paulsen MM, Paur I, Gjestland J, et al. Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. *Clinical Nutrition* 2020;39(12):3607–17.

(electrolyte levels, acid–base balance test) or nutrition supplements (oral rehydration solution and IV rehydration) in children seen in a paediatric university hospital.

CDSS-enabled nurses recorded fewer incorrect temperatures (risk difference = –0.8%, 95% CI: –0.9 to –0.6%) on wards. CDSS-supported paramedics were no more complete in their assessment of vital signs (respiratory rate, pulse rate and consciousness).

Documenting care

In two single studies, documentation of fall risk (risk ratio = 1.4, 95% CI: 0.03 to 73.7),⁹⁰ pressure-ulcer risk (risk ratio = 9.1, 95% CI: 1.95 to 42.5),⁹⁰ nutritional care planning, nutritional intake and treatment were all better when nurses were using CDSS.⁹⁵

Referring to expertise

Paramedics using CDSS in one study avoided unnecessary use of the ER by referring more patients to a community falls service rather than hospital (risk difference = 4.7%, 95% CI: 1.1. to 8.3%).⁶²

Care outcomes

CDSS were associated with better nurse or AHP influenced outcomes in less than half of the indicators reported (22/54, 40.7%) in six RCTs. In one indicator (delivery complications per 1000 births) CDSS reduced fewer harms than not using CDSS (whose harms also diminished over time).⁸⁶

Blood glucose control

Seven trials^{71,75–77,79–81} reporting on 19 separate indicators of glycaemic control suggest CDSS can improve:

- glucose levels in ICU nurses compared to non-tailored protocols⁷⁵
- proportion of patients with glycated haemoglobin (A1C) <7% (as well as systolic blood pressure and cholesterol levels)
- proportion of patients in target range
- number of occasions in target range and reduce the numbers over the target range
- control over 48 hours.

Blood coagulation management

Whilst CDSS-using nurses generated more ‘tests in range’ than unsupported doctors there were no differences in prevalence of INR ‘in range’ or time spent ‘in range’ between CDSS-enabled nurses and non-using doctors.

Antenatal and peripartum care

A single controlled before-and-after study showed that using a CDSS resulted in less of a reduction in delivery complications than not using one (risk difference = 2.4%, 95% CI: 1.1 to 3.7%) – although users and non-users both improved over time.

Managing those with chronic co-morbid conditions

For those patients with complex, co-morbid, conditions, CDSS use by nurses did not reduce ER use or hospitalisations or lead to rationalised, simpler, medication regimens.

Screening for obesity

A single trial revealed that trainee nurses using a CDSS saw more patients with obesity-related diagnoses and lower numbers of patients with missed obesity-related diagnoses.

Assessing for fall and pressure-ulcer risk factors

These complex and uncertain areas of nursing practice yielded mixed results and effects altered across differing study designs. A trial⁸⁹ saw CDSS users associated with fewer pressure ulcers (although

TABLE 10 Summary of economic costs and consequences results

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Difference (95% CI) ^a	
Cleveringa <i>et al.</i> 2010 ⁷²	CDSS use	Nurses		Diabetes-related costs (excluding CHD)-€ discounted	CDSS vs. usual care	1698.00 (187 to 3209) ^b	
	Usual care	Nurses					
	CDSS use			Cardiovascular disease cost-€ discounted	CDSS vs. usual care	-587.00 (-880 to -294) ^b	
	Usual care						
	CDSS use			Diabetic care protocol cost-€ discounted	CDSS vs. usual care	316.00 (315 to 318) ^b	
	Usual care						
	CDSS use			Total cost-€ discounted	CDSS vs. usual care	1,415.00 (-130 to 2961) ^b	
	Usual care						
	CDSS use			Total costs per QALY gained (Euro)	CDSS vs. usual care	38,243.00 ^b	
	Usual care						
	Geurts <i>et al.</i> 2017 ⁷³	CDSS use	Nurses	113 children	Average emergency department visit cost (Euro)	156.4	0.00
		Usual care	Nurses	109 children		156.4	
		CDSS use			Average diagnostics cost (Euro)	1.09	-0.46
		Usual care				1.55	
CDSS use				Average treatment cost (Euro)	4.48	1.90	
Usual care					2.58		
CDSS use				Average follow-up/hospitalisation (Euro)	134.	26.60	
Usual care					107.4		
CDSS use				Average costs of missed diagnoses/adverse events (Euro)	49.70	-32.10	
Usual care					81.8		
CDSS use			Average cost of CDSS implementation (Euro)	61.95	61.95		
Usual care				0.0			
CDSS use			Overall average cost	408	58.00		
Usual care				350			

continued

TABLE 10 Summary of economic costs and consequences results (continued)

Author and year	Interventions	Health professionals	Patient participants	Outcome measured	Outcome values reported	Difference (95% CI) ^a
Lattimer <i>et al.</i> 2000 ⁶⁵	CDSS	Nurses	Not applicable	Net savings [of CDSS use] in a year (£)	CDSS vs. usual care	13,185 (-77,509 to 123,824) ^b
	Usual care	Physicians	Not applicable			
	CDSS			Cost saved from inpatient stay	CDSS vs. usual care	51,059 ^b
	Usual care					
Snooks <i>et al.</i> 2014 ⁶²	CDSS	Paramedics		Implementing cost of CCDS in one month (in 100s £)	74	74
	Control	Paramedics				
	CDSS			Total cost of implementation in one month (in 100s £)	2,773 2,526	247 (-247 to 741) ^b
	Control					
CDSS	Control			Net resources saved by CCDS per patient year (£)		39 ^b
	Control					
CDSS	Control			Net cost resources saved by CCDS per patient year (£)		208-308 ^b
	Control					
CDSS	Control			Mean length of Job cycle time (minutes)	CDSS vs. control	8.9 min (2.3 to 15.3) ^b
CDSS	Control			Mean length of episode of care (minutes)	CDSS vs. control	-5.7 min (-38.5 to 27.2) ^b
Control						

a outcomes reported as reaching statistical significance.

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://dx.doi.org/10.1371/journal.pone.0106436>.

65. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;320(7241):1053-7.

72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;33(2):258-63.

73. Geurts D, de Vos-Kerkhof E, Polinder S, *et al.* Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;176(2):173-81.

non-users' also saw pressure ulcers reduced by a smaller amount). CBA studies saw *more* patients with pressure ulcers amongst CDSS users⁹³ but lower levels of malnutrition in patients cared for by CDSS-supported nurses⁹³ or no difference (in pressure ulcers or falls) in CDSS users and non-users.⁸³ In the single time series study that exists, where CDSS implementation constituted the interruption, there were fewer pressure ulcers or falls when nurses were using the decision support.

Triage

CDSS-using nurses and paramedics using CDSS made fewer calls needing advice from general practitioners (GP) (risk difference = -34.2%, 95% CI: -36 to -33%), lower numbers of patients visited at home by a GP (risk difference = -5.5%, 95% CI: -6.9 to -4.2%), and fewer admissions to hospital within 3 days of nurse input (risk difference = -0.98%, 95% CI: -1.8 to -0.2%). They were also no more likely to 'leave a patient at the scene without conveying to an emergency department' (risk difference = 5.2%, 95% CI: -1.7 to 12.1%). The proportion of 'correct' triage prioritisation judgments was higher when professionals used CDSS (risk difference = 24.7%; 95% CI: 18.8 to 30.6%).

Health-related quality of life and satisfaction with care

Only two studies attempted to measure HRQOL and one also reported patient satisfaction with care; neither study demonstrated differences in HRQOL or patient satisfaction attributable to CDSS use.

Do CDSS increase knowledge and shape positive behaviours?

Professions exist in part due to their own claims of specialist – superior to non-professional – knowledge and modes of behaviour.⁹⁸ It follows then those professional decisions are in part shaped by their knowledge and associated with behaviours. CDSS as a technology to support professional decision impacts on only some of the component parts of professional knowledge and behaviour ([Table 6](#)).

CDSS, on the basis of four RCTs, positively influenced perceptions of frequency of needing to deviate from protocols, positive knowledge and attitude changes. Conversely, measured knowledge itself and key elements of 'planned behaviour'⁹⁹ (intention, attitude, self-efficacy, and subjective and moral norms) were no different in CDSS-using professionals and non-users. Using more research knowledge in decisions, as a consequence of CDSS exposure, was equivocal: one RCT suggests it can be improved, whilst studies with nurses and physiotherapists suggests not.

Do CDSS improve safety in nurse, midwife and AHP performance?

CDSS do not by default make services safer ([Table 7](#)). The complex socio-technical systemic location of CDSS and the work required to embed and sustain systems mean unintended consequences are not uncommon. Aside from simply providing 'ineligible' or suboptimal advice, unintended consequences can also include (1) errors in information entry and retrieval associated with a poor human-computer interface, and (2) inflexible digital systems leading to errors in coordination and communication.¹⁰⁰ Of course, errors also occur in unsupported practice. CDSS, however, have the potential to hardwire such errors into the socio-technical system of clinical practice, making mistakes systematic and systemic. Iatrogenic harm can also arise from ignoring the complex socio-technical system that the technology must operate in: e-iatrogenesis.^{38,101} No studies focused specifically on the effects of CDSS use on patient safety in hospital care. In primary and first-response care, CDSS use by nurses is associated with lower probability of cardiovascular events in people having their diabetes managed by nurses (as opposed to CDSS-unsupported physicians) – an 11% risk difference (95% CI -18 to -4). However, serious adverse incidents and deaths in people who had fallen and were attended by CDSS-informed paramedics were unaffected.

The effects of 'good' design features in CDSS

We examined the effects of incorporating 16 design or system features known to positively influence performance and outcomes in 32 of the studies reported⁴¹ ([Tables 8 and 9](#)). None were associated with greater effectiveness of the CDSS and all were highly uncertain (as seen in their wide 95% CIs in [Table 8](#)).

Economic costs and benefits

Four randomised trials^{62,65,72,73} included 20 economic indicators of costs and consequences (Table 9). The costs of managing cardiovascular disease were lower in CDSS-using groups (cost difference = -€587, 95% CI -880 to -294) but cost more when supporting diabetes care protocol implementation (cost difference = €326, 95% CI 315 to 318). Clinical work took longer: 'mean length of job cycle time' was significantly higher (difference in minutes = 8.9, 95% CI 2.3 to 15.3) in CDSS users (see Table 9). Cost per quality adjusted life-year (QALY) was €38,243 (£32,333) – more than NICE's normal cost-effectiveness threshold of £20,000 to 30,000 per QALY gained for implementation in the UK NHS (Table 10).

Insights from developers, intended users and implementers of CDSS

Coherence

Coherence, or the extent to which an intervention is understood as meaningful, achievable and desirable, is an important factor in adoption. Participants emphasised a number of key differences from CDSS-unsupported care and used these to increase understanding and help encourage professional adoption:

- Reducing documentation whilst simultaneously utilising best evidence. One example was encouraging appropriate pain assessment whilst assessing pressure-ulcer risks.
- CDSS as 'teacher' – forcing users to engage with new information – such as guidelines – in practice.
- Making decisions 'visible' and facilitating appropriate action – an example being greater propensity for escalation in response to NEWS score above the Trust's threshold for action.
- Technology making information more 'palatable' to younger clinicians who were perceived as having preferences for technology-delivered information.
- CDSS facilitating QI and audit and accreditation to a desired standard: something that was much harder prior to the technology being introduced. An example provided was the measurement of 'care not done' such as a Waterlow pressure risk assessment within four hours of admission.
- Standardisation – of care planning, collecting and organising data. Examples would be the assessment, recording and suggested actions associated with the North American Nursing Diagnostic assessment system.

The extent to which these perceived advantages were shared by everyday users of the system was unknown. On probing, it was clear that leaders recognised variability in take-up and adoption, implying not all clinicians may feel the same way. Variation was often attributed to user characteristics such as age ('older users being reluctant') and clinical role ('pharmacist *reviewing* medications invokes less CDSS-use than *initiating* medications') rather than the systems themselves or the fit between decision-work and system suitability.

Interviewees highlighted the importance of fostering shared understanding of CDSS functions, stated benefits, limitations and links to concepts such as safety, accountability and (unwarranted) variability as levers for adoption and sustainability. One system developer and evaluator emphasised the understanding required to know when *not* to use the CDSS:

*it's critical that the users of any of these tools understand the limitations. So, they understand when to use it when **not** to use it (participant's emphasis). So, in that patient in front of them, how do they assess how safe and appropriate it is to use that tool?*

Participant 4. National Policy Team on Digital Health

Collective action

In terms of the work required to bring the CDSS into use (collective action), it was clear that for all the interviewees that adjustment and pragmatic relaxing of some prerequisites (such as the overriding of

alerts) – arising from reflexive monitoring – was required to sustain and encourage use. Implicit in many accounts was a recognition that systems sometimes possessed, and generated, unwanted characteristics that needed active countering. Examples included:

- fine tuning or muting alerts to reduce alert fatigue
- ensuring that over-riding a CDSS recommendation was handled ‘sensitively’ in any post-decision scrutiny – healthcare is uncertain and sometimes over-riding may be necessary or appropriate given the uniqueness of the individual to whom the CDSS is applied
- guarding against ‘garbage in and garbage out’ in both knowledge-based (poor-quality protocols) and machine learning systems (poor-quality ‘training’ data).

The extent to which CDSS developers and proponents expected work to adapt to the CDSS (as opposed to the CDSS having the flexibility to fit into work-as-done) was a strong thread in all the interviews. All the participants reported an awareness of ‘workflow’ (variously referred to as ‘context’, ‘work’, ‘environment’ or ‘practice’) as an important component in the adoption and enactment of CDSS advice.

For example, we had some clinical decision support in a discharge planning assessment ... Depending on the boxes [ticked] and the information provided, we gave it some guidance and then if it was greater than a score of 10 it would create a referral to the discharge team, to help with complex discharge planning. Now staff didn't even use the discharge planning assessment, never mind the associated decision support associated with it.

Participant 1. CNIO large metropolitan teaching hospital

A pharmacist system-developer relayed the importance of understanding the decision tasks and ‘work’ to which the CDSS was to be applied.

Or they may split the work amongst the team so that instead of having one nurse doing a drug round for all 28 patients, they might have four nurses, only giving drugs to six or seven patients. So I think when you're looking for from my perspective I look at medication safety and I always try to understand, well, if this is happening what are the things that are contributing to that? It's not just the individual, you know? Yes, there clearly is an individual aspect into this. But is it to do also with them finding and retrieving medicines? Is it to do with interruptions within the environment? Is it to do with the team dynamics the way they work? Is it to do with the trolley, so you know some of the equipment that they use actually isn't conducive to finding medicines, is not arranged in the right way, so ... trying to understand how they interact with the environment around them to do the task at hand.

Participant 2. CNIO large semi-urban NHS acute trust

All the accounts highlighted the need for supportive infrastructure and new ‘roles’ to enable sustained use of the systems.

What we need is more infrastructure and resources to support better implementation of these technologies, and these are tools, these are absolutely our tools, but they need people to use these tools properly. You can't just give it to the organisations, so we need more specialists who are able to look at the data and analyse who can develop different ways of configuring the system and adapt it to what we need.

Participant 2. CNIO large semi-urban NHS acute trust

Cognitive participation

To encourage cognitive participation, interviewees emphasised formal mechanisms for engagement and increased understanding: training, education, professional committees in areas such as guidance development (rule/knowledge-based systems) and implementation. The accounts revealed a strongly multidisciplinary work environment and decisions that involved multiple professions. However, the training, the committees, and the support infrastructure were almost always mono-professional: all of the respondents' accounts relayed work in which nurses and AHPs primarily restricted their work scope and communications to other nurses and AHPs.

All the interviewees had roles that involved responsibility for either CDSS introduction or maintaining and/or developing it in services. Aside from the aforementioned mono-professional focus, all discussed strategies that seemingly involved 'listening' to users, and encouraging 'buy-in', whilst simultaneously emphasising the advantages of 'forcing' CDSS use as part of work flow:

Before [CDSS], we were relying on people's memory, how to manage, literally more than one patient every five minutes arriving at triage for example. So, it did change the way people work because it forced them to assess people in a set way and always ask the same questions.

Participant 4. National Policy Team on Digital Health

Reflexive monitoring

Various accounts relayed this as a valuable mechanism and 'reality check' on assumed implementation and the use of formal methods to encourage cognitive participation in clinicians. All the accounts highlighted training and training materials as 'intended' mechanisms for supporting initiation and use. One pharmacist's account, though, revealed both the limitations of self-reported behaviour and the benefits of evaluating efforts:

The thing I would say though, while there's a lot of these resources available, I don't know if everybody is accessing it. Certainly, our junior doctors, we did a survey a little while back to try and understand how supported they were in terms of prescribing safely on the electronic system and a lot of them thought they were quite happy. They felt quite confident about using the system to prescribe safely and accurately but when we asked them about training types and resources available, a lot of people did not know about them or never used it or didn't attend the classroom training.

Participant 2. CNIO large semi-urban NHS acute trust

Exogenous factors outside formal strategy were highlighted by a participant CNIO in a large metropolitan teaching hospital with almost 100% adoption of an EHR with decision support capabilities. Strategically, the historic roll-out of a trust-wide EHR system was seen as only partly successful; but the COVID pandemic had catalysed pragmatic adoption and positive feedback. One participant suggested that the CDSS and EHR adoption process was sped up by '5 years'. The technology was also being used by the trust to prevent – or at least question – returning to 'business as usual' even as the pandemic context changed and more face-to-face care and treatment were possible.

Despite informal reflexive monitoring (primarily communal and individual appraisal) figuring strongly in accounts, the relative absence of formal systematisation-evaluation in accounts was striking. Only one of the interviewees referred to formal evaluation of CDSS effects on work and outcomes: a time series evaluation of a system – they had developed – focused on a relatively rare disease in the emergency department; perhaps tellingly, they were a system developer and a clinical academic. Two CNIOs mentioned 'informal evaluation', mainly the collection of anecdotal evidence and committee discussion of first-hand experience of use and adoption. Despite the lack of formal evaluation, participants recognised the *need* for rapid and relevant evaluation – often in the form of real-time monitoring:

If 80% of nurses are clicking past and just ignoring it, what's the clinical risk in that? Do we need to go and change practice and process and education to train and encourage people to actually read these things? And I don't think we do enough of that yet.

Participant 1. CNIO large metropolitan teaching hospital

Where it's really important like NEWS scoring, they also need to be monitored for compliance. It's no point having CDSS if you're not going to monitor if they're effective.

Participant 3. CAHPIO mixed acute and community, semi-urban NHS trust

Other insights

As with studies in the synthesis, users' accounts of system development and implementation did not contain references to formal or explicit theory. Phrases such as 'listening to users', 'understanding the work' and 'alert fatigue being counter-productive' implied some awareness of theoretically important concepts, but none of the stakeholders located their points within relevant theories of systems or implementation.

The dangers of ignoring interactional workability and unwanted artefacts of CDSS, such as generating more work for other parts of the healthcare system, were highlighted by one leader. To the extent that adoption itself can be compromised:

You do a nutritional screening tool and then the dietician says 'stop doing it because actually we're getting too many referrals', that's not the right reason to give it up; and that for me is about how you use the data to inform your future and your resourcing and your requirements and escalating risk where appropriate.

Participant 3. CAHPIO mixed acute and community, semi-urban NHS trust

Standardisation, safety and applying research evidence were the reported primary drivers for initiation, although the extent to which variability, safety and lack of evidence use were historical problems in their organisations – or indeed improved with the advent of CDSS – was not clear.

CDSS as a vehicle for improving knowledge application to practice was highlighted. Whilst two participants referred explicitly to research-based knowledge in the guidance underpinning the CDSS recommendations, for others it was codified, collective (committee) knowledge that was the basis for advice.

Where participants highlighted economic aspects of CDSS use, they generally focused on cost-effectiveness – but in the lay (costs-consequence) rather than formal economic sense (£ per QALY). One participant highlighted how economic context can hinder implementation:

You get paid a set amount for each out-patient appointment that's face-to-face ... let's say it's £100 for each patient, and a follow-up is £60 per patient. But there was a lot of, 'so what is this, it's not face-to-face physically but it's face-to-face virtually, is that still classed as a face-to-face appointment, is there a different cost associated. A lesser cost associated with a virtual consultation than a physical one?' So, it's not all just about tech it's about cost as well.

Participant 1. CNIO large metropolitan teaching hospital

Potential recipients of CDSS-mediated advice

As part of the process of calibrating the results with users (a form of stakeholder engagement) we used a voting process to derive the 'top 5' (see Box 1) frequently asked questions (FAQs) that might act as prompts for conversations between CDSS professional users and/or those in receipt of, or participating in, CDSS-enabled healthcare decision-making.

1. Does the CDSS make the decisions for my nurse/AHP and is my care now decided by a computer?
2. Will CDSS mean that I will have less time talking to someone?
3. Are nurses/AHPs appropriately trained to use CDSS?
4. Can I check that the CDSS is using correct information about me (e.g. the medications I take/up-to-date information on my medical history)?
5. Will the CDSS share my information with anyone else (e.g. Big Pharma)?

BOX 1 Top five questions from CDSS-care recipients

None of the studies included in the synthesis reported addressing these uncertainties from the perspective of people likely to receive the CDSS-enabled care as part of the implementation of the systems.

These five FAQs, whilst useful as a potential start point for further (primary) research, must be interpreted cautiously and are not indicative of representative uncertainties. The aim of engaging with the users was to sense-check our results; the questions did not arise from adequately scaled, sampled, or theoretically scrutinised research efforts.

Discussion

Reasons often cited for CDSS – and present in our stakeholder accounts – include safety, reducing unwarranted variation in practice and outcomes, and greater efficiency and effectiveness associated with the application of evidence to patient care. Our results suggest CDSS cannot yet be relied upon to make healthcare safer or increase standardisation or the application of high-quality research (or indeed, other kinds of knowledge) in nurse and AHP-led decision-informed healthcare. Whilst services *may* be safer, vary less and evidence feature in choices in CDSS-enabled care *some* of the time, we still do not necessarily know why. Evidence on efficiency is extremely sparse and inconclusive.

Two specific issues highlighted in our previous review remain relevant. First, knowledge-based CDSS need high-quality knowledge if they are to improve processes and outcomes. Until AI and machine learning becomes more widespread in NMAHP decision-making, knowledge-based systems will rely on protocols, algorithms and ‘if-then-else’ rules. The evidential basis for many of these rules and protocols is often unclear. CDSS need populating with well-evaluated rules in which the clinical significance of any pre-CDSS evaluation of rules, logic and generative mechanisms for the knowledge underpinning systems is established. CDSS are a form of ‘complex intervention’ and whilst guidance on developing complex interventions was less developed in 2009, the latest iteration of guidance makes the necessity of testing logic and mechanisms before adding more complexity by translating evidence into technologies explicit.¹⁰⁴

Second, the need to unpack the heterogeneity and mixed results by examining the effects of components of systems remains. We looked at features known to be associated with effectiveness in CDSS aimed primarily at doctors (see [Table 8](#)). There was little impact on NMAHP-focused CDSS. Other quality improvement and implementation behaviour-change methods – notably audit and feedback – have recognised that using theory explicitly in design and evaluation^{105,106} and conducting iterative programmes of head-head evaluation of key modifiable characteristics can lead to stronger and more sustained implementation of technologies.¹⁰⁷ We think this iterative development in NMAHP-focused CDSS would arguably lead to compounded learning and more efficient implementation of more effective systems.

The variable adoption of systems evaluated in our included studies suggests that CDSS are not uniformly acceptable to all clinicians. Many of the systems, and some of our stakeholder interviews, imply that CDSS often appear as a non-negotiable aspect of works experienced *after* the decision has already been made to adopt a system – usually by managers who will not have to work with the system for real-time decision-making. We found no instances of outright resistance to the imposition of CDSS in the primary studies. This was not surprising. Dune in his study of evaluations of CDSS systems in the NHS also found that whilst outright resistance was rare, other forms of sub-optimal engagement were more common. Clinicians doing the ‘bare minimum’, simply ‘ticking boxes’, deferring assessments (and avoiding computerised input), or providing poor excuses for missing specified time limits for activities were all features of professional-culture-influenced, subtle, resistance.¹⁰⁸

What’s changed since the last review of CDSS and NMAHPs?

The volume of studies focusing on CDSS has tripled since the last systematic review undertaken with a similar focus in 2009. The body of comparative research now contains a number of randomised comparisons between CDSS and unsupported practice, which is welcome. However, just less than half of the 35 included studies occur in a single site and all remain at high risk of biased conclusions. The increased number and comparative nature of studies has *not* led to more information to inform decision-making – whether as a clinical decision-maker or someone with responsibility for commissioning, implementing or purchasing systems in healthcare environments.

The studies included in the review suggest that CDSS have potential to improve both processes and outcomes – especially in nursing, and a more limited range of professions allied to medicine – for specific

decision-focused work. However, the limited empirical evidence means that the reproducibility and systematic realisation of this potential are hard to predict. Aside from the limited and contradictory empirical research, systems and evaluations lack an explicit theoretical basis. The net effect is that the evidence base still feels essentially scattergun as a corpus of work on which to base local improvement efforts. When systems work (which, in our review, is approximately half of the time) the reasons why can only be speculated upon. But as importantly, when these systems fail (which, in our review, was approximately half of the time) the often considerable financial and opportunity costs are difficult to justify as – to use an oft quoted idiom in healthcare – lessons cannot be learned. Lists of features known to increase effectiveness of CDSS systems^{41,42} are only partially useful in a complex socio-technical context in which professional knowledge, work as done (rather than work as imagined by CDSS designers or implementers), interdisciplinary norms and dynamics influence technology use and decision-making itself. Varghese and colleagues in a review of CDSS suggest uncertain and dynamic clinical environments shape the significant differences in impact of CDSS.¹⁰⁹ Many studies – 73% in Varghese *et al.*'s review – are single-site studies from a single unit, practice or hospital.¹⁰⁹ We might exhort designers, implementers and evaluators to consider the system-level factors that might generate differences in organisational dynamics, but it is only middle-range theories (such as NPT or NASS) that allow technologists, clinicians and researchers to think systematically about the diagnostic, remedial, proactive and evaluative tweaks needed to repair, improve and optimise systems.

For complex interventions such as CDSS, the context surrounding the system is as important as the mechanisms by which they generate outcomes. Whilst not a feature of the shifting research evidence base, the *context* for technology use in healthcare has changed dramatically since spring 2020 and the COVID-19 pandemic. Technologies that were an exceptional component of service delivery, such as video consultation, have become part of mainstream service provision – rapidly accepted and seemingly sustained.¹¹⁰ The evidence in our synthesis has not captured this shift, but it is possible that many of the barriers to implementation or positive effects may be less important in the current delivery context.

Study strengths and weaknesses

A strength of our review is our inclusive approach to searching for a broad range of studies of differing research designs to reflect the potential breadth of nursing, midwifery and AHP decision-informed care and treatment. We have also undertaken a tighter – methodologically speaking – systematic review focusing only on randomised controlled comparisons:¹⁰² the results are not substantially different. This suggests that risk of bias associated with study design is unlikely to be driving the overall research picture.

A related strength of the synthesis overall was the, limited, use of calibration interviews. These helped us unpack some of the apparent effects and both reinforced the importance of good design and implementation and served to highlight the relative absence of both in the included studies.

Including nurses, midwives and AHPs meant that differences in decision context could be examined. Specifically, we thought that certain uncertainty-related decision types (such as triage) or decision components (recording assessments) might be able to be controlled for and we would be able to look for differences between professions faced with similar uncertainties or tasks and using similar CDSS. To some extent this was realised – for example, in relation to triage. But the dearth of inter-professional studies for similar work using CDSS prevented any meaningful comparison. To the best of our knowledge this attempt to control for similar decisions in different contexts with the same system has only been undertaken once by Pope and colleagues in a qualitative case study of CDSS aimed at nurses in emergency and urgent care.⁴⁸ They found that consistency in the use of CDSS (i.e. as the designers intended) was as dependent on the implementation and resourcing to support the work required to enact the system as the technology itself. A theme that was apparent in our stakeholder accounts and also a viable hypothesis arising from the studies included in our synthesis is the lack of explicit theory used to design or evaluate reported implementation.

This was the first large-scale review of CDSS technology for both nurses and AHPs since 2009. We envisaged a larger number of studies, reflecting advances in computing power and CDSS types

(e.g. machine learning). In contrast to the rapid development of machine learning and changes to tech-enabled choices in the commercial world (such as predictive analytics from Amazon, Google etc.) the research evidence for CDSS in nursing, midwifery and AHPs seemed somewhat dated and lacking behind wider societal technology adoption. Nonetheless, our findings are based on three-times the number of studies in our earlier review, suggesting further and faster growth in the evidence base may be possible.

Viewing our synthesis through the lens of NPT proved feasible and highlighted issues that may have been missed if we focused only on our results in a pragmatic way. Thinking of implementation using higher-order concepts as part of systems development and implementation planning may be more useful than trying to simply 'build in' system characteristics. Whilst these are associated with positive outcomes in the wider CDSS literature they were not indicative of improvement in studies of nursing, midwifery and AHPs – suggesting, it is work/roles/professional lines of demarcation that matter more than the architecture or ingredients of decisions. Interviews with system developers, adopters and leaders suggest that – whilst the language may not map onto NPT terminology exactly – concepts of coherence (understanding/sense-making), cognitive participation (building and sustaining), collective action (enacting) and reflexive monitoring (evaluation) are present in accounts.

As with any review of research evidence, our synthesis is a function of the evidence. Research into CDSS aimed at nurses, midwives and AHPs is overwhelmingly low quality, with an absence of theoretical perspectives on design or implementation that could help explain apparent effects/failure or which others could systematically build on. Empirically, studies overwhelmingly focused on nurses and 'nursing decisions' in a professionally demarcated way. This is unfortunate as healthcare work is essentially multi-disciplinary and team-based – especially in acute environments. The evidence on the economics of CDSS aimed at nurses and AHPs is patchy and scarce. Our synthesis is unable to adequately answer policy questions of value-for-money associated with systems.

A key working assumption behind the review was that the comparative research evidence base would have matured sufficiently in quantity and quality to produce a substantive series of 'effects' and 'differences' between systems with less uncertainty than our last review. We felt describing these effects (and associated uncertainty) in a rigorous and systematic way, and having sufficient volume of similar research – amenable to statistical meta-analysis – would offer useful insights for decision-makers faced with commissioning and choosing CDSS. This assumption did not hold up.

We searched only for comparative research studies and failed to include study designs that – whilst less suited to statistical synthesis – might have yielded a richer insight into the implementation processes and mediating effects, mechanisms and contexts that were clearly influential in explaining the mixed picture of uncertain results. As the evidence base matures and the use of theory and well-conducted process evaluations increases, alternatives to traditional systematic review approaches – for example, realist synthesis – may provide more insight and actionable recommendations for implementation, particularly as a basis for nested evaluation alongside CDSS roll-outs.

Beyond reductionist ideas of 'satisfaction' with systems, a sophisticated and nuanced view of user perspective was largely absent in studies. Studies since the 1980s have highlighted the complex way in which people interact and socially create meaning, satisfaction, justifications and reinforcement for their use of technology generally and CDSS specifically.^{48,103} This is not a trivial omission; how people feel about systems influences their later behaviours and the consequent effects of systems. Artificially simplifying user perspectives on CDSS use to abstract concepts such as 'satisfaction' has limited utility for those seeking targets for optimising and evaluating systems.

The lack of common reporting standards for CDSS made comparison between studies to assess similarities difficult and we cannot rule out the possibility that there was more homogeneity between systems and therefore greater potential for statistical synthesis. In the absence of detailed, structured, reporting we were conservative in our estimates. Out of pragmatic necessity, we abstracted descriptions and system features and effects to higher-order concepts such as 'knowledge-based' (features) or 'care processes' (outcomes). This meant discarding potentially 'richer' information. Similarly, studies rarely addressed issues of context and

exogenous factors shaping use, implementation and effects. This meant issues of 'context' were hard to unpack and incorporate into our analysis. Other kinds of reviews – for example, realist syntheses – may have produced more information on what works for whom, when and in what circumstances.

A final weakness relates to our limited range of interviews with what can be termed 'stakeholders' in CDSS. Our original plans had to be changed as a result of lack of engagement arising from the redeployment of staff (and changed priorities) arising from the COVID-19 pandemic. Consequently, the numbers, extent and richness of the data were less than we would have liked. Nonetheless, the limited work yielded some valuable insights and the approach was promising and we will seek to maximise this technique in future syntheses generally and seek approvals and funds to try again with the communities of practice that surround CDSS design, adoption and implementation.

What could we have done differently?

We started from an optimistic estimate of fit-for-purpose research evidence and sufficient quality for inclusion. Our initial pre-synthesis exploratory work suggested substantially more evaluations than when we undertook our last similar review in the 1990s. This sanguine position was justified on the grounds that we expected the numbers of evaluations of CDSS to mirror wider technological and digital expansion in healthcare; for example, EHRs. The growth – at least rhetorically – of technology in the delivery of nursing, midwifery and AHP decision-informed care has not led to similar growth in evaluations.

Our planned analysis was SECONDARY|primary; a synthesis of a rich and detailed extensive pool of well-conducted comparative evaluations that might even merit meta-analysis of common systems, followed by smaller-scale qualitative interviews with those with a stake in decision support. On reflection, a secondary|PRIMARY balance may have been more productive. Whilst the pool of experimental/quasi-experimental research has more than tripled since the last comparable review in 2009 it is still very limited and heterogeneous and still unsuited to statistical synthesis and aggregation. Our primary data collection was hampered by the effects of the COVID-19 pandemic on accessing services and users of CDSS in healthcare. We struggled to recruit, arrange and conduct interviews remotely with the numbers of CDSS implementers, service supporters and service-users. But our interviews and virtual meetings suggest that there is still much to be gained from understanding the experiences and perceptions of those who develop, implement, use and receive/take part in CDSS-mediated professional advice/choices. The uncertainty in the quantitative evaluations synthesised, the depth of perceptions and experience in the limited calibration undertaken, and the role that theory – and theory revised in the light of empirical and perceptual description – could play in improving systems and their implementation make the need for well-conducted, theory-informed, primary qualitative research more urgent.

Study/trial registration details

The main study protocol was registered with PROSPERO¹ [number: CRD42019147773] and a protocol for our (sub)review of randomised controlled trials was published in the Cochrane Library.¹⁰²

Implications for decision-makers

At the system level, the passion of advocates for decision support systems as part of 'digital' healthcare is evident in social media and dedicated professional forums. Amongst policy-makers, the investment and intent to use more technology are regularly expressed by ministers and government. In the UK for example, it is a governmental ambition that:

*care professionals should be able to use decision support tools to provide the best care, medicine or device for a patient based on accurate and available data.*¹¹¹

Whilst developers, service leaders and advocates may push for greater tech-based decision support, our synthesis suggests that the decision-maker experience and real-world results may not warrant

such enthusiasm. Governments and health systems investing in CDSS-enabled care should adequately fund well-designed evaluations with sufficient data and methodological support to illuminate the return on investment from CDSS and enable truly informed choices about purchasing or development. There is an urgent need for well-designed health economic evaluations of CDSS to inform system and organisational-level decisions.

Organisations considering implementing or investing in CDSS aimed at nursing, midwifery and AHP work should be aware that there is no evidence on the 'best' systems to purchase or invest in. Claimed effects are likely to be accompanied by considerable uncertainty and an equal amount of effort needs to be put into the implementation and resourcing of the 'work' needed to initiate, embed and sustain CDSS in a varied workforce whose decision-informed work is heavily contextualised and shapes technology use and its interaction with practice and clinical experience. Whilst some of the technical skills in ICT and data science required to instigate or adopt a CDSS might be planned for, it is our experience that skills required to embed CDSS in work effectively are less evident in many services: theory-informed quality improvement and behaviour-change expertise; human factors and other engineering-focused skills; evaluation of complex interventions. Organisations should pay as much attention to building teams with these skills to support CDSS embedding as they might ICT or clinical champions/advocates. Such teams should include an active patient focus and voice to aid adoption. The five FAQs from our patients – if addressed by providers – might also usefully help encourage 'coherence' (shared understanding between potential users/adopter in NPT). Marketing and information aimed at users (both professional and patient) should aim to provide answers tailored to a local service context.

Individual clinicians are often told that the recommendations of CDSS are just that: recommendations. It is tempting to downplay the need for critical appraisal of CDSS recommendations; indeed, prescriptive models of searching for research evidence to apply to patient care decisions have been taken by some to imply that (computerised decision support) 'systems' do not require appraisal – their major advantage.¹¹² The architects of these recommendations are more cautious and are clear that some responsibilities lie on both the system developers and implementers:

The only more compiled source would be a [computerised decision support] system, such as an electronic medical [sic.] record, in which the individual patient's characteristics were automatically linked to the current best evidence that matched their specific circumstances, with caregivers being reminded or notified of key aspects of management. Such computerised decision support systems are currently few and far between, and those in existence often fall short of ensuring that the evidence supporting the system is the best available and is kept up to date.¹¹³

And the CDSS user:

Users of evidence reports at any level of the 5S pyramid need to be aware of the underlying methods of assembly and assure themselves that these methods are sound. At each level, the standards for evidence generation, retrieval, selection, and analysis should be explicit and at the highest evidence standard possible. For example, systems based on guidelines for patient care should be explicit about the source of the guidelines, and the guidelines should be based on systematic reviews of the pertinent evidence to date.¹¹³

Whilst recognising that, for ad hoc individual clinical decisions and clinicians, critical appraisal of a CDSS system's evidence base is unrealistic, we recommend that where a CDSS system is being commissioned or implemented, then those with responsibility for others' use of CDSS-generated advice assure themselves of the quality of the evidence underpinning the system. Ideally guidance should come from the most reliable, trustworthy and clinically useful research evidence. It was clear from our interviews with system implementers that some systems in use in settings are adapted and based on the collective knowledge of 'experts', often from within the hospital, or standardised terminological systems (for example, NANDA) where the effects on clinical outcomes of decisions informed by them are largely unknown.

Research recommendations

Evaluators (as well as system designers) should consider appropriate middle-range implementation theory as a means of negotiating the complex socio-technical relationship between technology and associated work. As with quality improvement¹⁰⁶ and implementation-related behaviours,¹⁰⁵ testing theory-based predictions systematically and building on extant theory in design and evaluation of CDSS will do much to reduce the probability of a similarly 'broad, but shallow' pool of empirical studies facing researchers updating this synthesis in another 10 years.

Second, future syntheses should be made easier and more informative by journals and authors adhering to guidelines for conducting and reporting CDSS when publishing findings. These guidelines for CDSS, in general, already exist; there is no reason to believe that they cannot be applied to nursing, midwifery and AHP-focused CDSS.^{114,115}

Whilst the growth in the number of comparative evaluations in the past decade is welcome, simple head-to-head comparisons rarely help us understand and learn from failure and success of systems. CDSS are a complex healthcare intervention and the development and evaluation of the systems needs to reflect this complexity. Well-established guidance exists for developing and evaluating such interventions and should form an explicit part of studies.¹⁰⁴ This is particularly the case where the comparative evidence is – relatively – well developed (for example, triage decisions or drug dosing), where stronger use of explicit theory could lead to better-designed studies and more insight – especially from modest CDSS effects or 'failure'.

The absence of PPI involvement in primary studies of CDSS should be remedied. In an era in which shared decision-making is an expected norm in many areas of healthcare practice, it is hard to conceive of good reasons why patients should not have a stronger role in informing the design, implementation and evaluation of CDSS. Missing opportunities for better PPI means researchers may unwittingly perpetuate existing inequalities in areas such as technological access or literacy between demographic and professional groups.

Given the well-established potential for CDSS to introduce harm into clinical work, studies should incorporate the possibility of both e-iatrogenesis and harm into their designs. As with any other health technology evaluation, they should measure safety events and unintended consequences in their research. This is particularly important in any economic analysis of the costs and benefits of systems.

Like issues of safety, future research should seek to build the economic evidence surrounding CDSS. As well as the sorts of cost-effectiveness analysis that would allow potential adopters to choose between different systems – studies we acknowledge may be complex to design and conduct for in-house evaluation teams – even the simplest cost-consequence analysis could and should consider the costs of implementing the systems. These are often missing from studies and – as with quality improvement in healthcare generally – have the effect of making systems seem more attractive than they might be if the true costs were known.¹¹⁶

Conclusion

Our review of available evidence comparing CDSS support to unsupported (by CDSS) decision-related performance and outcomes in nurses and – to a far lesser extent – midwives and AHPs synthesises 35 studies and reveals a contradictory picture. Studies reported that around half the process measures improved and half were worse or no different. CDSS-attributable outcomes fared little better, with only ≈40% of outcomes reported actually improving with CDSS use and, in one study, producing fewer delivery complications less efficiently than not using the system. Characteristics previously believed to improve general CDSS effectiveness were not associated with improved processes or outcomes. The

within-study variation in take-up and wide confidence intervals around effects suggest two things. First, 13 years since the last substantive non-medical review, uncertain effects are still evident. Second, simply adding more studies into the heterogeneous bundle of extant evaluations, without systematically testing and refining relevant mid-range theories, will not lead to the step change in replicable implementation that policy-makers and advocates of digital technology in health desire so urgently.

This is the first systematic review that set out to control for the same systems used in different professional and organisational contexts. The limited empirical and theoretical treatment of the complex socio-technical context in which CDSS operate means we are unable to offer definitive conclusions about which systems best improve processes and outcomes in NMAHP work. The synthesis does allow us to add to the growing realisation that CDSS planners, implementers and evaluators should use well-developed and rigorously tested mid-range theory that both already exists and is designed to produce technologies better able to work within complex socio-technical systems. Doing this *before* systems are designed and evaluated, rather than having to use theory to scrutinise processes and effects, would be more efficient. Justifying greater investment in implementing CDSS, unaccompanied by well-designed, theoretically informed evaluation, is made more challenging by the absence of economic comparisons of the costs and benefits of different systems aimed at NMAHPs. Studies that focus on real-world implementation of CDSS should – by default – incorporate some form of health economic evaluation.

CDSS are often sold on the promise that they can support and release the potential of NMAHPs' decisions in multidisciplinary healthcare to increase quality efficiently. Our synthesis suggests there is still some way to go before this is a routine reality for both clinicians and patients.

Acknowledgements

We would like to thank the CDSS experts and patients that gave their time to help us contextualise our results at a point in our history when providing time for others is a generous gift.

Contributions of authors

Carl Thompson (<https://orcid.org/0000-0002-9369-1204>) drafted this synthesis and was responsible for the initial research idea and bid.

Teumzghi Mebrahtu (<https://orcid.org/0000-0003-4821-2304>) and **Carl Thompson** (<https://orcid.org/0000-0002-9369-1204>) were responsible for the review design, extraction and analysis; all team members at some point helped screen titles and abstracts, informed our analysis, and helped draft work and articles.

Sarah Skyrme (<https://orcid.org/0000-0002-6173-0117>) and **Alison Ledward** carried out the virtual meeting of PPG members and production of patient FAQs on CDSS.

Sarah Skyrme (<https://orcid.org/0000-0002-6173-0117>) also interviewed our system experts.

Ethical approval for the qualitative components of the project was sought and granted by the University of Leeds Faculty of Medicine and Health Ethics Committee HREC 20-002 - Effects of computerised clinical decision support systems (CDSS) on nursing and Allied Health Professional performance and patient outcomes: A systematic review and user contextualisation.

Data-sharing statement

All data requests should be submitted to the corresponding author for consideration. Access to available anonymised data may be granted following review.

Ethics statement

Ethical approval for the study was granted by the School of Healthcare Research Ethics Committee, University of Leeds, on 20 October 2020: approval reference HREC 20-002.

This article

The contractual start date for this research was in October 2019. This article began editorial review in October 2021 and was accepted for publication in July 2022. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The *Health and Social Care Delivery Research* editors and publisher have tried to ensure the accuracy of the authors' article and would like to thank the reviewers for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this article.

Disclaimer

The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

This manuscript reports on one component of the research award Effects of computerised clinical decision support systems (CDSS) on nursing and Allied Health Professional performance and patient outcomes: A systematic review and user contextualisation. For more information about this research please view the award page [<https://www.fundingawards.nihr.ac.uk/award/NIHR127926>]

References

1. Thompson C, Randell R, Keenan AM, *et al.* Effects of computerised clinical decision support systems (CDSS) on nursing and allied health professional performance and patient outcomes: a systematic review and user contextualisation. PROSPERO 2019 [cited background]. URL: www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42019147773.
2. Covell DG, Uman GC, Manning PR. Information needs in office practice: are they being met? [published Online First: 1985/10/01]. *Ann Intern Med* 1985;**103**(4):596–9. <https://doi.org/10.7326/0003-4819-103-4-596>.
3. Thompson C, Cullum N, McCaughan D, *et al.* Nurses, information use, and clinical decision making – the real world potential for evidence-based decisions in nursing [published Online First: 2004/07/16]. *Evid Based Nurs* 2004;**7**(3):68–72. <https://doi.org/10.1136/ebn.7.3.68>.
4. Garg AX, Adhikari NKJ, McDonald H, *et al.* Effects of computerized clinical decision support systems on practitioner performance and patient outcomes a systematic review. *JAMA* 2005;**293**(10):1223–38. <https://doi.org/10.1001/jama.293.10.1223>.
5. Cooper RJ, Bissell P, Ward P, *et al.* Further challenges to medical dominance? The case of nurse and pharmacist supplementary prescribing. *Health* 2012;**16**(2):115–33. <https://doi.org/10.1177/1363459310364159>.
6. Noordegraaf M. Hybrid professionalism and beyond: (New) Forms of public professionalism in changing organizational and societal contexts. *J Prof Organ* 2015;**2**(2):187–206. <https://doi.org/10.1093/jpo/jov002>.
7. Hallgren KA, Bauer AM, Atkins DC. Digital technology and clinical decision making in depression treatment: current findings and future opportunities [published Online First: 2017/04/28]. *Depress Anxiety* 2017;**34**(6):494–501. <https://doi.org/10.1002/da.22640>.
8. Hughes DJ, Allen D. *Nursing and the Division of Labour in Healthcare*. Basingstoke: Palgrave; 2002.
9. Hammond KR, Kelly KJ, Schneider RJ, *et al.* Clinical inference in nursing: revising judgments. *Nurs Res* 1967;**16**(1):38–45.
10. Buckingham CD, Adams A. Classifying clinical decision making: a unifying approach [published Online First: 2000/11/30]. *J Adv Nurs* 2000;**32**(4):981–9.
11. Standing M. Clinical judgement and decision-making in nursing - nine modes of practice in a revised cognitive continuum [published Online First: 2008/03/21]. *J Adv Nurs* 2008;**62**(1):124–34. <https://doi.org/10.1111/j.1365-2648.2007.04583.x>.
12. Elstein AS, Schwartz A. Clinical problem solving and diagnostic decision making: selective review of the cognitive literature [published Online First: 2002/03/23]. *BMJ* 2002;**324**(7339):729–32. <https://doi.org/10.1136/bmj.324.7339.729>.
13. Osheroff JA, Teich JM, Middleton B, *et al.* A roadmap for national action on clinical decision support [published Online First: 2007/01/11]. *J Am Med Inform Assoc* 2007;**14**(2):141–5. <https://doi.org/10.1197/jamia.M2334>.
14. Sim I, Gorman P, Greenes RA, *et al.* Clinical decision support systems for the practice of evidence-based medicine. *J Am Med Inform Assoc* 2001;**8**(6):527–34. <https://doi.org/10.1136/jamia.2001.0080527>.
15. Sutton RT, Pincock D, Baumgart DC, *et al.* An overview of clinical decision support systems: benefits, risks, and strategies for success. *NPJ Digit Med* 2020;**3**(1):17. <https://doi.org/10.1038/s41746-020-0221-y>.

16. Lytle KS, Short NM, Richesson RL, *et al.* Clinical decision support for nurses: a fall risk and prevention example [published Online First: 2015/11/17]. *Comput Inform Nurs* 2015;**33**(12):530–7; quiz E1. <https://doi.org/10.1097/CIN.0000000000000192>
17. Khong PC, Hoi SY, Holroyd E, *et al.* Nurses' clinical decision making on adopting a wound clinical decision support system [published Online First: 2015/06/13]. *Comput Inform Nurs* 2015;**33**(7):295–305. <https://doi.org/10.1097/CIN.0000000000000164>.
18. Gross DP, Armijo-Olivo S, Shaw WS, *et al.* Clinical decision support tools for selecting interventions for patients with disabling musculoskeletal disorders: a scoping review [published Online First: 2015/12/17]. *J Occup Rehabil* 2016;**26**(3):286–318. <https://doi.org/10.1007/s10926-015-9614-1>.
19. Martin Ruiz ML, Valero Duboy MA, Torcal Loriente C, *et al.* Evaluating a web-based clinical decision support system for language disorders screening in a nursery school [published Online First: 2014/05/30]. *J Med Internet Res* 2014;**16**(5):e139. <https://doi.org/10.2196/jmir.3263>
20. Mahabir C, Kuperman G, Lantigua R, *et al.* Improving depression screening using CDS and patient access to information using PHRs. *Comput Inform Nurs* 2014;**32**(8):364–65. <https://doi.org/10.1097/01.NCN.0000453183.49477.ee>.
21. Berner ES. *Clinical Decision Support Systems: Theory and Practice*. 2nd ed. New York, NY: Springer; 2007.
22. Cresswell K, Majeed A, Bates DW, *et al.* Computerised decision support systems for healthcare professionals: an interpretative review [published Online First: 2012/01/01]. *Inform Prim Care* 2012;**20**(2):115–28. <https://doi.org/10.14236/jhi.v20i2.32>.
23. Rahbari NN, Reissfelder C, Koch M, *et al.* The predictive value of postoperative clinical risk scores for outcome after hepatic resection: a validation analysis in 807 patients [published Online First: 2011/06/16]. *Ann Surg Oncol* 2011;**18**(13):3640–9. <https://doi.org/10.1245/s10434-011-1829-6>.
24. Cook DA, Pencille LJ, Dupras DM, *et al.* Practice variation and practice guidelines: attitudes of generalist and specialist physicians, nurse practitioners, and physician assistants. *PLOS ONE* 2018;**13**(1):e0191943. <https://doi.org/10.1371/journal.pone.0191943>.
25. Stride SL, Hundley VA, Way S, *et al.* Identifying the factors that influence midwives' perineal practice at the time of birth in the United Kingdom. *Midwifery* 2021;**102**:103077. <https://doi.org/10.1016/j.midw.2021.103077>.
26. Cheyne H, Dowding DW, Hundley V. Making the diagnosis of labour: midwives' diagnostic judgement and management decisions. *J Adv Nurs* 2006;**53**(6):625–35. <https://doi.org/10.1111/j.1365-2648.2006.03769.x>.
27. Royal College of Nursing. *Improving Digital Literacy 2020* [cited background]. URL: www.rcn.org.uk/professional-development/publications/pub-006129.
28. Bright TJ, Wong A, Dhurjati R, *et al.* Effect of clinical decision-support systems: a systematic review. *Annals of Internal Medicine* 2012;**157**(1):29–43. <https://doi.org/10.7326/0003-4819-157-1-201207030-00450> %.
29. Moja L, Kwag KH, Lytras T, *et al.* Effectiveness of computerized decision support systems linked to electronic health records: a systematic review and meta-analysis [published Online First: 2014/10/17]. *Am J Public Health* 2014;**104**(12):e12–22. <https://doi.org/10.2105/AJPH.2014.302164>.
30. Randell R, Mitchell N, Dowding D, *et al.* Effects of computerized decision support systems on nursing performance and patient outcomes: a systematic review. *J Health Serv Res Policy* 2007;**12**(4):242–51. <https://doi.org/10.1258/135581907782101543>.

31. Gillaizeau F, Chan E, Trinquart L, *et al.* Computerized advice on drug dosage to improve prescribing practice. *Cochrane Database Syst Rev* **2013**(11):CD002894.
32. Tan K, Dear PR, Newell SJ. Clinical decision support systems for neonatal care. *Cochrane Database Syst Rev* **2005**(2):CD004211.
33. Coiera E, Westbrook J, Wyatt J. The safety and quality of decision support systems [published Online First: 2006/10/20]. *Yearb Med Inform* 2006:20–5.
34. Dunn Lopez K, Gephart SM, Raszewski R, *et al.* Integrative review of clinical decision support for registered nurses in acute care settings. *J Am Med Inform Assoc* 2016;**24**(2):441–50. <https://doi.org/10.1093/jamia/ocw084>.
35. Dowding D, Mitchell N, Randell R, *et al.* Nurses' use of computerised clinical decision support systems: a case site analysis. *J Clin Nurs* 2009;**18**(8):1159–67. <https://doi.org/10.1111/j.1365-2702.2008.02607.x>.
36. Randell R, Mitchell N, Dowding D, *et al.* Effects of computerized decision support systems on nursing performance and patient outcomes: a systematic review [published Online First: 2007/10/11]. *J Health Serv Res Policy* 2007;**12**(4):242–9. <https://doi.org/10.1258/135581907782101543>.
37. Porter A, Dale J, Foster T, *et al.* Implementation and use of computerised clinical decision support (CCDS) in emergency pre-hospital care: a qualitative study of paramedic views and experience using Strong Structuration Theory. *Implement Sci* 2018;**13**(1):91. <https://doi.org/10.1186/s13012-018-0786-x>.
38. Weiner JP, Kfuri T, Chan K, *et al.* 'e-Iatrogenesis': the most critical unintended consequence of CPOE and other HIT [published Online First: 2007/03/03]. *J Am Med Inform Assoc* 2007;**14**(3):387–8; discussion 89. <https://doi.org/10.1197/jamia.M2338>.
39. Langhorne P, Ramachandra S. Organised inpatient (stroke unit) care for stroke: network meta-analysis. *Cochrane Database Syst Rev* 2020(4). <https://doi.org/10.1002/14651858.CD000197.pub4>.
40. Sintchenko V, Coiera E. Decision complexity affects the extent and type of decision support use. *AMIA Annu Symp* 2006;724–8.
41. Roshanov PS, Fernandes N, Wilczynski JM, *et al.* Features of effective computerised clinical decision support systems: meta-regression of 162 randomised trials. *BMJ* 2013;**346**.
42. Van de Velde S, Heselmans A, Delvaux N, *et al.* A systematic review of trials evaluating success factors of interventions with computerised clinical decision support. *Implement Sci* 2018;**13**(1):114. <https://doi.org/10.1186/s13012-018-0790-1>.
43. May C, Finch T, Mair F, *et al.* Understanding the implementation of complex interventions in health care: the normalization process model [published Online First: 2007/09/21]. *BMC Health Serv Res* 2007;**7**:148. <https://doi.org/10.1186/1472-6963-7-148>.
44. Greenhalgh T, Wherton J, Papoutsis C, *et al.* Beyond adoption: a new framework for theorizing and evaluating nonadoption, abandonment, and challenges to the scale-up, spread, and sustainability of health and care technologies [published Online First: 2017/11/03]. *J Med Internet Res* 2017;**19**(11):e367. <https://doi.org/10.2196/jmir.8775>.
45. Cresswell KM, Worth A, Sheikh A. Actor-Network Theory and its role in understanding the implementation of information technology developments in healthcare. *BMC Medical Inform Decis Mak* 2010;**10**(1):67. <https://doi.org/10.1186/1472-6947-10-67>.
46. May CR, Cummings A, Girling M, *et al.* Using Normalization Process Theory in feasibility studies and process evaluations of complex healthcare interventions: a systematic review [published Online First: 2018/06/09]. *Implement Sci* 2018;**13**(1):80. <https://doi.org/10.1186/s13012-018-0758-1>.

47. Finch TL, Rapley T, Girling M, et al. Improving the normalization of complex interventions: measure development based on normalization process theory (NoMAD): study protocol [published Online First: 2013/04/13]. *Implement Sci* 2013;**8**:43. <https://doi.org/10.1186/1748-5908-8-43>.
48. Pope C, Halford S, Turnbull J, et al. Using computer decision support systems in NHS emergency and urgent care: ethnographic study using normalisation process theory [published Online First: 2013/03/26]. *BMC Health Serv Res* 2013;**13**:111. <https://doi.org/10.1186/1472-6963-13-111>.
49. Kawamoto K, Houlihan CA, Balas EA, et al. Improving clinical practice using clinical decision support systems: a systematic review of trials to identify features critical to success. *BMJ* 2005;**330**(7494):765. <https://doi.org/10.1136/bmj.38398.500764.8F>.
50. May CR, Finch T, Ballini L, et al. Evaluating complex interventions and health technologies using normalization process theory: development of a simplified approach and web-enabled toolkit [published Online First: 2011/10/04]. *BMC Health Serv Res* 2011;**11**:245. <https://doi.org/10.1186/1472-6963-11-245>.
51. Cochrane Effective Practice Organisation of Care (EPOC). EPOC Resources for review authors: What study designs can be considered for inclusion in an EPOC review and what should that be called? 2017 [cited background]. URL: <http://epoc.cochrane.org/resources/epoc-resources-review-authors> (accessed 15 April 2020).
52. Cochrane Effective Practice and Organisation of Care (EPOC). Data collection form. *EPOC resources for review authors* 2017 [cited background]. URL: <https://epoc.cochrane.org/resources/epoc-resources-review-authors> (accessed 30 March 2020).
53. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019)*. Cochrane; 2019.
54. Cochrane Effective Practice and Organisation of Care (EPOC). Suggested risk of bias criteria for EPOC reviews. *EPOC resources for review authors* 2017 [cited background]. URL: <https://epoc.cochrane.org/resources/epoc-resources-review-authors>.
55. Heinze G, Schemper M. A solution to the problem of separation in logistic regression. *Stat Med* 2002;**21**(16):2409–19. <https://doi.org/10.1002/sim.1047>.
56. Lipscomb M. Abductive reasoning and qualitative research [published Online First: 2012/09/07]. *Nurs Philos* 2012;**13**(4):244–56. <https://doi.org/10.1111/j.1466-769X.2011.00532.x>.
57. Roberts K, Dowell A, Nie JB. Attempting rigour and replicability in thematic analysis of qualitative research data; a case study of codebook development [published Online First: 2019/03/30]. *BMC Med Res Methodol* 2019;**19**(1):66. <https://doi.org/10.1186/s12874-019-0707-y>.
58. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;**3**(2):77–101. <https://doi.org/10.1191/1478088706qp063oa>.
59. Miles MB, Huberman AM. *Qualitative Data Analysis: An Expanded Sourcebook*. 2nd edn. Thousand Oaks, CA, London: Sage; 1994.
60. Ryan GW, Bernard HR. Techniques to identify themes. *Field Methods* 2003;**15**(1):85–109. <https://doi.org/10.1177/1525822X02239569>.
61. Bennett P, Hardiker N. A Quantitative study investigating the effects of computerised clinical decision support in the emergency department. *Stud Health Technol Inform* 2016;**225**:53–7.
62. Snooks HA, Carter B, Dale J, et al. Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLOS ONE* 2014;**9**(9):e106436. <https://doi.org/10.1371/journal.pone.0106436>.
63. Wells B. *Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice*. Swansea University; 2013.

64. Lattimer V, George S, Thompson F, *et al.* Safety and effectiveness of nurse telephone consultation in out of hours primary care: randomised controlled trial. *BMJ* 1998;**317**(7165):1054–9.
65. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;**320**(7241):1053–7.
66. Cortez S. *Measuring active Clinical Decision Support Influence on Nursing Research Utilization*. University of Phoenix; 2014.
67. Fitzmaurice DA, Hobbs FD, Murray ET, *et al.* Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Arch Intern Med* 2000;**160**(15):2343–8.
68. Kroth PJ, Dexter PR, Overhage JM, *et al.* A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *AMIA Annu Symp* 2006:444–8.
69. Rood E, Bosman RJ, van der Spoel JI, *et al.* Use of a computerized guideline for glucose regulation in the intensive care unit improved both guideline adherence and glucose regulation. *J Am Med Inform Assoc* 2005;**12**(2):172–80. <https://doi.org/10.1197/jamia.M1598>.
70. Sassen B, Kok G, Schepers J, *et al.* Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *J Med Internet Res* 2014;**16**(10):e211.
71. Cleveringa FGW, Gorter KJ, Van Donk MD, *et al.* Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;**31**(12):2273–5.
72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;**33**(2):258–63.
73. Geurts D, de Vos-Kerkhof E, Polinder S, *et al.* Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *Eur J Pediatr* 2017;**176**(2):173–81.
74. Lee NJ, Chen ES, Currie LM, *et al.* The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. <https://doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, *et al.* Comparison of three protocols for tight glycemic control in cardiac surgery patients. *Diabetes Care* 2009;**32**(5):757. <https://doi.org/10.2337/dc08-1851>.
76. Canbolat O, Kapucu S, Kilickaya O. Comparison of routine and computer-guided glucose management for glycemic control in critically ill patients. *Crit Care Nurse* 2019;**39**(4):20–7. <https://doi.org/10.4037/ccn2019431>.
77. Cavalcanti AB, Silva E, Pereira AJ, *et al.* A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *J Crit Care* 2009;**24**(3):371–8. <https://doi.org/10.1016/j.jcrc.2009.05.005>.
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;**21**(2):106–15. <https://doi.org/10.4037/ajcc2012956>.
79. Hovorka R, Kremen J, Blaha J, *et al.* Blood glucose control by a model predictive control algorithm with variable sampling rate versus a routine glucose management protocol in cardiac surgery patients: a randomized controlled trial. *J Clin Endocrinol Metab* 2007;**92**(8):2960–4. <https://doi.org/10.1210/jc.2007-0434>.

80. Mann EA, Jones JA, Wolf SE, *et al.* Computer decision support software safely improves glycemic control in the burn intensive care unit: a randomized controlled clinical study. *J Burn Care Res* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, *et al.* Multicentric, randomized, controlled trial to evaluate blood glucose control by the model predictive control algorithm versus routine glucose management protocols in intensive care unit patients. *Diabetes Care* 2006;**29**(2):271. <https://doi.org/10.2337/diacare.29.02.06.dc05-1689>.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clin Lab Haematol* 1997;**19**(3):203–7.
83. Byrne CM. *Impact of Prospective Computerized Clinical Decision Support Information and Targeted Assistance on Nursing Home Resident Outcomes*. State University of New York at Albany; 2005.
84. Duclos A, Touzet S, Restier L, *et al.* Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *Eur J Clin Nutr* 2015;**69**(7):769–75. <https://doi.org/10.1038/ejcn.2014.288>.
85. Forberg U, Unbeck M, Wallin L, *et al.* Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implement Sci* 2016;**11**. <https://doi.org/10.1186/s13012-016-0375-9>.
86. Dalaba MA, Akweongo P, Aborigo RA, *et al.* Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLOS ONE* 2015;**10**(5):e0125920. <https://doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, *et al.* Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *J Am Med Inform Assoc* 2008;**15**(1):107–13. <https://doi.org/10.1197/jamia.M2164>.
88. McDonald MV, Feldman PH, Barron-Vaya Y, *et al.* Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *J Eval Clin Pract* 2017;**22**(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, *et al.* A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;**50**(4):475–86. <https://doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis [published Online First: 2011/12/17]. *J Am Med Inform Assoc* 2012;**19**(4):615–20. <https://doi.org/10.1136/amiajnl-2011-000504>.
91. Dykes PC, Carroll DL, Hurley A, *et al.* Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;**304**(17):1912–8.
92. Dykes PC, Burns Z, Adelman J, *et al.* Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA Netw Open* 2020;**3**(11):e2025889–e89.
93. Fossum M, Alexander GL, Ehnfors M, *et al.* Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;**80**(9):607–17. <https://doi.org/10.1016/j.ijmedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, *et al.* A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *J Adv Nurs* 2019;**75**(11):3058–67.

95. Paulsen MM, Paur I, Gjestland J, *et al.* Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: a randomized controlled trial. *Clin Nutr* 2020;**39**(12):3607–17.
96. Dykes PC, Burns Z, Adelman J, *et al.* Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA Netw Open* 2020;**3**(11):e2025889–e89. <https://doi.org/10.1001/jamanetworkopen.2020.25889>.
97. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;**320**(7241):1053–7.
98. Abbott AD. *The System of Professions: An Essay on the Division of Expert Labor*. Chicago, IL: University of Chicago Press; 1988.
99. Fishbein M, Ajzen I. *Predicting and Changing Behavior: The Reasoned Action Approach*. New York, NY; Hove: Psychology Press; 2009.
100. Coiera E, Ash J, Berg M. The unintended consequences of health information technology revisited [published Online First: 2016/11/11]. *Yearb Med Inform* **2016**(1):163–9. <https://doi.org/10.15265/IY-2016-014>.
101. Sittig DF. A socio-technical model of health information technology-related e-iatrogenesis [published Online First: 2008/11/13]. *AMIA Annu Symp Proc* 2008:1209–10.
102. Mebrahtu TF, Bloor K, Ledward A, *et al.* Effects of computerised clinical decision support systems (CDSS) on nursing and allied health professional performance and patient outcomes. *Cochrane Database Syst Rev* **2021**(3). <https://doi.org/10.1002/14651858.CD014699>.
103. Greenhalgh T, Hinder S, Stramer K, *et al.* Adoption, non-adoption, and abandonment of a personal electronic health record: case study of HealthSpace [published Online First: 2010/11/18]. *BMJ* 2010;**341**:c5814. <https://doi.org/10.1136/bmj.c5814>.
104. O'Cathain A, Croot L, Duncan E, *et al.* Guidance on how to develop complex interventions to improve health and healthcare. *BMJ Open* 2019;**9**(8):e029954. <https://doi.org/10.1136/bmjopen-2019-029954>.
105. Atkins L, Francis J, Islam R, *et al.* A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems [published Online First: 2017/06/24]. *Implement Sci* 2017;**12**(1):77. <https://doi.org/10.1186/s13012-017-0605-9>.
106. Grol R, Baker R, Moss F. Quality improvement research: understanding the science of change in health care [published Online First: 2002/11/27]. *Qual Saf Health Care* 2002;**11**(2):110–1. <https://doi.org/10.1136/qhc.11.2.110>.
107. Ivers NM, Sales A, Colquhoun H, *et al.* No more 'business as usual' with audit and feedback interventions: towards an agenda for a reinvigorated intervention [published Online First: 2014/01/21]. *Implement Sci* 2014;**9**:14. <https://doi.org/10.1186/1748-5908-9-14>.
108. Dune R. *Evaluating Evaluations of Clinical Decision Support Systems: Case Studies From NHS Clinical Settings*. Open University; 2016.
109. Varghese J, Kleine M, Gessner SI, *et al.* Effects of computerized decision support system implementations on patient outcomes in inpatient care: a systematic review [published Online First: 2017/10/17]. *J Am Med Inform Assoc* 2018;**25**(5):593–602. <https://doi.org/10.1093/jamia/ocx100>.
110. Hutchings R. *The Impact of Covid-19 on The Use of Digital Technology in the NHS*. London: Nuffield Trust; 2020. URL: www.nuffieldtrust.org.uk/files/2020-08/the-impact-of-covid-19-on-the-use-of-digital-technology-in-the-nhs-web-2.pdf (accessed 31 May 2022).
111. Department of Health and Social Care. The future of healthcare: our vision for digital, data and technology in health and care London: gov.uk; 2018 [Policy Paper]. URL: www.gov.uk/government/publications/the-future-of-healthcare-our-vision-for-digital-data-and-technology-in-health-and-care/

- [the-future-of-healthcare-our-vision-for-digital-data-and-technology-in-health-and-care](#) (accessed 1 July 2021).
112. Dicenso A, Bayley L, Haynes RB. Accessing pre-appraised evidence: fine-tuning the 5S model into a 6S model [published Online First: 2009/09/26]. *Evid Based Nurs* 2009;**12**(4):99–101. <https://doi.org/10.1136/ebn.12.4.99-b>.
 113. Haynes RB. Of studies, syntheses, synopses, summaries, and systems: the '5S' evolution of information services for evidence-based healthcare decisions [published Online First: 2007/01/11]. *Evid Based Med* 2006;**11**(6):162–4. <https://doi.org/10.1136/ebm.11.6.162-a>.
 114. Kawamoto K, McDonald CJ. Designing, conducting, and reporting clinical decision support studies: recommendations and call to action. *Ann Intern Med* 2020;**172**(11_Supplement):S101–9. <https://doi.org/10.7326/M19-0875>.
 115. National Health Services (NHS). Digital Health: NHSX launches assessment criteria for digital health tools 2020 [cited discussion. URL: www.digitalhealth.net/2020/10/nhsx-launches-assessment-criteria-for-digital-health-tools/] (accessed 29 October 2020).
 116. Thompson C, Pulleyblank R, Parrott S, *et al*. The cost-effectiveness of quality improvement projects: a conceptual framework, checklist and online tool for considering the costs and consequences of implementation-based quality improvement [published Online First: 2015/07/24]. *J Eval Clin Pract* 2016;**22**(1):26–30. <https://doi.org/10.1111/jep.12421>.

Appendix 1 Example Ovid MEDLINE(R) 1946 to 12 February 2021 search strategy

1. exp Decision Making/ (207895)
2. decision support techniques/ (20911)
3. (decision* adj2 making).ti,ab,kf. (159754)
4. (decision* adj2 support*).ti,ab,kf. (24230)
5. (decision* adj2 aid*).ti,ab,kf. (6501)
6. or/1-5 (354546)
7. exp Computers/ (79322)
8. exp information systems/ (238259)
9. exp Informatics/ (537355)
10. Internet/ (74916)
11. Software/ (112580)
12. Cell Phone/ (8821)
13. Mobile Applications/ (6962)
14. exp Telemedicine/ (32559)
15. Medical Records Systems, Computerized/ (19076)
16. exp Electronic Health Records/ (21793)
17. computer*.ti,ab,kf. (313610)
18. electronic*.ti,ab,kf. (291368)
19. (internet or web or online or on-line).ti,ab,kf. (310071)
20. (software or computer program*).ti,ab,kf. (193359)
21. (automate* or automation).ti,ab,kf. (136436)
22. (pda or pdas).ti,ab,kf. (13229)
23. personal digital assistant*.ti,ab,kf. (1012)
24. (app or apps).ti,ab,kf. (31717)
25. (application* adj2 mobile*).ti,ab,kf. (4834)
26. (iPad* or iPhone* or smartphone* or smart phone* or smart device* or mobile phone or android phone* or cellphone* or cell phone*).ti,ab,kf. (26450)
27. (tablet adj2 (pc or device* or comput*)).ti,ab,kf. (1603)
28. ((hand held or handheld) adj2 (pc or device* or comput*)).ti,ab,kf. (2669)
29. (telehealth or telecare or telemedicine or ehealth or mhealth).ti,ab,kf. (29130)
30. or/7-29 (1674343)
31. 6 and 30 (66042)
32. exp Decision Making, Computer-Assisted/ (149528)
33. Decision Support Systems, Clinical/ (8302)
34. (computer assisted adj2 (decision* or diagnos* or therap* or support or treatment? or management)).ti,ab,kf. (1545)
35. (computer aided adj2 (decision* or diagnos* or therap* or support or treatment? or management)).ti,ab,kf. (3921)
36. (decision adj2 support adj2 (system* or tool*)).ti,ab,kf. (9917)
37. (decision making adj2 (system* or tool*)).ti,ab,kf. (2560)
38. Expert Systems/ (3420)
39. (expert adj2 system*).ti,ab,kf. (3613)
40. Reminder Systems/ (3568)
41. ((computer* or electronic* or CDSS) adj2 (reminder* or alert*)).ti,ab,kf. (1210)
42. ((medication or medicine or treatment or therapy) adj2 (reminder* or alert*)).ti,ab,kf. (857)
43. reminder system*.ti,ab,kf. (875)
44. Medical Order Entry Systems/ (2303)

45. ((computer* or electronic*) adj2 order entry).ti,ab,kf. (1874)
46. (computer adj2 decision support*).ti,ab. (412)
47. CPOE.ti,ab,kf. (1139)
48. or/32-47 (177952)
49. 31 or 48 [all computerised clinical decision support systems terms] (228840)
50. Allied Health Personnel/ (11925)
51. Allied Health Occupations/ (587)
52. Physical Therapist Assistants/ (16)
53. Physical Therapy Specialty/ (2889)
54. Speech-Language Pathology/ (3172)
55. Occupational Therapy/ (13482)
56. Nutritionists/ (1290)
57. dietetics/ (7837)
58. Anesthesiologists/ (1163)
59. podiatry/ (2273)
60. exp Osteopaths/ (321)
61. osteopathic physicians/ (321)
62. anesthesiologist*.ti,ab,kf. (22810)
63. podiatrist*.ti,ab,kf. (910)
64. prosthetist*.ti,ab,kf. (397)
65. chiropodist*.ti,ab,kf. (132)
66. orthoptist*.ti,ab,kf. (319)
67. orthotist*.ti,ab,kf. (220)
68. osteopath*.ti,ab,kf. (5983)
69. radiographer*.ti,ab,kf. (1803)
70. art therapist*.ti,ab,kf. (89)
71. drama therapist*.ti,ab,kf. (3)
72. music therapist*.ti,ab,kf. (368)
73. (allied adj2 health adj2 (profession* or worker* or personnel or occupation* or staff)).ti,ab,kf. (3421)
74. ((physical or occupational or language or speech or physio*) adj2 therap*).ti,ab,kf. (50227)
75. physiotherapist*.ti,ab,kf. (8544)
76. dietetic*.ti,ab,kf. (9828)
77. dietitian*.ti,ab,kf. (6580)
78. nutritionist*.ti,ab,kf. (3020)
79. Patient care team/ (66483)
80. ((multidisciplinary or multi-disciplinary or multiprofessional or multi-professional or interdisciplinary or interprofessional) adj2 team*).ti,ab,kf. (32126)
81. Emergency Medical Technicians/ (5756)
82. Emergency Medical Services/ (43736)
83. Ambulances/ (6210)
84. Air Ambulances/ (2874)
85. paramedic*.ti,ab,kf. (8537)
86. HEMS.ti,ab,kf. (767)
87. ems.ti,ab,kf. (13017)
88. emt.ti,ab,kf. (25232)
89. prehospital.ti,ab,kf. (13136)
90. pre-hospital.ti,ab,kf. (4836)
91. first responder*.ti,ab,kf. (2449)
92. emergency medical technician*.ti,ab,kf. (1168)
93. emergency services.ti,ab,kf. (4115)
94. ambulance*.ti,ab,kf. (11269)
95. field triage.ti,ab,kf. (275)

96. out-of-hospital.ti,ab,kf. (11317)
97. (nurse or nurses or nursing).ti,ab,kf. (462330)
98. exp nurses/ (89638)
99. exp nursing staff/ (67063)
100. Midwifery/ (19460)
101. (midwif* or midwiv*).ti,ab,kf. (25895)
102. or/50–101 [allied health professionals or nurses or midwives] (836031)
103. 49 and 102 [all CDSS and allied health professionals or nurses or midwives] (9549) (see *Appendix 2, Table 11*).

Appendix 2 Additional information on studies and interventions

TABLE 11 Additional information on studies and interventions

Author and year	Type of report	Funding source	Interventions and their details	CDSS function category
Beeckman <i>et al.</i> 2013 ⁸⁹	Journal article	None	<ul style="list-style-type: none"> • A standalone computer-based CDSS (Pre-vPlan-a six-step clinical practice to reduce pressure ulcers using CDSS): <ul style="list-style-type: none"> Step-1: Analysis of current practice, target group, and context Step-2: Match of research findings and/or existing guidelines to practice Step-3: description of the specific change outcome Step-4: Selection/development of the implementation strategies Step-5: Development and execution of the implementation process Step-6: Continuous evaluation and adaptation of the implementation • A standard protocol (a hard copy with no implementation strategy) of reducing pressure ulcers 	Disease prevention
Bennet <i>et al.</i> 2016 ⁶¹	Journal article	Not stated	<ul style="list-style-type: none"> • A standalone computer-based Triage CDSS [intervention period]: the CDSS was developed in-house by engineers and ED clinician to be used in triage in emergency department • Triage without CDSS [pre-intervention period] 	Triaging
Blaaha <i>et al.</i> 2009 ⁷⁵	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a predictive model algorithm that, automates blood glucose control in critically ill patients, by calculating the current insulin of each individual patient and generating advice on the new insulin infusion rate. • Control-1 (paper based-Matias protocol): continuous insulin infusion combined with iv insulin boluses (RMP) to maintain euglycemia (target range 4.4–6.1 mmol/liter) • Control-2 (paper based-Bath protocol): initial insulin infusion rate is dependent on the initial blood glucose. Subsequent changes to the infusion rate are made after comparing the latest blood glucose both to the target range and also to the previous blood glucose value 	Drug dosing
Byrne 2005 ⁸³	PhD Thesis	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: computerized information to alert nursing and other staff to the resident-specific risk factors • Usual care (CDSS non-use) 	Disease prevention reminders
Canbolat <i>et al.</i> 2019 ⁷⁶	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer based CDSS: automated blood glucose monitoring system using newly developed glycaemic control software. • Usual care: ICU nurses measured blood glucose levels and reported them to ICU physicians, who made dosing and treatment decisions based on their own knowledge and assessment. 	Drug dosing

continued

TABLE 11 Additional information on studies and interventions (continued)

Author and year	Type of report	Funding source	Interventions and their details	CDSS function category
Cavalcanti <i>et al.</i> 2009 ⁷⁷	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer/handheld-based CDSS: a computer program running on an ICU desktop or a handheld (electronic supplemental material) used for adjusting insulin doses. • Control-1 (Leuven protocol): continuous intravenous insulin infusion with adjustments according to a protocol developed and used by Van den Berghe <i>et al.</i> • Control-2 (conventional treatment): intermittent subcutaneous insulin administration according to a sliding scale. 	Drug dosing
Cleveringa <i>et al.</i> 2008 ⁷¹	Journal article	Private sector-commercial	<ul style="list-style-type: none"> • A standalone computer-based CDSS: an intervention with the following components:1) diabetes consultation hour run by a practice nurse,2) a CDSS that contained a diagnostic and treatment algorithm based on the Dutch type2 diabetes guidelines and provided patient-specific treatment advice,3) a recall system, and 4) feedback every 3 months regarding the percentage of patients meeting the treatment target. The primary care physicians (PCPs) were advised that they should prescribe new medication and refer patients if necessary. • Usual care: diabetes care provided by the Primary care physicians (PCP) or by a practice nurse under PCP responsibility 	Disease diagnosis and management
Cleveringa <i>et al.</i> 2010 ⁷²	Journal article	Private sector-commercial	<ul style="list-style-type: none"> • Same as Cleveringa <i>et al.</i> 2008 but a cost effectiveness study. 	Disease diagnosis and management
Cortez 2014 ⁶⁶	PhD Thesis	Not stated	<ul style="list-style-type: none"> • A standalone computer-based CDSS: clinical decision support in the form of evidence-based drop-down boxes in the nursing electronic documentation system • Control (no drop-down boxes) 	[chronic] Disease management
Dalaba 2015 ⁸⁶	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer [laptop]-based CDSS: intervention software that prompts health workers to provide pre-natal and delivery care and then gives recommendations or alerts the health worker when there are danger signs • Usual care: pre-natal and delivery care without the use of CDSS 	Disease management/prevention reminders
Dowding <i>et al.</i> 2012 ⁹⁰	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS [intervention period]: an organization-wide electronic health record (EHR), Kaiser Permanent Health-Connect • Usual care (CDSS non-assisted)-pre-intervention period 	Disease prevention
Duclos <i>et al.</i> 2015 ⁸⁴	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: computerized system aimed to (1) detect automatically the malnourished patient by calculating Weight/Height and Height/Age ratio from the weight, height and age of children at admission and (2) automatically alert doctors and dietitians when the child was below the normal ratio. The alert was presented to doctors as a 'red flag' on the display alongside electronically prescribed drugs; for dietitians a monitoring dashboard was updated daily so they could intervene without waiting for a doctor. • Usual care (CDSS non-assisted) 	Disease diagnosis reminder
Dumont <i>et al.</i> 2012 ⁷⁸	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a computerised insulin-dosing calculator • paper protocol (modified Portland protocol) 	Drug dosing

TABLE 11 Additional information on studies and interventions (continued)

Author and year	Type of report	Funding source	Interventions and their details	CDSS function category
Dykes <i>et al.</i> 2010 ⁹¹	Journal article	Public sector/not for profit	<ul style="list-style-type: none"> • A standalone computer-based CDSS: <ul style="list-style-type: none"> ○ Phase 1: qualitative inquiry to identify barriers and facilitators to fall risk communication and interventions. ○ Phase 2: developing fall prevention tool kit prototype by using the Morse Falls Scale (MFS) risk factors ○ Phase 3: identify valid icons for the fall prevention tool kit using an iterative process involving domain experts, end users, and an illustrator. ○ Phase 4: implement the CDSS/conducting the trial • Usual care 	Disease prevention
Dykes <i>et al.</i> 2020 ⁹²	Journal article	Public sector/not for profit	<ul style="list-style-type: none"> • A standalone computer-based CDSS: A Five-Phase Fall Tailoring Interventions for Patient Safety (TIPS) tool kit <ul style="list-style-type: none"> ○ Phase 1: Problem analysis: learn about the needs and preferences of patients and providers and other social-technical factors that relate to fall prevention ○ Phase 2-3: Design and development: implement content, display, and workflow integration strategies most likely to address requirements and overcome barriers ○ Phase 4: Implementation: conduct a pilot test of fall TIPS and compare for effectiveness in engaging patients and families in the 3-step fall prevention process ○ Phase 5: Evaluation: evaluate the toolkit's efficacy on patient activation, falls, and injurious falls • Usual care 	Disease prevention
Fitzmaurice <i>et al.</i> 2000 ⁶⁷	Journal article	Public sector/MRC	<ul style="list-style-type: none"> • A standalone computer-based CDSS: anticoagulation management system used by Nurses • Usual care: anticoagulation management by physicians without the use of CDSS 	Disease management
Forberg <i>et al.</i> 2016 ⁸⁵	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a reminders system for peripheral venous catheter (PVC) intravenous treatment/management • Usual care (CDSS non-assisted) 	Disease management reminders
Fossum <i>et al.</i> 2011 ⁹³	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a system composed of CDSS integrated into the EHR based on two research-based risk assessment instruments: the Risk Assessment Pressure Scale (RAPS) for pressure ulcer risk screening and the Mini Nutritional Assessment (MNA®) tool for screening nutritional status. • Usual care (CDSS non-assisted) 	Disease prevention
Geurts <i>et al.</i> 2017 ⁷³	Journal article	Public sector/non-for-profit	<ul style="list-style-type: none"> • A standalone computer-based CDSS: clinical dehydration scale and guidelines on treatment of acute gastroenteritis were incorporated in an electronic, easily accessible clinical decision support system, available at each desktop at the emergency department • Usual care 	Disease diagnosis and management
Hovorka <i>et al.</i> 2007 ⁷⁹	Journal article	Public sector/EC	<ul style="list-style-type: none"> • A standalone computer-based CDSS: enhanced model predictive control (eMPC) that adapts itself to the input-output relationship observed during tight glucose control. • Usual care (CDSS non-assisted) 	Drug dosing

continued

TABLE 11 Additional information on studies and interventions (continued)

Author and year	Type of report	Funding source	Interventions and their details	CDSS function category
Kroth <i>et al.</i> 2006 ⁶⁸	Journal article	Not stated	<ul style="list-style-type: none"> • A standalone computer-based CDSS: consists of a DatascopeAccutor Plus patient monitor (Datascope Corp., Paramus, NJ) and a bed side PC (keyboard & LCD screen). The DataScope delivers its results (Blood pressure, pulse, body temperature and Oximetry) to the PC via an RS-232 serial interface that uses a vendor developed, data-exchange protocol. • Usual care (CDSS non-assisted) 	Disease management
Lattimer <i>et al.</i> 1998 ⁶⁴	Journal article	Public sector/EC	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a computer based telephone advice system (TAS) used by nurses in primary care call management • Usual care (Doctors with no CDSS) 	Triaging
Lattimer <i>et al.</i> 2000 ⁶⁵	Journal article	Public sector/MRC	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a computer based telephone advice system (TAS) used by nurses in primary care call management • Usual care (doctors with no CDSS) 	Triaging
Lee <i>et al.</i> 2009 ⁷⁴	Journal article	Public sector	<ul style="list-style-type: none"> • A hand-held- or mobile-based CDSS: a personal digital assistant-based clinical decision support system for obesity used for the screening and management of obesity, smoking, and depression and consists of 3 parts: screening, automated generation of diagnosis, and care planning • Usual care (CDSS non-assisted): personal digital assistant-based clinical log without decision support features for obesity 	Disease diagnosis and management
Lv <i>et al.</i> 2019 ⁹⁴	Journal article	Public sector/non-for-profit	<ul style="list-style-type: none"> • A standalone computer-based CDSS: the software included the following basic modules: medication reminder, adherence management, alert of acute asthma exacerbations, assessment of exacerbation severity, treatment recommendation, keeping a health diary, instant communication with healthcare providers and health education • Usual care 	Disease prevention
Mann <i>et al.</i> 2011 ⁸⁰	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: an EndoTool® insulin dosing CDSS package for acute glycaemic control • Usual care (paper-based protocol) 	Drug dosing
McDonald <i>et al.</i> 2017 ⁸⁸	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: with three components: (i) an algorithm to identify patients with complex medication regimens at increased risk of medication problems or adverse outcomes; (ii) a clinical alert comprising an email to the nurse's mobile device flagging specific patients with complex medication regimens and signposting to a 'medication regimen complexity care management module'; (iii) a complex medication management module as part of the Patient Care Record System (PCRS) that suggested nursing goals and interventions for patients with multiple co-morbidities and complex medications. • Usual care: nurses with pen-based tablet computers running the PCRS electronic health record who could access referral, medication, care plan, assessment information. Using clinical judgment alone, they decided issues to communicate to doctors, PCRS problems to 'pull down' and their priority. Before and/or seeing a patient nurses reviewed their Plan of Care, reviews and updates current medications and documents progress on the PCRS sub modules/problems. 	Disease diagnosis and management; and, diagnosis alerts

TABLE 11 Additional information on studies and interventions (continued)

Author and year	Type of report	Funding source	Interventions and their details	CDSS function category
Paulsen <i>et al.</i> 2020 ⁹⁵	Journal article		<ul style="list-style-type: none"> • A standalone computer-based CDSS: MyFood digital decision support system is developed for use among hospitalized patients who are malnourished or at risk of malnutrition • Usual care 	Disease prevention
Plank <i>et al.</i> 2006 ⁸¹	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: a model predictive control system used for glycaemic control. It enables the prediction of the glucose excursion by a dose optimizer. The dose optimizer proposes future insulin infusion rates and tunes the rates until the predicted glucose excursion fits into a desired glucose excursion • Usual care (Routine Treatment Protocol (RTP)): blood glucose values were provided to the ICU staff as required by the routine glucose management protocol implemented in the respective ICU 	Drug dosing
Rood <i>et al.</i> 2005 ⁶⁹	Journal article	Not stated	<ul style="list-style-type: none"> • A standalone computer-based CDSS: guideline-based advice provided via a clinical information system (CIS) decision support software module (Event Manager) and a custom-made Visual Basic application integrated within the CIS. The application displayed glucose and insulin data and suggested current treatment and the interval to the next glucose measurement. • Usual care (paper-based guideline) 	Disease management
Roukema <i>et al.</i> 2008 ⁸⁷	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: nurses used a computer system that automatically identified children with high risk score and recommends laboratory test • Usual care: Physicians assess patients without the help of CDSS 	Diagnosis/management reminders
Sassen <i>et al.</i> 2014 ⁷⁰	Journal article	Not stated	<ul style="list-style-type: none"> • Web-based CDSS: clinical decision support system used to optimize shared decision-making and the self-management of patients • Control (no CDSS use) 	Disease management
Snooks <i>et al.</i> 2014 ⁶²	Journal article	Public sector	<ul style="list-style-type: none"> • A hand-held tablet [computer]-based CDSS: paramedics use CDSS to decide whether to take patients who had fallen to an Emergency Department or leave them at home with referral to a community-based falls service • Usual care: paper-based protocols comprised assessment, treatment on scene as required and default conveyance to the Emergency Department unless the patient refused to travel to hospital. 	Triaging
Vadher <i>et al.</i> 1997 ⁸²	Journal article	Public sector	<ul style="list-style-type: none"> • A standalone computer-based CDSS: nurses used oral anticoagulant control model to monitor doses • Usual care (trainee doctor without CDSS) 	Drug dosing
Wells 2013 ⁶³	PhD Thesis	Not stated	<ul style="list-style-type: none"> • A hand-held tablet [computer]-based CDSS: paramedics use CDSS to decide whether to take patients who had fallen to an emergency department or leave them at home with referral to a community-based falls service • Usual care: paper-based protocols comprised assessment, treatment on scene as required and default conveyance to the Emergency Department unless the patient refused to travel to hospital. 	Triaging

CDSS, computerised decision support system; EC, European Commission; ICU, intensive care unit; MRC, Medical research Council.

61. Bennett P, Hardiker N. A Quantitative Study Investigating the Effects of Computerised Clinical Decision Support in the Emergency Department. *Studies in health technology and informatics* 2016;225:53–7.
62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;9(9):e106436. doi: <https://dx.doi.org/10.1371/journal.pone.0106436>.
63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a Prehospital Setting: Processes of Adoption and Impact on Paramedic Role and Practice. Swansea University (United Kingdom), 2013.
64. Lattimer V, George S, Thompson F, *et al.* Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;317(7165):1054–9.
65. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;320(7241):1053–7.
66. Cortez S. Measuring active clinical decision support influence on nursing research utilization. University of Phoenix, 2014.
67. Fitzmaurice DA, Hobbs FD, Murray ET, *et al.* Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;160(15):2343–8.
68. Kroth PJ, Dexter PR, Overhage JM, *et al.* A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *Amia Annual symposium proceedings AMIA symposium* 2006:444–8.
69. Rood E, Bosman RJ, van der Spoel JI, *et al.* Use of a Computerized Guideline for Glucose Regulation in the Intensive Care Unit Improved Both Guideline Adherence and Glucose Regulation. *Journal of the American Medical Informatics Association* 2005;12(2):172–80. doi: 10.1197/jamia.M1598.
70. Sassen B, Kok G, Schepers J, *et al.* Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;16(10):e211.
71. Cleveringa FGW, Gorter KJ, Van Donk MD, *et al.* Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;31(12):2273–5.
72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;33(2):258–63.
73. Geurts D, de Vos-Kerkhof E, Polinder S, *et al.* Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;176(2):173–81.
74. Lee NJ, Chen ES, Currie LM, *et al.* The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;32(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.
75. Blaha J, Kopecky P, Matias M, *et al.* Comparison of three protocols for tight glycemic control in cardiac surgery patients. *Diabetes Care* 2009;32(5):757. doi: 10.2337/dc08-1851.
76. Canbolat O, Kapucu S, Kilickaya O. Comparison of routine and computer-guided glucose management for glycemic control in critically ill patients. *Critical Care Nurse* 2019;39(4):20–7. doi: 10.4037/ccn2019431.
77. Cavalcanti AB, Silva E, Pereira AJ, *et al.* A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;24(3):371–8. doi: <https://doi.org/10.1016/j.jccr.2009.05.005>.
78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;21(2):106–15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.
79. Hovorka R, Kremen J, Blaha J, *et al.* Blood glucose control by a model predictive control algorithm with variable sampling rate versus a routine glucose management protocol in cardiac surgery patients: a randomized controlled trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;92(8):2960–4. doi: 10.1210/jc.2007-0434.
80. Mann EA, Jones JA, Wolf SE, *et al.* Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;32(2):246–55.
81. Plank J, Blaha J, Cordingley J, *et al.* Multicentric, randomized, controlled trial to evaluate blood glucose control by the model predictive control algorithm versus routine glucose management protocols in intensive care unit patients. *Diabetes Care* 2006;29(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical and Laboratory Haematology* 1997;19(3):203–07.
83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
84. Duclos A, Touzet S, Restier L, *et al.* Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *European journal of clinical nutrition* 2015;69(7):769–75. doi: 10.1038/ejcn.2014.288.

85. Forberg U, Unbeck M, Wallin L, *et al.* Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implementation Science* 2016;11 doi: 10.1186/s13012-016-0375-9.
86. Dalaba MA, Akweongo P, Aborigo RA, *et al.* Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;10(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, *et al.* Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *Journal of the American Medical Informatics Association: JAMIA* 2008;15(1):107–13. doi: 10.1197/jamia.M2164.
88. McDonald MV, Feldman PH, Barron-Vaya Y, *et al.* Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;22(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, *et al.* A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;50(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;19(4):615–20. doi: 10.1136/amiajnl-2011-000504 [published Online First: 2011/12/17].
91. Dykes PC, Carroll DL, Hurley A, *et al.* Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;304(17):1912–18.
92. Dykes PC, Burns Z, Adelman J, *et al.* Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA network open* 2020;3(11):e2025889-e89.
93. Fossum M, Alexander GL, Ehnfors M, *et al.* Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;80(9):607–17. doi: <https://dx.doi.org/10.1016/j.ijmedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, *et al.* A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of Advanced Nursing* 2019;75(11):3058–67.
95. Paulsen MM, Paur I, Gjestland J, *et al.* Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. *Clinical Nutrition* 2020;39(12):3607–17.(Table 12).

Appendix 3 Study information on protocols and CDSS implementation theory/models

TABLE 12 Study information on protocols and CDSS implementation theory/models

Author and year	Implementation model/theory for CDSS	Protocol published ^a	Science of implementation published elsewhere ^a
Beeckman <i>et al.</i> 2013 ⁸⁹	Implementation model by Grol and Wensing (2005): a stepped approach to implementation, justifying the implementation need to reduce target group possible resistance. Starting with in-depth analysis of current practice, target group, and context a series of steps employed (1) matching research findings and/or existing guidelines to the relevant practice, (2) describing desired change outcomes, (3) selecting/developing implementation strategies, (4) developing and executing implementation processes, and (5) evaluation and adaptation of processes.	None	None
Bennet <i>et al.</i> 2016 ⁶¹	Although author discusses 'theory of decision-making' in nursing (e.g: four stages of medical decision-making by Elstein <i>et al.</i> (1978), page 44 of the thesis), it was not explicitly described that if implementation was based on any model or theory.	None	None
Blaha <i>et al.</i> 2009 ⁷⁵	None described	Yes (Trial ID: NCT00764712); protocol outcomes and manuscript outcomes match. However, the protocol does not provide any additional information.	None
Byrne 2005 ⁸³	Author discusses Rycroft-Malone <i>et al.</i> 's conceptual framework for adoption (page 43 of Thesis) but not explicitly mentioned that it was used as a model of implementation.	None	None
Canbolat <i>et al.</i> 2019 ⁷⁶	None described	None	None
Cavalcanti <i>et al.</i> 2009 ⁷⁷	None described	None	None
Cleveringa <i>et al.</i> 2008 ⁷¹	None described	Yes (Trial ID: ISRCTN21523044); protocol outcomes and manuscript outcomes match. However, the protocol does not provide any additional information.	None
Cleveringa <i>et al.</i> 2010 ⁷²	None described	Yes (Trial ID: ISRCTN21523044); Cost effectiveness outcomes in manuscript are not listed in protocol. The protocol does not provide any additional information.	None

continued

TABLE 12 Study information on protocols and CDSS implementation theory/models (continued)

Author and year	Implementation model/theory for CDSS	Protocol published ^a	Science of implementation published elsewhere ^a
Cortez 2014 ⁶⁶	None described	None	None
Dalaba 2015 ⁸⁶	None described	None	None
Dowding <i>et al.</i> 2012 ⁹⁰	None described	None	None
Duclos <i>et al.</i> 2015 ⁸⁴	None described	None	None
Dumont <i>et al.</i> 2012 ⁷⁸	None described	None	None
Dykes <i>et al.</i> 2010 ⁹¹	None described	Yes (Trial ID: NCT00675935); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None
Dykes <i>et al.</i> 2020 ⁹²	None described	Yes (Trial ID: NCT02969343); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None
Fitzmaurice <i>et al.</i> 2000 ⁶⁷	None described	None	None
Forberg <i>et al.</i> 2016 ⁸⁵	None described	Yes (Trial ID: ISRCTN44819426); manuscript outcomes match the listed secondary outcomes in the protocol. However, the protocol does not provide any additional information.	None
Fossum <i>et al.</i> 2011 ⁹³	None described	None	None
Geurts <i>et al.</i> 2017 ⁷³	None described	Yes (Trial ID: NTR2304); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None
Hovorka <i>et al.</i> 2007 ⁷⁹	None described	None	None
Kroth <i>et al.</i> 2006 ⁶⁸	None described	None	None
Lattimer <i>et al.</i> 1998 ⁶⁴	None described	None	None
Lattimer <i>et al.</i> 2000 ⁶⁵	None described	None	None
Lee <i>et al.</i> 2009 ⁷⁴	None described	None	None

TABLE 12 Study information on protocols and CDSS implementation theory/models (continued)

Author and year	Implementation model/theory for CDSS	Protocol published ^a	Science of implementation published elsewhere ^a
Ly <i>et al.</i> 2019 ⁹⁴	None described	Authors cite protocol number (ChiCTR1800016726) but not accessible.	None
Mann <i>et al.</i> 2011 ⁸⁰	None described	None	None
McDonald <i>et al.</i> 2017 ⁸⁸	None described	None	The authors published about 'Automating the medication regimen complexity index' (https://doi.org/10.1136/amia-jnl-2012-001272) a year before but hardly give any theoretical background or models for adoption other than just 'testing' if the technology works.
Paulsen <i>et al.</i> 2020 ⁹⁵	None described	Yes (Trial ID: NCT03412695); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	The authors published about 'Automating the medication regimen complexity index' (https://mhealth.jmir.org/2018/9/e175/) but hardly give any theoretical background or models for adoption other than just 'testing' if the technology works.
Plank <i>et al.</i> 2006 ⁸¹	None described	None	None
Rood <i>et al.</i> 2005 ⁶⁹	None described	None	None
Roukema <i>et al.</i> 2008 ⁸⁷	None described	None	None
Sassen <i>et al.</i> 2014 ⁷⁰	None described	Yes (Trial ID: NTR2584); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None
Snooks <i>et al.</i> 2014 ⁶²	None described	Yes (Trial ID: ISRCTN10538608); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None
Vadher <i>et al.</i> 1997 ⁸²	None described	None	None

Author and year	Implementation model/theory for CDSS	Protocol published ^a	Science of implementation published elsewhere ^a
Wells 2013 ⁶³	None described	Yes (Trial ID: ISRCTN10538608); manuscript outcomes match the listed outcomes in the protocol. However, the protocol does not provide any additional information.	None

a searches conducted using first author and title of project/manuscript.

61 Bennett P, Hardiker N. A quantitative study investigating the effects of computerised clinical decision support in the emergency department. *Studies in health technology and informatics* 2016;**225**:53–7.

62. Snooks HA, Carter B, Dale J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1): cluster randomised trial of computerised clinical decision support for paramedics. *PLoS ONE* 2014;**9**(9):e106436. doi: <https://dx.doi.org/10.1371/journal.pone.0106436>.

63. Wells B. Implementation of Computerised Clinical Decision Support (CCDS) in a prehospital setting: processes of adoption and impact on paramedic role and practice. Swansea University (United Kingdom), 2013.

64. Lattimer V, George S, Thompson F, *et al.* Safety and effectiveness of nurse telephone consultation in out of hours primary care: Randomised controlled trial. *British Medical Journal* 1998;**317**(7165):1054–9.

65. Lattimer V, Sassi F, George S, *et al.* Cost analysis of nurse telephone consultation in out of hours primary care: evidence from a randomised controlled trial. *BMJ* 2000;**320**(7241):1053–7.

66. Cortez S. Measuring active clinical decision support influence on nursing research utilization. University of Phoenix, 2014.

67. Fitzmaurice DA, Hobbs FD, Murray ET, *et al.* Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: a randomized, controlled trial. *Archives of Internal Medicine* 2000;**160**(15):2343–8.

68. Kroth PJ, Dexter PR, Overhage JM, *et al.* A computerized decision support system improves the accuracy of temperature capture from nursing personnel at the bedside. *Amia Annual symposium proceedings AMIA symposium* 2006:444–8.

69. Rood E, Bosman RJ, van der Spoel JI, *et al.* Use of a Computerized guideline for glucose regulation in the intensive care unit improved both guideline adherence and glucose regulation. *Journal of the American Medical Informatics Association* 2005;**12**(2):172–80. doi: 10.1197/jamia.M1598.

70. Sassen B, Kok G, Schepers J, *et al.* Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. *Journal of medical Internet research* 2014;**16**(10):e211.

71. Cleveringa FGW, Gorter KJ, Van Donk MD, *et al.* Combined task delegation, computerized decision support, and feedback improve cardiovascular risk for type 2 diabetic patients. *Diabetes Care* 2008;**31**(12):2273–5.

72. Cleveringa FGW, Welsing PMJ, Van Den Donk M, *et al.* Cost-effectiveness of the diabetes care protocol, a multifaceted computerized decision support diabetes management intervention that reduces cardiovascular risk. *Diabetes Care* 2010;**33**(2):258–63.

73. Geurts D, de Vos-Kerkhof E, Polinder S, *et al.* Implementation of clinical decision support in young children with acute gastroenteritis: a randomized controlled trial at the emergency department. *European Journal of Pediatrics* 2017;**176**(2):173–81.

74. Lee NJ, Chen ES, Currie LM, *et al.* The effect of a mobile clinical decision support system on the diagnosis of obesity and overweight in acute and primary care encounters. *ANS Adv Nurs Sci* 2009;**32**(3):211–21. doi: <https://dx.doi.org/10.1097/ANS.0b013e3181b0d6bf>.

75. Blaha J, Kopecky P, Matias M, *et al.* Comparison of three protocols for tight glycemic control in cardiac surgery patients. *Diabetes Care* 2009;**32**(5):757. doi: 10.2337/dc08-1851.

76. Canbolat O, Kapucu S, Kilickaya O. Comparison of routine and computer-guided glucose management for glycemic control in critically ill patients. *Critical Care Nurse* 2019;**39**(4):20–27. doi: 10.4037/ccn2019431.

77. Cavalcanti AB, Silva E, Pereira AJ, *et al.* A randomized controlled trial comparing a computer-assisted insulin infusion protocol with a strict and a conventional protocol for glucose control in critically ill patients. *Journal of Critical Care* 2009;**24**(3):371–8. doi: <https://doi.org/10.1016/j.jcrrc.2009.05.005>.

78. Dumont C, Bourguignon C. Effect of a computerized insulin dose calculator on the process of glycemic control. *Am J Crit Care* 2012;**21**(2):106–15. doi: <https://dx.doi.org/10.4037/ajcc2012956>.

79. Hovorka R, Kremen J, Blaha J, *et al.* Blood glucose control by a model predictive control algorithm with variable sampling rate versus a routine glucose management protocol in cardiac surgery patients: a randomized controlled trial. *The Journal of Clinical Endocrinology & Metabolism* 2007;**92**(8):2960–4. doi: 10.1210/jc.2007-0434.

80. Mann EA, Jones JA, Wolf SE, *et al.* Computer decision support software safely improves glycemic control in the burn intensive care unit: A randomized controlled clinical study. *Journal of Burn Care and Research* 2011;**32**(2):246–55.
81. Plank J, Blaha J, Cordingley J, *et al.* Multicentric, randomized, controlled trial to evaluate blood glucose control by the model predictive control algorithm versus routine glucose management protocols in intensive care unit patients. *Diabetes Care* 2006;**29**(2):271. doi: 10.2337/diacare.29.02.06.dc05-1689.
82. Vadher BD, Patterson DL, Leaning M. Comparison of oral anticoagulant control by a nurse-practitioner using a computer decision-support system with that by clinicians. *Clinical and laboratory haematology* 1997;**19**(3):203-07.
83. Byrne CM. Impact of prospective computerized clinical decision support information and targeted assistance on nursing home resident outcomes. State University of New York at Albany, 2005.
84. Duclos A, Touzet S, Restier L, *et al.* Implementation of a computerized system in pediatric wards to improve nutritional care: a cluster randomized trial. *European journal of clinical nutrition* 2015;**69**(7):769–75. doi: 10.1038/ejcn.2014.288.
85. Forberg U, Unbeck M, Wallin L, *et al.* Effects of computer reminders on complications of peripheral venous catheters and nurses' adherence to a guideline in paediatric care—a cluster randomised study. *Implementation Science* 2016;**11** doi: 10.1186/s13012-016-0375-9.
86. Dalaba MA, Akweongo P, Aborigo RA, *et al.* Cost-effectiveness of clinical decision support system in improving maternal health care in Ghana. *PLoS ONE* 2015;**10**(5):e0125920. doi: <https://dx.doi.org/10.1371/journal.pone.0125920>.
87. Roukema J, Steyerberg EW, van der Lei J, *et al.* Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source. *Journal of the american medical informatics association: JAMIA* 2008;**15**(1):107–13. doi: 10.1197/jamia.M2164.
88. McDonald MV, Feldman PH, Barron-Vaya Y, *et al.* Outcomes of clinical decision support (CDS) and correlates of CDS use for home care patients with high medication regimen complexity: a randomized trial. *Journal of Evaluation in Clinical Practice* 2017;**22**(1):10–19.
89. Beeckman D, Clays E, Van Hecke A, *et al.* A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: a two-armed randomized controlled trial. *Int J Nurs Stud* 2013;**50**(4):475–86. doi: <https://dx.doi.org/10.1016/j.ijnurstu.2012.09.007>.
90. Dowding DW, Turley M, Garrido T. The impact of an electronic health record on nurse sensitive patient outcomes: an interrupted time series analysis. *J Am Med Inform Assoc* 2012;**19**(4):615–20. doi: 10.1136/amiainl-2011-000504 [published Online First: 2011/12/17].
91. Dykes PC, Carroll DL, Hurley A, *et al.* Fall prevention in acute care hospitals: a randomized trial. *Jama* 2010;**304**(17):1912–18.
92. Dykes PC, Burns Z, Adelman J, *et al.* Evaluation of a patient-centered fall-prevention tool kit to reduce falls and injuries: a nonrandomized controlled trial. *JAMA network open* 2020;**3**(11):e2025889-e89.
93. Fossum M, Alexander GL, Ehnfors M, *et al.* Effects of a computerized decision support system on pressure ulcers and malnutrition in nursing homes for the elderly. *Int J Med Inf* 2011;**80**(9):607–17. doi: <https://dx.doi.org/10.1016/j.ijmedinf.2011.06.009>.
94. Lv S, Ye X, Wang Z, *et al.* A randomized controlled trial of a mobile application-assisted nurse-led model used to improve treatment outcomes in children with asthma. *Journal of advanced nursing* 2019;**75**(11):3058–67.
95. Paulsen MM, Paur I, Gjestland J, *et al.* Effects of using the MyFood decision support system on hospitalized patients' nutritional status and treatment: A randomized controlled trial. *Clinical Nutrition* 2020;**39**(12):3607–17.

